

Prospects of Plant Tissue Culture in Orchid Propagation: A Review

Nand Lal¹, Madhulika Singh²

How to cite this article:

Nand Lal, Madhulika Singh. Prospects of Plant Tissue Culture in Orchid Propagation: A Review. Indian J Biol 2020; 7(2):103-110.

Abstract

Orchids (Family Orchidaceae) are ornamentally valuable and unique group of flowering plants. They are valued mostly for their long lasting and diverse morphology of flowers. Some orchids are highly prized due to their application in food, flavor and medicinal industry. Seed germination and vegetative propagation are the methods through which orchid plants regenerate in nature. Both these methods of regeneration are quite slow to fulfill orchid demand. Plant tissue culture technique has been used extensively in multiplication of orchids. Although seed germination produces heterogenous population but most of orchid *species* are multiplied by asymbiotic seed germination. Propagation through protocorm-like-bodies (PLBs) or direct from explant regeneration is quite more advantageous to multiply orchids. Artificial seed technology is another method used for orchid multiplication, transport and conservation of endangered orchid *species*. Culture of medicinal orchids for production of bioactive metabolites could be used as substitute for dependence on natural population.

Keywords: Orchids; Propagation; Protocorm like bodies; Thidiazuron; Secondary metabolite; Synthetic seed.

Introduction

Orchids (Family Orchidaceae) are generally characterized with long lasting, diverse morphology and exquisite flowers that have been extensively used in floriculture since time immemorial. At present, in world's flower industry it account with an annual sales more than US\$4 billion.⁶⁹ The family includes about 736 genera with more than 28,000 *species* which are mostly distributed in the wet tropics worldwide except few isolated islands, polar and desert regions.^{14,69} India has rich orchid diversity and approximately 158 genera and 1331 *species* flourishing up to an elevation of 5000m are recorded.⁵⁴ In comparison to plants from other families the mycorrhizal specificity, pollinator specialization and very low germination of seeds restricts the distribution of orchid *species* narrowly and are quite more susceptible to habitat distribution.⁷⁰

Author's Affiliation: ¹Professor, Department of Life Sciences, ²Assistant Professor, Department of Biotechnology, Chhatrapati Shahu Ji Maharaj University, Kanpur, Uttar Pradesh 208012, India.

Corresponding Author: Nand Lal, Professor, Department of Life Sciences, Chhatrapati Shahu Ji Maharaj University, Kanpur, Uttar Pradesh 208012, India.

E-mail: nl_pr@yahoo.co.in

The family is considered as most diverse family among the flowering plants.¹⁵ They grow on all kinds of habitats, except the aquatic systems and on their growth habitat they may be characterized as terrestrial, epiphytic or saprophytic. Most of the orchids are epiphytic in nature, while the rest are terrestrial. Although orchids are grown mainly for ornamental applications but, some are employed as herbal medicines and food (tubers of *Cynorchis* and *Eulophia*) by many different cultures and tribes.⁴ Vanillin obtained from *Vanilla planifolia*, an

odourous constituent is a very useful commercial product. It is used for flavouring in baking, ice-creams, custards, pudding and in aromatherapy and perfume industry.³⁵ *Gastrodia elata* has medicinal application in treating tetanus, headaches, dizziness and epilepsy.⁶² *Macodes*, *Ludisia*, *Anoectochilus* and *Goodyera* genera have attractive foliage and are known as jewel orchids. The wild population of orchids can multiply sexually by seed, and asexually by vegetative propagation. However, the vegetative propagation is very slow and only few seeds germinate in spite millions seed produced in each capsule. The seeds have simple seed coat structure and reduced embryos. The embryos are minute size generally varies in length from 0.05 to 6 mm, width 0.01 to 0.93 mm, and 0.3 to 14 µg in weight.² The seeds have varied suspensor morphology, lack cotyledons, endosperm and depicts availability of insufficient nutrient reserves needed for germination. The association of orchid seeds with appropriate fungus for germination is necessary to fulfill the essential physiochemical stimulus needed for seed germination and seedling development. Collection of wild population as a source of stock plant for various socio-economic trades instead of their commercial multiplication is a serious threat to several *species* of orchids. Low multiplication of wild population and extensive uses in floral and other industries has resulted in depletion of population of many *species* at an alarming rate and some of them are at the verge of extinction.^{16,42} In recent past, in vitro culture methods of propagation have been used to reduce collection pressures on wild population. In addition to asymbiotic seed germination, orchids are regenerated through explants such as leaves, root rhizome, shoot meristem, axillary buds, pseudonode etc.¹⁶ In this communication, we have reviewed the recent progress made in plant tissue culture for multiplication, conservation and genetic modification of orchids.

1. Asymbiotic Seed germination

In nature orchid seed germination is dependent on association with mycorrhizal fungus. Knudson³² demonstrated that no mycorrhizal association is needed on in vitro germination of orchid seeds and by using simple nutrient medium along with sucrose Knudson pioneered the germination of several epiphytic orchids. Multiplication of orchids through in vitro seed germination technique have been developed for several *species* and are most commonly used method for multiplying different orchid *species* and hybrids representing diverse habits and habitats.^{3,49} Under culture conditions

germination of seeds are affected by several physico-chemical factors and it is requisite to optimize seed germination for each *species* of orchid. A number of media such as Murashige and Skoog,⁴³ Mitra et al.,⁴¹ Kundson³³ and Gamborg et al.,²¹ etc. have been used for seed germination and propagation through tissue culture. Growth regulators like auxins and cytokinins are added into medium for germination and growth of seedlings of several orchids.⁵⁴

The organic additives such as peptone, banana extract, coconut water, casein hydrolyzate, yeast extract etc. accelerate seed germination and protocorm formation.²⁹ Availability of natural source of carbohydrates, vitamins, amino acids, inorganic ions and phytohormones in such additives fulfill the nutritional requirement of germinating seeds which in wild are fulfilled by mycorrhizal association.^{25,54}

Both immature and mature seeds can be germinated using tissue culture technique but germination of mature seeds is quite more difficult in comparison to immature seeds. Germination of seeds generally depends on their nutrient reserves. Several variations in reserve food composition between immature and mature seeds have been reported by researchers. In *Corallorhiza maculata* the immature seeds have starch as reserve food while mature seeds contain lipids.²⁷ This was suggested that during the process of seed maturation starch and other carbohydrates of immature seeds is converted into lipids of mature seeds. The lack of metabolic machinery in mature seeds is another hindrance in germination of mature seeds.²⁶ Dormancy of mature seeds and accumulation of germination inhibitors in the seed coat also restrict germination of mature seeds. The dormancy of seeds can be broken by low temperature treatment²⁷ or by using sucrose.⁶⁶ In the germination process, seeds first form protocorm and protocorm proliferates to form more protocorm or can differentiate root and shoot.

2. Multiplication from vegetative explants

Plant tissue culture techniques have been used for regeneration of orchids during last several years and a few *species* of orchids have been multiplied either through direct shoot regeneration^{16,54,58} or from protocorm like bodies formation from culture of vegetative explants.^{16,54} Orchid propagation through seed germination give rise to heterozygous plants, while micropropagation through various vegetative parts of the plants can control this problem, however multiplication of orchids through micropropagation is limited due to obstacle such

as low population of plants in nature, phenolics exudation, field transplantation, somaclonal variation etc. Of the various reports regeneration from the explants selected from plants growing in nature or greenhouse is very few in comparison from the explants selected from in vitro raised seedlings (Table 1). Explants such as apical meristem, node, pseudonode, shoot, leaf, leaf tip, rhizome etc. have been used for clonal multiplication. Different cytokinins such as BAP (N6 -benzylaminopurine), Kinetin, 2-iP (2-isopentyladenine), TDZ (Thidiazuron), either alone or in combination with auxin 2,4-D (2,4 dichlorophenoxy acetic acid), NAA (α -naphthalene acetic acid), IAA have been tested for shoot proliferation and PLBs formation (Table 1). In *Oncidium flexuosum*, TDZ was reported to be effective in direct PLB formation from leaf explants under dark while plants with well developed shoots and roots grew only after transfer of PLBs to under light and growth regulator free medium.³⁸ Roy et al.,⁵¹ reported that low concentration (0.909 μ M) of TDZ induced more PLBs from seedling derived pseudostem segments of *Cymbidium giganteum* while in high concentration, PLB multiplication increased but reduced plantlet and root development from the PLBs. They also

suggested that in dual phase culture system where a thin layer of the liquid medium overlaid the semi-solid medium, low concentration of TDZ is most appropriate for inducing healthy plantlets from pseudostem segments of *Cymbidium giganteum*. In *Dendrobium longicornu* shoot formation from the nodal explants was observed in medium containing 15 μ M BAP and 5 μ M NAA while PLBs were developed when the medium contained BAP in combination with 2,4-D.²⁰ In some orchids light is also considered among the various factors that affect PLB induction. In *Phalaenopsis* absence of light induce PLB formation while differentiation of PLBs into plantlets was favored by light. However, most studies in *Phalaenopsis* describe presence of light for PLB regeneration. Induction of PLBs is also affected by the type of light. According to Mehraj et al.³⁹ with the use of blue and white light emitting diodes (LED) combined with trehalose as carbohydrate source or red and white LED with sucrose as carbohydrate is favorable for PLB regeneration. Hybridization is very useful technique for producing orchid cultivar with attractive morphology, color and fragrance of flowers. Mass multiplication of orchid hybrids can be made possible certainly by plant tissue culture technique.¹²

Table 1: Micropropagation through vegetative explants culture of some orchids.

Orchid	Explant	Medium and Growth Regulator(s)	Result/Response	Reference
<i>Arundina graminifolia</i>	Node explants derived from three-year-old field grown plants	$\frac{1}{2}$ MS + 6.97 μ M kinetin (Kn), or 15% coconut water (CW) or 13.3 μ M BA 1/2MS + 44.4 μ M BA	sprouting of the axillary bud PLB from base of bud	(36)
Cattleya 'Almakee'	epidermal (=foliar) peel root explants from the greenhouse plants	Mitra with Kn (1mg L-1) + NAA (1 mgL-1)	PLB and somatic embryogenesis	(53)
<i>Coelogyne cristata</i>	segment of PLBs (3 mm ² in size)	$\frac{1}{2}$ MS with 1.0 mg l-1 NAA and 0.5 mg l-1 BA with Coconut Powder	Shoot formation	(44)
<i>Cymbidium giganteum</i>	Pseudostem from seedling	MS with 0.909 μ M TDZ	PLB formation	[51]
<i>Cymbidium goeringii</i>	Rhizome explants from seedlings	MS with 20 μ M 2,4-D and 2 μ M TDZ MS with 2 μ M NAA	Shoot formation Rooting of shoots	(46)
<i>Dendrobium</i> sp.	Leaf tip	MS + 0.5 mg l-1 NAA + 0.5 mg l-1 BAP	Shoot initiation and plant formation	(24)
<i>Dendrobium</i> sp.	Leaf tip	MS + 10 mg l-1 2, 4-D	PLB	(24)
<i>Dendrobium lituiflorum</i>	Axillary buds from 5-month-old in vitro grown plants	MS medium supplemented with 0.5 mg l-1 2,4-D	PLB	(17)

<i>Doritaenopsis</i>	Root tip from in vitro plantlets	modified MS +2.3 μ M TDZ	PLB	(47)
<i>Dendrobium chrysotoxum</i>	Transverse sections (2 mm thickness) of stem-nodes from in vitro raised seedlings	semisolid and liquid Murashige and Skoog medium supplemented with BA 4.44 μ M and KIN 4.65 μ M	Shoot buds on cytokinin containing media or (PLBs) on NAA containing media both solid and liquid.	(31)
<i>Geodorum densiflorum</i>	Rhizome tips from in vitro seedlings	MS + 2 mg/l BAP	Shoot formation	(5)
<i>Rhynchosstylis retusa</i>	Leaf	MS with 1.5 mg/l BAP and 1.5 mg/l Kinetin	PLB and somatic embryogenesis	(28)
<i>Oncidium flexuosum</i>	Leaf	modified MS medium with 1.5 μ M TDZ under dark	PLB formation	(38)
<i>Phalaenopsis amabilis</i> cv. 'Cool Breeze'	thin sections of Inflorescence axis	NDM (New Dogashima Medium) medium 0,5 + mg/l-1 NAA, 2% sucrose, 10% coconut water, 2 gl-1 peptone and 1gl-1 activated charcoal	PLB formation	(60)
<i>Thunia marshalliana</i>	Pseudonodes from plants grown in glasshouse	MS with 22.0 μ M BAP	Shoot regeneration	(58)
<i>Vanilla planifolia</i>	Leaf from field growing plant	MS medium + 4.52 μ M 2,4-D + 2.22 μ M BAP	Shoot regeneration from callus	(30)
<i>Vanilla planifolia</i>	Shoot tip from plants growing in greenhouse	MS with 1.0mg/l BAP + 150ml coconut water	Shoot regeneration	(34)

Table 2: Production of Secondary metabolite from in vitro cultures of some medicinal orchids.

Orchid	Culture type	Secondary metabolite/activity	Reference
<i>Ansellia africana</i>	In vitro regenerated PLBs	phenolic acids exhibiting a high degree of antioxidant activity	(7)
<i>Coelogyne ovalis</i>	Leaves of micropropagated plantlets	phenolics, flavonoids and anthocyanins) and antioxidant activity	(59)
<i>Dendrobium thyrsiflorum</i>	Shoot	Antioxidant activity	(6)
<i>Dendrobium aphyllum</i>	In vitro plantlets	Antioxidant activity	(9)
<i>Dendrobium fimbriatum</i>	In vitro regenerated shoot, root and leaves	β -sitosterol	(48)
Dendrobium hybrid(Dendrobium enopi x Dendrobium Pink Lady)	LED illuminated Protocorm-Like Bodies	Flavonoid, Phenolics and anticancer activity	(67)
<i>Habenaria edgeworthii</i>	callus suspension cultures	Phenolics and antioxidant activity	(22)
<i>Malaxis acuminata</i>	In vitro regenerated leaf and shoot	Dietary fatty acids, α -hydroxy acids, phenolic acids, sterols, amino acids, sugars and glycosides	(10)

3. Production of Synthetic seed and in vitro conservation

Artificial seed technology is considered as an effective and extensively used method for propagation of plant species exhibiting reproductive barrier.^{1,37,57} The difficulty in propagation of orchids through seeds can be replaced by using artificial seed technology. Transportation and delivery of propagules can be easily carried out through artificial seeds. Several orchid species

such as *Cymbidium devonianum*,¹⁸ *Cymbidium eburneum*, *Cymbidium hookerianum*,²³ *Geodorum densiflorum*,¹⁹ *Paphiopedilum wardii*,⁶⁸ *Dendrobium wardianum*,⁵² *Ansellia africana*⁸ etc. have been mass propagated successfully through artificial seed. Its application in conservation of desirable, rare and endangered orchids has been reported by several researchers. Plantlets of *Vanda coerulea* was obtained by encapsulation of six month old PLBs regenerated from leaf explants, the encapsulated PLBs was stored for 100 days at 4°C.⁵⁵ Artificial

seeds of *Cymbidium aliofolium* was prepared by encapsulating protocorms with 4% (w/v) sodium alginate and 0.2 mol/l calcium chloride solution showed 100% germination and also the encapsulated protocorm could be stored for 28 days at 4°C.⁵⁰ Reducing the nutrient strength in the encapsulated matrix and low temperature improves storage period of PLBs of *Cymbidium devonianum*.¹⁸ In *Dendrobium nobile* incorporation of sucrose and mannitol (7.5 and 12 %) in the encapsulating matrix showed storage of encapsulated PLBs till 60 days.⁴²

4. In vitro culture and secondary metabolite production

In addition to extensive use of orchids as ornamentals many orchids are used extensively as herbal medicines. A number of reports are available on medicinal uses of orchids. Some orchids such as *Dendrobium nobile*, *Gastrodia elata* and *Bletilla striata* are routinely used in Traditional Chinese Medicine.^{45,56} The genus *Anoctochilu*, *Calanthe*, *Coelogyne*, *Cymbidium*, *Cypripedium*, *Dendrobium*, *Ephemerantha*, *Gymnadenia*, *Habenaria*, *Nevilia* and *Thunia* are common orchid genera which exhibit biosynthesis and accumulation of medicinally useful metabolites.⁴⁵ Phytochemicals such as alkaloids, bibenzyl derivatives, flavonoids, phenanthrenes etc. have been reported from orchids.⁵⁶ The populations of medicinal orchids are decreasing at alarming rate due to their exploitation in medicinal uses. The in vitro culture of plant cells, tissues and organs has been demonstrated as a pivotal source for production of useful structurally complex and high value natural biomolecules.⁶⁴ The use of plant tissue culture is a substitute for dependence on natural habitat as well as conventional cultivation of orchid plants for the production of bioactive secondary metabolites. Accumulation of secondary metabolites in in vitro cultures of orchids has been reported in some studies (Table 2). Factors such as type of explants, composition and type of culture medium, growth regulators and their concentration, concentration of carbon source, culture age etc. influence biosynthesis and accumulation of secondary metabolites in in vitro culture.^{45,56} Though some studies have been done on secondary metabolite production from in vitro culture of orchids but still extensive research is needed with orchid culture to optimize factors that influence secondary metabolite production.

5. Genetic engineering and orchid improvement

Improvement of orchids for some major traits such as flower morphology, colour, fragrance, cut-flower longevity, resistance to diseases, stress etc. through

transgenic technologies can be advantageous over the breeding of orchids through traditional methods. Modification of orchids for desirable character by using plant genetic engineering has been demonstrated in a few orchid species.⁴⁰ Longevity of flowers of *Dendrobium chrysotoxum* in vase life has been enhanced by *Agrobacterium* mediated transfer of an antisense ACC oxidase gene into protocorms and the ACC oxidase inactivated in regenerated transgenic plants.¹¹ The bright yellow of the wild type color of *Oncidium* 'Gower Ramsey' flower was changed to more attractive light and white-yellow by using *Agrobacterium tumefaciens* mediated transfer of RNA interference for β -ring carotene hydroxylase (BCH2) gene.⁶⁵ Shoot multiplication of *Phalaenopsis amabilis*⁶¹ and *Dendrobium lasianthera*⁶³ has been enhanced by inserting Knat 1 (Knotted 1 like *Arabidopsis thaliana*) gene. *Phalaenopsis* orchid with enhanced resistance to Cymbidium Mosaic Virus (CymMV) and *Erwinia carotovora* was developed by double transformation of PLBs. Using Particle bombardment initially PLBs were transformed with CymMV coat protein cDNA (CP) then re-transformed by *Agrobacterium* mediated transfer of sweet pepper ferredoxin-like protein cDNA (Pflp).¹³ Though remarkable advances have been made for production of transgenic orchids however, extensive studies remain to be done.

6. Conclusion and Future prospects

Orchids are considered as largest and most diverse group of angiosperms. Apart from their use in ornamental industry they are also used in medicinal and food industry and many orchid species are considered as endangered due to their extensive collection and several other anthropological activities. Vegetative and sexual multiplication process is extremely slow to meet its demand worldwide. There is serious need to multiply orchids through alternate method and some orchids so far have been multiplied through in vitro culture technique. Though, plant tissue culture technique has been used mostly for asexual seed germination but multiplication of orchids through seeds generates heterozygous plants. Establishment of protocol for regeneration of orchids from various vegetative explants are required. Improvement in flower quality by strengthening of orchid biotechnology can advance floriculture industry. In vitro production of secondary metabolites from medicinal orchids is another aspect which is initiated in few orchid species and extensive emphasis should be given on metabolites engineering techniques to enhanced production and identification of secondary metabolite.

References

1. Ara H, Jaiswal U, Jaiswal VS. Synthetic seed: prospects and limitations. *Curr Sci* 2000; 78: 1438-1444.
2. Arditti J. Factors affecting the germination of orchid seeds. *Botanical Review* 1967; 33: 1-97.
3. Arditti J. Orchid seed germination and seedling culture - a manual. In: Arditti J, ed. *Orchid biology, reviews and perspectives II*. Ithaca, NY: Cornell University Press; 1982. p. 244-370.
4. Arditti J. *Fundamentals of Orchid Biology*. John Wiley & Sons, New York; 1992.
5. Bhadra SK, Hossain MM. In vitro germination and micropropagation of *Geodorum densiflorum* (Lam.) Schltr., an endangered orchid species. *Plant Tiss Cult* 2003; 13: 165-171.
6. Bhattacharyya P, Kumaria S, Job N, Tandon P. Phyto-molecular profiling and assessment of antioxidant activity within micropropagated plants of *Dendrobium thyrsiflorum*: a threatened, medicinal orchid. *Plant Cell Tiss Org Cult* 2015; 122: 535-550.
7. Bhattacharyya P, Kumar V, Doležal GJK, Staden JV. Deciphering the phenolic acid reserves and antioxidant activity within the protocorm like bodies of *Ansellia africana*: A vulnerable medicinal orchid. *Industrial Crops and Products*. 2019; 135: 21-29.
8. Bhattacharyya P, Kumar V, van Staden J. In vitro encapsulation based short term storage and assessment of genetic homogeneity in regenerated *Ansellia africana* (Leopard orchid) using gene targeted molecular markers. *Plant Cell Tiss Org Cult* 2018; 133: 299-310.
9. Bhattacharyya P, Paul P, Kumaria S, Tandon P. Transverse thin cell layer (t-TCL)-mediated improvised micropropagation protocol for endangered medicinal orchid *Dendrobium aphyllum* Roxb: an integrated phytomolecular approach. *Acta Physiol Plant* 2018; 40: 137 DOI: 10.1007/s11738-018-2703-y
10. Bose B, Choudhury H, Tandon P, Kumaria S. Studies on secondary metabolite profiling, anti-inflammatory potential, in vitro photoprotective and skin-aging related enzyme inhibitory activities of *Malaxis acuminata*, a threatened orchid of nutraceutical importance. *Journal of Photochemistry and Photobiology B: Biology* 2017; 173: 686-695. <https://doi.org/10.1016/j.jphotobiol.2017.07.010>
11. Bunnag S, Pilahome W. Agrobacterium - mediated transformation of *Dendrobium chrysotoxum* Lindl. *Afr J Biotechnol* 2012; 11: 2472-2476.
12. Cardoso CJC, Zanello CA, Chen JT. An overview of orchid protocorm-like bodies: mass propagation, Biotechnology, Molecular Aspects, and Breeding. *Int J Mol Sci* 2020; 21: 985-1016. doi:10.3390/ijms21030985
13. Chan YL, Lin KH, Sanjaya, Liao LJ, Chen WH, Chan MT. Gene stacking in *Phalaenopsis* orchid enhances dual tolerance to pathogen attack *Transgenic Res* 2005; 14: 279-288. DOI 10.1007/s11248-005-0106-5
14. Chase MW, Cameron KM, Freudestein JV, Pridgeon AM, Salazar G, Berg CV, Schuiteman A. An update classification of orchidaceae. *Bot J Linn Soc* 2015; 177: 151-174.
15. Christenhusz MJM, Byng JW. The number of known plant species in the world and its annual increase. *Phytotaxa* 2016; 261: 201-217.
16. Chugh S, Guha S, Rao IU. Micropropagation of orchids: A review on the potential of different explants. *Scientia Horti* 2009; 122: 507-520. doi:10.1016/j.scienta.2009.07.016
17. Das MC, Kumaria S, Tandon P. In vitro propagation and conservation of *Dendrobium lituiflorum* Lindl through Protocorm-Like Bodies. *J Plant Biochem Biotechnol* 2008; 17: 177-180.
18. Das MC, Kumaria S, Tandon P. Storage and high conversion frequency of encapsulated protocorm-like bodies of *Cymbidium devonianum* (orchid). *J Horti Sci Biotechnol* 2011; 86: 611-615.
19. Datta KB, Kanjilal B, De Sarker D. Artificial seed technology: development of a protocol in *Geodorum densiflorum* (Lam) Schltr.-An endangered orchid. *Curr Sci* 1999; 76: 1142-1145.
20. Dohling S, Kumaria S, Tandon P. Multiple shoot induction from axillary bud cultures of the medicinal orchid, *Dendrobium longicornu*. *AoB Plants* 2012; pls032; doi:10.1093/aobpla/pls032
21. Gamborg OL, Miller RA, Ojima K. Nutrient requirement of suspension cultures of soyabean root cells. *Expt Cell Res* 1968; 50: 151-158.
22. Giri L, Dhyani P, Rawat S, Bhatt ID, Nandi SK, Rawal RS, Pande V. In vitro production of phenolic compounds and antioxidant activity in callus suspension cultures of *Habenaria edgeworthii*: A rare Himalayan medicinal orchid. *Indust Crops Prod* 2012; 39: 1-6.
23. Gogoi K, Kumaria S, Tandon P. Cryopreservation of *Cymbidium eburneum* Lindl. and *C. hookerianum* Rchb. f., two threatened and vulnerable orchids via encapsulation-dehydration. *In Vitro Cell Dev Biol - Plant* 2013; 49: 248-254.
24. Goswami K, Yasmin S, Nasiruddin KM, Khatun F, Akte J. In Vitro Regeneration of *Dendrobium*

- sp. of Orchid Using Leaf Tip as Explant. *J Environ Sci Nat Resources* 2015; 8: 75-78.
25. Gupta A. Asymbiotic seed germination in orchids: role of organic additives. *Intl Advan Res J Sci, Engineer Tech* 2016; 3: 143-147.
 26. Harrison CR. Ultrastructural and histochemical changes during the germination of *Cattleya auratiaca* (Orchidaceae). *Botanical Gazette* 1977; 138: 41-45.
 27. Harvais G. Growth requirements and development of *Cypripedium reginae* in axenic cultures. *Canadian J Bot* 1973; 51: 327-332.
 28. Islam SS, Bhattacharjee B. Plant regeneration through somatic embryogenesis from leaf and root explants of *Rhynchostylis retusa* (L.) Blume. *Appl Biol Res* 2015; 17: 158-165.
 29. Jaime, A, Teixeira, DS. Response of hybrid *Cymbidium* (Orchidaceae) protocorm like bodies to 26 plant growth regulators. *Botanica Lithuanica* 2014; 20: 3-13.
 30. Janarthanam B, Seshadri S, Plantlet regeneration from leaf derived callus of *Vanilla planifolia* Andr. *In Vitro Cell Dev Biol - Plant* 2008; 44: 84-89.
 31. Kaur S. In vitro regeneration of shoot from Nodal explants of *Dendrobium chrysotoxum* Lindl. *J Hort Res* 2017; 25: 27-34. DOI: 10.1515/johr-2017-0003
 32. Knudson L. Nonsymbiotic germination of orchid seeds. *Botanical Gazette* 1922; 73: 1-25.
 33. Knudson L. A new nutrient solution for germination of orchid seeds. *American Orchid Soc Bulletin* 1946; 15: 214-217.
 34. Kalimuthu K, Senthikumar R, Murugalatha N. Regeneration and mass multiplication of *Vanilla planifolia* Andr. - a tropical orchid. *Curr Sci* 2006; 91: 1401-1403.
 35. Lubinsky P, Bory S, Hernandez JH, Kim SC, Gomez-Pompa A. Origins and dispersal of cultivated *Vanilla* (*Vanilla planifolia* Jacks. [Orchidaceae]). *Economic Bot* 2008; 62: 127-138.
 36. Martin KP. Micropropagation of the bamboo orchid (*Arundina graminifolia* D. Don) Hochr.) through protocorm-like-bodies using node explants. *Propagation Ornament Plants* 2007; 7: 97-100.
 37. Mathur J, Ahuja PS, Lal N, Mathur AK. Propagation of *Valeriana wallichii* DC. Using apical and axial shoot buds. *Plant Sci* 1989; 60: 111-116.
 38. Mayer JLS, Stancato GC, AppezzatoDa-Gloria B. Direct regeneration of protocorm-like bodies (PLB) from leaf apices of *Oncidium flexuosum* Sims (Orchidaceae). *Plant Cell Tiss Org Cult* 2010; 103: 411-416.
 39. Mehraj H, Alam MM, Habiba SU, Mehbub H. LEDs combined with CHO sources and CCC priming PLB regeneration of *Phalaenopsis*. *Horticulture* 2019; 5: 34-45.
 40. Mii M, Chin DP. Genetic transformation of orchids. *Acta Hort* 2010; 878: 461-466. DOI: 10.17660/Acta Hort.2010.878.59
 41. Mitra GC, Prasad RN, Choudhury AR. Inorganic correlated salts and differentiation of protocorms in seed of an orchid (*Dendrobium fimbriatum*) and correlated changes in its free amino acid context. *Indian J Exp Biol* 1976; 14: 350-351.
 42. Mohanty P, Nongkling P, Das MC, Kumaria S, Tandon P. Short-term storage of alginate-encapsulated protocorm-like bodies of *Dendrobium nobile* Lindl.: an endangered medicinal orchid from north-east India. *3 Biotech* 2013; 3: 235-239. DOI 10.1007/s13205-012-0090-4
 43. Murashige T, Skoog F. A revised medium for rapid growth and bio-assays with tobacco tissue cultures. *Plant Physiol* 1962; 15: 473-497.
 44. Naing AH, Chung JD, Park IS, Lim KB. Efficient plant regeneration of the endangered medicinal orchid, *Coelogyne cristata* using protocorm-like bodies. *Acta Physiol Plant* 2011; 33: 659-666. <https://doi.org/10.1007/s11738-010-0586-7>
 45. Pant B. Medicinal orchids and their uses: Tissue culture a potential alternative for conservation. *African J Plant Sci* 2013; 7: 448-467.
 46. Park HY, Kang KW, Kim DH, Sivanesan I. In vitro propagation of *Cymbidium goeringii* Reichenbach fil. through direct adventitious shoot regeneration. *Physiol Mol Biol Plants* 2018; 24: 307-313. <https://doi.org/10.1007/s12298-017-0503-2>
 47. Park SY, Murthy HN, Paek KY. Protocorm-like body induction and subsequent plant regeneration from root tip cultures of *Doritaenopsis*. *Plant Sci.* 2003; 164: 919-923.
 48. Paul P, Joshi M, Gurjar D, Shailajan S, Kumaria S. In vitro organogenesis and estimation of β -sitosterol in *Dendrobium fimbriatum* Hook.: An orchid of biopharmaceutical importance *South African J Bot* 2017; 113: 248-252.
 49. Paul S, Kumaria S, Tandon P. An effective nutrient medium for asymbiotic seed germination and large-scale in vitro regeneration of *Dendrobium hookerianum*, a threatened orchid of northeast India *AoB Plants* 2012; plr032; doi:10.1093/aobpla/plr032
 50. Pradhan S, Tiruwa B, Subedee BR, Pant B. In vitro germination and propagation of a threatened medicinal orchid, *Cymbidium aloifolium* (L.) Sw. through artificial seed. *Asian Pacific J Tropical Biomedicine* 2014; 4: 971-976. <https://doi.org/10.12980/>

- APJTB.4.2014APJTB-2014-0369
51. Roy AR, Sajeev S, Pattanayak A, Deka BC. TDZ induced micropropagation in *Cymbidium giganteum* Wall. Ex Lindl. and assessment of genetic variation in the regenerated plants. *Plant Growth Regul* 2012; 68: 435-445.
 52. Sharma A, Tandon P, Kumar A. Regeneration of *Dendrobium wardianum* Warner. (Orchidaceae) from synthetic seeds. *Indian J Exp Biol* 1992; 30: 747-748.
 53. Sharma V. Regenerative competence of thin cell layer (epidermal peel) for in vitro propagation of *Cattleya 'Almakee'*. *Asian J Microbiol Biotech Env Sci* 2018; 20: S65-S69.
 54. Sarmah D, Kolukunde S, Sutradhar M, Singh BK, Mandal T, Mandal N. A review on: in vitro cloning of orchids. *Int J Curr Microbiol App Sci* 2017; 6: 1909-1927. <https://doi.org/10.20546/ijcmas>
 55. Sarmah DK, Borthakur M, Borua PK. Artificial seed production from encapsulated PLBs regenerated from leaf base of *Vanda coerulea* Griff. ex. Lindl.: An endangered orchid. *Curr sci* 2010; 98: 686-690.
 56. Singh A, Duggal S. Medicinal Orchids: An Overview. *Ethnobotanical Leaflets* 2009; 13: 351-363.
 57. Singh M, Yadav C, Bano F. Plant Regeneration from Alginate encapsulated nodes of *Andrographis paniculata* (Burm. F.) Wallich ex Nees. A potential medicinal herb. *Vegetos* 2018; 31: 28-33.
 58. Singh M, Kumaria S, Tandon P. In vitro multiplication of *Thunia marshalliana* Rcbh. f. through mature seeds and pseudonode culture. *Indian J Biotech* 2014; 13: 268-273.
 59. Singh N, Kumaria S. Deciphering the role of stress elicitors on the differential modulation of chalcone synthase gene and subsequent production of secondary metabolites in micropropagated *Coelogyne ovalis* Lindl., a therapeutically important medicinal orchid. *South African J Bot* 2020; <https://doi.org/10.1016/j.sajb.2020.06.019>.
 60. Sinha P, Hakim M, Alam M. Efficient micropropagation of *Phalaenopsis amabilis* (L.) BL. cv. 'Cool Breeze' using inflorescence axis thin sections as explants. *Propagation Ornament Plants* 2007; 7: 9-15.
 61. Semiarti E, Indrianto A, Purwantoro A, Isminingsih S, Suseno N, Ishikawa T, Yoshiaka Y, Machida Y, Machida C. Agrobacterium-mediated transformation of the wild orchid *species Phalaenopsis amabilis*. *Plant Biotechnol* 2007; 24: 265-272.
 62. Tsai CF, Huang CL, Lin YL, Lee YC, Yang YC, Huang NK. The neuroprotective effects of an extract of *Gastrodia elata*. *J Ethnopharmacology* 2011; 138: 119-125.
 63. Utami ESW, Hariyanto S, Manuhara YSW. *Agrobacterium tumefaciens*-mediated transformation of *Dendrobium lasianthera* J.J.Sm: An important medicinal orchid. *J Genetic Engineer Biotech* 2018; 16: 703-709.
 64. Verpoorte R, Contin A, Memelink J. Biotechnology for the production of plant secondary metabolites. *Phytochem Reviews* 2002; 1: 13-25.
 65. Wang HM, To KY, Lai HM, Jeng ST. Modification of flower colour by suppressing β -ring carotene hydroxylase genes in *Oncidium*. *Plant Biol (Stuttg)* 2016; 18: 220-229. doi: 10.1111/plb.12399
 66. Weatherhead MA, Zee SY, Barretto G. Some observations on the early stages of development of *Eulophia yushuiana*. *Memoirs of the Hong Kong Natural History Society* 1986; 17: 85-90.
 67. Yeowa LC, Chewa BL, Sreeramanan S. Elevation of secondary metabolites production through light-emitting diodes (LEDs) illumination in protocorm-like bodies (PLBs) of *Dendrobium* hybrid orchid rich in phytochemicals with therapeutic effects. *Biotech Rep* 2020; 27: e00497 <https://doi.org/10.1016/j.btre.2020.e00497>
 68. Zeng S, Wu K, da Silva JAT, Zhang J, Chen Z, Xia N, Duan J. Asymbiotic seed germination, seedling development and reintroduction of *Paphiopedilum wardii* Sumerh., an endangered terrestrial orchid. *Sci Hortic* 2012; 138:198-209.
 69. Zhang S, Yang Y, Li J, Qin J, Zhang W, Huang W, Hu H. Physiological diversity of orchids. *Plant Diversity* 2018; 40:196-208.
 70. Zhang Z, Yan Y, Tian Y, Li J, He J-S, Tang Z. Distribution and conservation of orchid *species* richness in China. *Biol Conservation* 2015; 181: 64-72.