

## Acute Myocardial Infarction Presenting with Atypical Symptoms

Priyanka V.N.<sup>1</sup>, Anil Kumar V.R.<sup>2</sup>, Nagamalesh U.M.<sup>3</sup>

### Author's Affiliation:

<sup>1</sup>Assistant Professor <sup>2</sup>Associate Professor, Emergency Medicine, Ramaiah Medical College, Bengaluru, Karnataka 560054, India.  
<sup>3</sup>Sr. Consultant, Department of Cardiology, Ramaiah Narayana Hrudayalaya Heart centre, Bangalore, Karnataka 560054, India.

### Corresponding Author:

**Anil Kumar V R**, Associate Professor, Emergency Medicine, Ramaiah Medical College, Bengaluru, Karnataka 560054, India.

E-mail: [dranil417@gmail.com](mailto:dranil417@gmail.com)

Received on 26.07.2018,

Accepted on 17.09.2018

### Abstract

**Introduction:** Chest pain is the most common key presentation of acute coronary syndrome presenting to Emergency department. But few patients may present with atypical symptoms and pose a great challenge to the Emergency physicians. Diagnosis has to be made early since maximal mortality occurs within first few hours if treatment is not instituted early. A cross-sectional study was conducted in our hospital to study the proportion of atypical symptoms in patients with acute coronary syndrome. **Methods:** A Cross-sectional study was conducted in our hospital from September 2017 to April 2018. All patients aged >20 years presenting to emergency department and fulfilling the following diagnostic criteria were included. The WHO diagnostic criteria were used for diagnosis of Acute coronary syndrome [9]. 1) Clinical history of ischemic type of chest pain 2) Changes in serial ECG tracings a) ST-segment elevation of more than 1mm in 2 limb leads. b) ST-segment elevation of more than 2mm in 2 or more contiguous chest leads. c) ST segment depression >0.5mm at J point in >2 contiguous leads with positive troponin I. 3) Positive Serum cardiac biomarkers (troponin I) The diagnosis was confirmed if 2 out of 3 above components were positive. ECG was done for all patients presenting to ER with symptoms suggestive of acute coronary syndrome. Cardiac enzymes were sent in patients with nonspecific ECG changes. **Results:** 323 patients were included in our study. Out of which, 269 patients presented with typical chest pain and 54 (16.71%) patients presented with atypical symptoms. The incidence in male and female population was 44.4% and 55.5% respectively. The incidence of atypical symptoms increased with increasing age and was highest in the age group of 71-80 yr. **Conclusion:** It is crucial to consider acute coronary syndrome as a differential diagnosis in patient with atypical presentation & Emergency physicians should have a knowledge of various atypical presentations. As missed Myocardial Infarction has dire consequences, continued efforts are required to reduce incidence, overall mortality & morbidity.

**Keywords:** Atypical Symptoms; Acute Coronary Syndrome; Atypical MI.

### Introduction

Chest pain is the most common key presentation of acute coronary syndrome presenting to Emergency department [1]. But few patients may present with atypical symptoms and pose a great challenge to the Emergency physicians [2,3]. Failing to consider acute coronary syndrome in patients with these atypical symptoms is a pitfall. Diagnosis has to be made early since maximal mortality

occurs within first few hours if treatment is not instituted early [2,4-8]. A cross-sectional study was conducted in our hospital to study the proportion of atypical symptoms in patients with acute coronary syndrome. Our Objective was to emphasize the need for timely consideration of acute coronary syndrome as one of the Differential Diagnosis in patients presenting to Emergency department with atypical symptoms.

## Methods

A Cross-sectional study was conducted in our hospital on patients who presented to Emergency department with symptoms of Acute coronary syndrome from September 2017 to April 2018. All patients aged >20 years presenting to emergency department and fulfilling the following diagnostic criteria were included. Typical symptoms included substernal chest pain/discomfort with/without radiation to left arm, neck or jaw. Any symptoms which falls short of typical features would constitute atypical chest pain. The WHO diagnostic criteria were used for diagnosis of Acute coronary syndrome [9].

1. Clinical history of ischemic type of chest pain
2. Changes in serial ECG tracings.
  - a) ST-segment elevation of more than 1mm in 2 limb leads.
  - b) ST-segment elevation of more than 2mm in 2 or more contiguous chest leads.
  - c) ST segment depression >0.5mm at J point in >2 contiguous leads with positive troponin I.
3. Positive Serum cardiac biomarkers (troponin I)

The diagnosis was confirmed if 2 out of 3 above components were positive. ECG was done

for all patients presenting to ER with symptoms suggestive of acute coronary syndrome. Cardiac enzymes were sent in patients with nonspecific ECG changes.

### Statistical Methods

Descriptive statistics of patients' characteristics and atypical symptoms was analyzed & summarized in terms of percentage

## Results

323 patients were included in our study. Out of which, 269 patients presented with typical chest pain and 54 (16.71%) patients presented with atypical symptoms. Twenty-four male patients and 30 female patients had atypical symptoms. The incidence in male and female population was 44.4% and 55.5% respectively. The incidence of atypical symptoms increased with increasing age and was highest in the age group of 71-80 yr, with an incidence of 27.77%.

In patients with atypical symptoms, we found that 21 patients had diabetes and 14 patients had hypertension. Various atypical symptoms included dyspnoea, epigastric pain, palpitation, generalised weakness, nausea/vomiting, sweating. Jaw pain, syncope. Thirty-seven patients with STEMI and 17 patients with NSTEMI had atypical symptoms.

**Table 1:** Age distribution

Age in years	Typical symptoms	Atypical symptoms
<30	3	-
31-40	8	-
41-50	34	5(9.2%)
51-60	67	9(16.6%)
61-70	71	12(22.2%)
71-80	47	15(27.77%)
>80	39	13(24%)
Total	269(83.28%)	54(16.71%)

**Table 2:** Gender distribution

Sex	Typical symptoms	Atypical symptoms
Male	141	24(44.4%)
Females	128	30(55.5%)
Total	269	54

**Table 3:** Distribution of risk factors

Risk factors	Typical symptoms	Atypical symptoms
Hypertension	138(51%)	14(25.9%)
Diabetes	155(57.6%)	21(38.8%)

**Table 4:** Various atypical symptoms

Atypical symptoms	No of patients	Percentage %
Dyspnoea	21	38.8
Upper abdomen pain	11	20.37
Nausea/ vomiting	4	7.4
Generalised weakness	4	7.4
Jaw pain	2	3.7
Syncope	3	5.5
Sweating	6	11.1
Palpitation	3	5.5

## Discussion

Heberden was the first to describe typical symptoms of acute Myocardial Infarction. Chest pain is the classic presenting feature of acute coronary syndromes. The World Health Organisation requires the presence of chest pain as one of the main components for diagnosis of acute coronary syndromes [9]. Acute coronary syndromes presenting with atypical symptoms are less likely to receive a timely ECG & further intervention. Despite diagnostic advances, missed acute Myocardial Infarction constitutes for 2-10% of all cases. These patients tend to have longer delays in diagnosis, treatment & therefore have an increased mortality rate [2,4-8]. Atypical presentation of acute coronary syndromes is a significant independent predictor of in-hospital mortality [10]. Myocardial infarction without chest pain had 23.3% in-hospital mortality rate as compared to 9.3% among patients with chest pain [2].

In our study the proportion of patients with atypical symptoms was 16.71%. Several studies have reported the incidence of atypical presentations between 4.7% to 33% [2,4-8,11,12]. We found that the incidence is higher in women (55.5%) comparable with other studies [8,3,11]. The incidence increased with age with highest incidence of 51.4% in patients aged >70 years. According to Global Registry of Acute Coronary Events (GRACE) and National Registry of Myocardial Infarction (NRFMI)-2, incidence of atypical presentation was as high as 33% and 44% respectively in patients aged >75 yrs [2,4,13,17]. According to Global Registry of Acute Coronary Events (GRACE) there was higher female proportion correlating to our study results.

We found dyspnoea was the most common atypical presentation followed by epigastric pain. Various other atypical symptoms included jaw pain, generalised weakness, syncope, sweating and palpitation. Several studies have shown that

dyspnoea is the most frequent symptom followed by gastrointestinal symptoms. [4,11].

The incidence of atypical presentation was 38.8% in diabetics according to our study. In various other studies the incidence of diabetics presenting with atypical symptoms ranges from 35% to 47% [13-16]. Various studies have concluded that diabetes is an independent predictor for the presence of silent MI [13-16].

The mechanism of atypical presentation is multifactorial. Diabetes may have a diminished awareness of ischemic pain which could result in atypical manifestation. This may be attributed to autonomic neuropathy and prolongation of the anginal perpetual threshold. However, the importance of cardiac autonomic neuropathy in silent Myocardial infarction is still debated. Droste et al have described "defective warning system", an altered threshold to pain perception or other noxious stimuli to be related to silent infarcts [17].

Many studies have shown that the region of infarction is not associated with atypical manifestations, although patients with IWMI tend to often present with gastrointestinal symptoms than those with AWTMI [3,18]. In our study, 24 patients had inferior wall MI, 13 patients had anterior wall MI and 17 were diagnosed with NSTEMI. Epigastric pain was the common atypical symptom in IWMI and dyspnoea was common in AWTMI. NSTEMI patients presented with non-specific symptoms like generalised weakness, palpitation, syncope, sweating.

## Conclusion

The incidence of atypical presentation of acute coronary syndrome in Emergency department is high & accurate diagnosis remains a great challenge to the Emergency Physicians. Hence, it is crucial to consider acute coronary syndrome as a differential

diagnosis in such patients & Emergency physicians should have a knowledge of various atypical presentations. High index of suspicion especially in elderly with comorbidities helps in early detection and management. As missed Myocardial Infarction has dire consequences, continued efforts are required to reduce incidence, overall mortality & morbidity.

### References

1. Bean WB. Masquerades of myocardial infarction. *Lancet*. 1997;1:1044-46.
2. Canto JG, Shlipak MG, Rogers WJ, Malmgren JA, Frederick PD, Lambrew CT, Ornato JP, Barron HV, Kiefe CI. Prevalence, Clinical Characteristics and Mortality Among Patients With Myocardial Infarction Presenting Without Chest Pain. *JAMA*. 2000;283:3223-9
3. Chowta K N, Prijith P D, Chowta M N. Modes of presentation of acute myocardial infarction. *Indian J Crit Care Med* 2005;9:151-4.
4. Brieger D, Eagle KA, Goodman SG, Steg PG, Budaj A, White K, Montalescot G. Acute coronary syndromes without chest pain, an underdiagnosed and undertreated high-risk group: insights from the Global Registry of Acute Coronary Events. *Chest*. 2004;126:461-69.
5. Dorsch MF, Lawrance RA, Sapsford RJ, Durham N, Oldham J, Greenwood DC, Jaackson BM, Morrell C, Robinson MB, Hall AS, EMMACE Study group. Poor prognosis of patients presenting with symptomatic myocardial infarction but without chest pain. *Heart* 2001;86:494-8.
6. Zdziennicka J, Siudak Z, Zawislak B, Dziewierz A, Rakowski T, Dubiel J, Dudek D. Patients with non-ST-elevation myocardial infarction and without chest pain are treated less aggressively and experience higher in-hospital mortality. *Kardiologia polska* 2007;65:769-75.
7. Herlitz J, Karson BW, Richter A, Strombom U, Hjalmarson A. Prognosis for patients with initially suspected acute myocardial infarction in relation to presence of chest pain. *Clin Cardiol* 1992;15:570-6.
8. Goldstein RE, Boccuzzi SJ, Cruess D. Prognosis after hospitalization for acute myocardial infarction not accompanied by typical ischemic chest pain. The Multicentre Diltiazem Postinfarction Trial Research Group. *Am J Med* 1995;99:123-31.
9. Gillum RF, Fortmann SP, Prineas RJ, Kottke TE. International diagnostic criteria for acute myocardial infarction and acute stroke. *Am Heart J*. 1984;108:150-58.
10. El-Menyer A, Zubaid M, Sulaiman K, Almahmeed W, Singh R, Alsheikh-ali A A, Suwandi J, GRACE investigators. Atypical presentation of acute coronary syndrome: A significant independent predictor of in-hospital mortality. *Journal of cardiology* 2011;57:165-71.
11. Jones ID, Slovis CM. Emergency department evaluation of the chest pain patient. *Emerg Med Clin North Am* 2001;19:269-82.
12. Grosmaître P, Vavasseur O, Yachouh E, Courtial Y, Jacob X, Meyran S, Lantelme P. Significance of atypical symptoms for the diagnosis and management of myocardial infarction in elderly patients admitted to emergency departments. *Archives of cardiovascular disease* 2013;106:586-92.
13. Sukhija R, Dhanwal D, Gambhir DS, Dewan R. Silent myocardial ischaemia in patients with type II diabetes mellitus and its relation with autonomic dysfunction. *Indian Heart J*. 2000;52:540-546.
14. Hwang SY, Park EH, Shin ES, Jeong MH. Comparison of factors associated with atypical symptoms in younger and older patients with acute coronary syndromes. *J Korean Med Sci*. 2009;24:789-94.
15. Arenja N, Mueller C, Ehl NF, Brinkert M, Roost K, Reichlin T, Sou SM, Hochgruber T, Osswald S, Zellweger MJ. Prevalence, extent, and independent predictors of silent myocardial infarction. *Am J Med*. 2013;126:515-22.
16. Scognamiglio R, Negut C, Ramondo A, Tiengo A, Avogaro A. Detection of coronary artery disease in asymptomatic patients with type 2 diabetes mellitus. *J Am Coll Cardiol*. 2006;47:65-71.
17. Droste C, Roskamm H. Pain mechanisms in symptomatic and silent ischemia. *Isr J Med Sci*. 1989;25:487-92.
18. Shakoor M T, Falik S, Shah F. Incidence of atypical presentation of myocardial infarction. *Pakistan Heart Journal* 2008;41:15-20.