

Original Research Article

Skin Adnexal Tumors: An Experience at Tertiary Care Centre

Sonam S. Nandyal¹, Sharvani R. Setty², Rekha B. Puranik³, Sujatha S. Giriyan⁴,
Narayan Murthy C⁵

¹Associate Professor, ²Assistant Professor, ³Professor and Head, Department of Pathology, Basaveshwara Medical College and Hospital, Chithradurga, Karnataka 577502, India. ³Professor, ⁴Professor and Head, Department of Pathology, Karnataka Institute of Medical Sciences, Hubli, Karnataka 580022, India.

How to cite this article:

Sonam S. Nandyal, Sharvani R. Setty, Rekha B. Puranik et al. Skin Adnexal Tumors: An Experience at Tertiary Care Centre. Indian J Pathol Res Pract. 2019;8(3):307-316.

Abstract

Background: Skin adnexal tumors are large and diverse group of uncommon neoplasms usually misdiagnosed clinically due to their large variety and variants. Histopathology usually helps in establishing the diagnosis. **Aims:** To study the spectrum and morphological features of skin adnexal tumors (SATs) at our centre. **Methods:** This is both retrospective and prospective descriptive study over a period of 5 years (June 2005 to June 2010). The study included all the biopsies and excision specimens submitted for histopathological examination. **Results:** Most of the SATs were benign (25/31) with head and neck being the most common location (67.74%). Male to female ratio was 1:1.8. There was equal occurrence of tumors of sweat gland and hair follicle differentiation (45%). The most common varieties of tumors encountered in the present study included proliferating trichilemmal tumor and hidradenoma. The concordance between clinical and histopathological diagnosis was found to be 6.45% approximately. **Conclusions:** SATs are relatively uncommon. As they are often clinically misdiagnosed, histopathology is mandatory for the diagnosis. Although they are not numerically important or therapeutically problematic, nevertheless they form the spectrum of interesting lesions.

Keywords: Adnexal tumor; Proliferating trichilemmal tumor; Hidradenoma; Pilomatricoma.

Corresponding Author:

Sharvani R. Setty, Assistant Professor,
Dept of Pathology, Basaveshwara Medical
College and Hospital, Chitradurga, Karnataka
577501, India.

E-mail: sharvanisetty11@gmail.com

Received on 11.04.2019,

Accepted on 30.04.2019

Introduction

Skin is a complex tissue with an intricate adnexae. Skin adnexae give rise to a perplexing number of neoplasms probably more than the accounted

numbers. These tumors essentially differentiate in the line of hair follicles, sebaceous glands, eccrine and apocrine sweat glands [1]. Skin adnexal tumors (SATs) are rare lesions and in view of the fact that they are so infrequently encountered in practice, they may cause difficulty in diagnosis.

The embryological relations, anatomical distribution, combinations of neoplasms and the associations of SATs are all genuine biological phenomena [2]. Understanding the concept of histogenesis is essential for diagnosis of these tumors [3].

Apart from their rarity, difficulties in diagnosis also result due to their large variety, non specific clinical appearance, their frequent differentiation along two or more adnexal lines simultaneously and their complex nomenclature [1,4,5]. Benign SATs have a tendency to mimic malignant lesions and usually manifest in any age group. Malignant SATs are rare, locally aggressive and have the potential for nodal involvement and distant metastasis with a poor clinical outcome. Therefore establishing a diagnosis of malignancy in SAT is important for therapeutic and prognostic purpose [6].

There is scarcity of literature on the clinicopathological features of SATs. Histopathological study is mandatory to establish diagnosis which is the most valuable means of diagnosis in dermatopathology, inspite of its own limitations. Literature search showed very few studies from India, majority of them being isolated case reports. Keeping in view of these facts, the present study documents our experience with SATs with emphasis on clinical data, histomorphology, diagnostic approach and brief review of literature.

Materials and Methods

This was a descriptive study conducted at the department of Pathology in a tertiary care hospital over a period of 5 years from 2005 to 2010. The study included all the biopsies and excision specimens submitted for histopathological examination. Data was collected after a meticulous study of inpatient and outpatient dermatology, surgery records and histopathology records. The specimens were fixed in 10% formalin and the gross features were examined and sampled depending on the size of tumor. Tissue was routinely processed and sections of 3 to 5 micron thickness were taken and stained with hematoxylin and eosin. Wherever necessary relevant special stain was done. The facility for

histochemical staining for enzymes was not available in our laboratory setup.

Results

During the 5 year study period, 135 skin tumors were noted, of which 31 were adnexal in origin. Prevalence of SATs compared to total surgical pathology specimens was found to be 0.12%. The tumors were categorized as per WHO classification. Majority of tumors in our series were benign (25 cases - 80.6%) with equal occurrence of hair follicle and sweat gland tumors (45%)(Table 1). Age range was wide from 11 to 80 years. One of the most notable findings in the present study was female predominance with male to female ratio of 1:1.8. Head and neck was the most frequently involved site for SATs accounting for 67.74%. The tumor size ranged from 0.5 cm to 7 cm. Most of the SATs (70.96%) were <2.5 cm in maximum dimensions. SATs were clinically suspected in 6.45% of patients before excision.

Tumors with Hair follicle differentiation

Hair follicle tumors had a female predominance and proliferating trichilemmal tumor (PTT) was the most common type. Next in frequency were pilomatricoma, trichoepithelioma and Malignant PTT (Table 2).

There were 7 PTTs among which 5 occurred in females and 2 in males. Age range was 14-60 years. Most common site was scalp (5 cases). Most common clinical diagnosis was sebaceous cyst. Histologically, all cases were well circumscribed and composed of squamoid cells showing trichilemmal keratinization and peripheral palisading. The tumor cells showed mild pleomorphism as well as dyskeratosis (Figs. 1,2). Calcification was noted in 4 of them. Presence of squamous eddies and cholesterol clefts were the other changes observed.

Four pilomatricomas were encountered in 3 females and 1 male subject. In one case, a 30 year old female presented with recurrence of axillary pilomatricoma despite wide surgical excision (Fig. 3). Histologically all 4 tumors were well

Table 1: Distribution of tumors according to line of differentiation and behaviour

| | Benign | Malignant | Total |
|---|--------|-----------|-------|
| Tumors with Hair follicle differentiation | 13 | 1 | 14 |
| Tumors with sweat gland differentiation | 11 | 3 | 14 |
| Tumors with Sebaceous differentiation | 1 | 2 | 3 |

Table 2: Tumors with hair follicle differentiation

| No of cases | Age/sex | Site | Clinical details | Histopathological Diagnosis |
|-------------|---------|-------------------|--|---|
| 1 | 30 yr/F | scalp | Swelling | Proliferating trichilemmal tumor |
| 2 | 40 yr/F | Scalp | Non healing ulcer - SCC | Proliferating trichilemmal tumor |
| 3 | 14 yr/F | Scalp | Sebaceous cyst | Proliferating trichilemmal tumor |
| 4 | 60 y/F | Scalp | Sebaceous cyst | Proliferating trichilemmal tumor |
| 5 | 45 yr/F | scalp | Ulcer - SCC | Proliferating trichilemmal tumor |
| 6 | 55 y/M | Foot | Swelling - Sebaceous cyst | Proliferating trichilemmal tumor |
| 7 | 60 y/M | Axilla | Cystic swelling Sebaceous cyst | Proliferating trichilemmal tumor with pilar cyst |
| 8 | 18 y/F | Face | Swelling sebaceous cyst | Pilomatricoma |
| 9 | 26 y/F | Suboccipital | Hard swelling - osteoma | Pilomatricoma with extensive calcification |
| 10 | 30 y/F | Axilla | Recurrent bony hard swelling | Giant Pilomatricoma with extensive calcification |
| 11 | 35 y/M | Leg | Mobile Swelling Sebaceous cyst | Pilomatricoma with giant cell reaction |
| 12 | 86 y/F | Dorsum of nose | Asymptomatic nodule Adnexal tumor | Trichoepithelioma |
| 13 | 28 y/F | Face | Multiple asymptomatic nodules - BCC | Trichoepithelioma |
| 14 | 28 y/F | Scalp | Ulcerative lesion - SCC | Malignant proliferating trichilemmal tumor |




Fig. 1: Proliferating trichilemmal tumor: Scalp swelling measuring 4x3 cm



Fig. 3: Recurrent calcified Pilomatricoma: Axillary bony hard swelling measuring 7x4 cm




Fig. 2: Proliferating trichilemmal tumor: squamoid cells showing trichilemmal keratinisation (abrupt keratinization) and peripheral palisading. The tumor cells show mild pleomorphism as well as dyskeratosis (H&E 10X).




Fig. 4: Pilomatricoma: Leg Swelling measuring 2x1 cm.

circumscribed, composed of islands of basophilic cells and shadow cells (Figs. 4,5). Extensive calcification was seen in 2 of them and one tumor showed giant cell reaction to keratin.

Both trichoepitheliomas presented as nodular lesions over face. Histologically, showed islands of basaloid cells in the dermis with peripheral palisading (Fig. 6). Horn cysts, abortive hair follicles and presence of fibrous stroma were noted.

We came across one rare case of malignant PTT in 28 year old female who presented with an ulcer over scalp and the tumor had infiltrative margins composed of pleomorphic squamoid cell islands




Fig. 5: Pilomatricoma: composed of islands of basophilic cells and shadow cells (H & E, 10X).




Fig. 6: Trichoepithelioma: Well circumscribed dermal tumor with islands of basophilic cells (H & E, 10X).




Fig. 7: Malignant proliferating trichilemmal tumor: pleomorphic squamous cell islands with areas of abrupt keratinisation and atypical frequent mitotic figures(H&E, 10X).

with areas of abrupt keratinisation and atypical frequent mitotic figures (Fig. 7).

Tumors with sweat gland differentiation

Benign sweat gland tumors comprised of hidradenoma, which formed the bulk of cases followed by one case each of cylindroma (Fig. 8), syringoma, Syringocystadenoma Papilliferum (SCP), apocrine hidrocystoma (Fig. 9) and chondroid syringoma (Fig. 10) (Table 3).

Hidradenoma was seen in 4 females and 2 males. Age range was 18-75 years. Three each were




Fig. 8: Cylindroma: The tumor show dual population of cells with jigsaw puzzle pattern (H & E, 10X). Hyaline basement membrane showed PAS positivity (inset).




Fig. 9: Apocrine hidrocystoma: Multiloculated cystic lesion lined by columnar secretory cells showing decapitation (H & E,40X)




Fig 10: Chondroid syringoma: Epithelial cells arranged in solid sheets with formation of tubular lumina with abundant mucoid stroma (H & E,10X).

noted on the trunk and face. Common clinical presentations were swelling, ulcer and proliferative growth. All the tumors were well circumscribed lobular masses composed of clear cells and eosinophilic cells arranged in solid areas, tubule formations, with areas of cystic change. Cytoplasm of clear cells showed PAS positivity indicating presence of glycogen (Fig. 11). One of the tumor showed extensive squamous differentiation.

SCP was seen over back in a 38 year old male with epidermis showing papillomatosis and the cystic invaginations extending downward from epidermis lined by two layers of cells, the luminal cells being columnar showing apical snouting. Stroma of papillary core had dense lymphoplasmacytic infiltrate (Fig. 12).

Both hidradenocarcinomas mimicked metastatic adenocarcinoma histologically. One case was a recurrent swelling over the scalp in a 11 year old female (Figs. 13,14). Histologically the tumor

cells were arranged in lobules and sheets and in tubular formations filled with secretions. Cells displayed mild nuclear pleomorphism with clear cytoplasm and brisk mitosis. In both cases detailed histomorphological study and complete clinical diagnostic workup was done to exclude metastasis to the skin. Apocrine carcinoma of axilla had infiltrating margins with tumor cells displaying cytonuclear pleomorphism and brisk mitosis (Figs. 15). The cells were arranged in sheets, islands, tubules and singles. Luminal cells showed decapitation secretion. PAS-positive diastase-resistant material was identified in the cells.

Tumors with sebaceous gland differentiation

Sebaceoma presented as nodule over nose in 55 year old female and showed extensive squamous differentiation (Fig. 16). Both sebaceous carcinomas were noted over lower eyelid with suspected clinical diagnosis of chalazion and basal cell carcinoma.

Table 3: Tumors with sweat gland differentiation

| No of cases | Age/sex | Site | Clinical details | Histopathological Diagnosis |
|-------------|---------|-------------------------|---------------------------|-----------------------------|
| 1 | 75 y/F | Cheek | Ulcer- BCC | Hidradenoma |
| 2 | 18 y/F | Abdominal wall | Swelling | Hidradenoma |
| 3 | 60 y/F | Forehead | Nodular ulcer | Hidradenoma |
| 4 | 62 y/M | Scalp | Proliferative growth | Hidradenoma |
| 5 | 32 y/F | Neck | Swelling | Hidradenoma |
| 6 | 57 y/M | Back | Swelling - sebaceous cyst | Hidradenoma |
| 7 | 32 y/F | Scalp | Swelling -Adnexal tumor | Cylindroma |
| 8 | 38 y/F | Back | Papilloma | SCP |
| 9 | 27 y/F | Eyelid | Asymptomatic nodule | Syringoma |
| 10 | 48 y/M | Forehead | Swelling- Dermoid cyst | Chondroid syringoma |
| 11 | 62 y/M | External auditory canal | Nodule | Apocrine hidrocystoma |
| 12 | 11 y/F | Scalp | Recurrent swelling | Hidradenocarcinoma |
| 13 | 45 y/M | Abdominal wall | Nodulo-ulcerative growth | Hidradenocarcinoma |
| 14 | 60 y/M | Axilla | Cauliflower like growth | Apocrine carcinoma |

Table 4: A comparative analysis of Indian literature with regard to the line of differentiation of adnexal tumors

| Studies | Year | Total number of cases (N) | Tumors with Sebaceous differentiation | Tumors with Sweat gland differentiation | Tumors with Hair follicle differentiation |
|---------------------------|------|---------------------------|---------------------------------------|---|---|
| Vaishnavi and Dharkar [7] | 1974 | 48 | 03 | 38 | 07 |
| Kartha et al. [7] | 1980 | 83 | 03 | 45 | 35 |
| Reddy et al. [7] | 1982 | 85 | 18 | 54 | 13 |
| Solanki RL et al. [8,9] | 1989 | 94 | 22 | 50 | 22 |
| Nair PS [10] | 2008 | 33 | 02 | 19 | 12 |
| Saha A et al. [4] | 2011 | 23 | 03 | 13 | 07 |
| Jindal and Patel [11] | 2012 | 25 | 01 | 13 | 11 |
| Gayathri et al. [12] | 2012 | 29 | 03 | 15 | 11 |
| Radhika et al. [13] | 2013 | 35 | 7 | 17 | 11 |
| Pantola et al. [12] | 2013 | 70 | 4 | 42 | 24 |
| Sharma A et al. [14] | 2014 | 56 | 12 | 24 | 20 |
| Vani D et al. [15] | 2015 | 51 | 10 | 22 | 19 |
| Pujani M et al. [12] | 2016 | 25 | 4 | 14 | 7 |
| Present study | | 31 | 3 | 14 | 14 |




Fig. 11: Hidradenoma: solid areas composed of clear cells and eosinophilic cells (H&E,). Clear cells show PAS positivity (inset).



Fig. 14: Hidradenocarcinoma: Nodulo ulcerative growth measuring 7x6cm over abdominal wall




Fig. 12: Syringocystadenoma Papilliferum: Cystic invagination with by papillary projections lined by two rows of cells. Plasma cells seen in the core (H&E,10X).




Fig. 15: Apocrine carcinoma: Cauliflower like growth measuring 7x6 cm in axillary region




Fig. 13: Hidradenocarcinoma: Recurrent scalp swelling.




Fig. 16: Sebaceoma: well circumscribed mass with more than half of the cells being basaloid cells and few islands of mature sebaceous cells with extensive keratinization (H&E,10X).

Discussion

Only 31 cases seen in a 5 year period indicates that SATs are uncommon. A comparative analysis of many studies from Indian literature has been shown in Table IV with regard to the line of differentiation of SATs. In majority of Indian studies the most frequent line of differentiation encountered was sweat gland differentiation followed by hair follicle differentiation with the least frequent being sebaceous gland differentiation (Table 4). On the contrary present study showed equal occurrence of tumors of hair follicle and sweat gland differentiation (45%). In the present study 80.6% tumors were benign and 19.4% were malignant. Radhika et al., Reddy et al., Yaqoob N and Samaila have reported 77.14%, 69.41%, and 88.5% benign tumors respectively [7,13,15,16].

In our study SATs had a wide age range from 11 to 80 years with a female predominance. Female:male ratio was 1.8:1. Female : male ratio as observed by Nair PS and Saha et al. was 2.3:1 and 1.88:1 respectively [4,10]. Female predominance was noted in majority of Indian studies [6,13,14,17]. The higher frequency of SATs in females may be explained by the fact that they are cosmetically more conscious.

The predominance of SATs in head and neck region is a well documented fact supported by many studies which was also observed in our study (67.74%) [6,11,14,16,17]. This is accounted by the fact that this region is rich in adnexal units. In the present study, common SATs were PTT and hidradenoma. Radhika et al., Solanki RL et al., Reddy et al., Sharma et al. and Vani D et al. observed hidradenoma as the most common benign tumor [6,7,9,13,14].

PTT most commonly occur on the scalp and have a distinct predilection for women which concurs with our study [18]. Recently PTTs has been classified into benign, low grade, high grade and malignant tumors [18]. The usual clinical diagnosis of PTT is squamous cell carcinoma and sometimes PTTs may mimic SCC due to presence of cellular atypia [19]. Without paying closer attention to the trichilemmal type of keratinization, many of these lesions may be mistakenly diagnosed as SCC [20]. Presence of pre existing pilar cyst, good circumscription, trichilemmal keratinisation, absence of significant atypia, peripheral palisading and hyalinised basement membrane favour diagnosis of PTT [1]. Immunohistochemical positivity for CD34 and calretinin indicate the outer root sheath origin of PTT with low Ki 67 index [21]. Pilar cyst and PTT can

occur together and lesions intermediate between pilar cyst and PTT have been evidenced. The key factor in proliferation may be vascularisation of cyst epithelium possibly secondary to focal injury [22]. There was one such lesion in our study in a 60 year old male who presented with scalp swelling and showed presence of both these lesions. Malignant PTT is an uncommon histopathological entity. No absolute clinical criteria can distinguish a benign PTT from a malignant PTT. Hence the entire lesion must undergo histopathological examination [20].

Pilomatricoma was first described in 1880 by Malherbe and Chenantais as a tumor of the sebaceous glands [23]. Pilomatricoma, although considered a tumor of hair matrical differentiation, is now believed to be a complex panfollicular neoplasm, in that it can exhibit signs of differentiation of the entire follicle that is towards inner sheath, infundibulum and isthmic epithelium [23]. In 75-79% of the times the clinical diagnosis is incorrect at the time of initial consultation [24]. Usual clinical diagnosis include calcified sebaceous cyst, osteoma, neurofibroma etc [25]. The term "giant pilomatricoma" has been used for lesions greater than 5 cm in diameter, although the term is very subjective [24]. Older lesions become solid with predominance of shadow cells, keratin debris, multinucleated giant cells and dystrophic calcification [1]. Calcification occur frequently and osseous metaplasia of stroma is occasionally found [22]. In our study calcification was seen in 2 cases.

Trichoepithelioma is a major histological differential diagnosis for keratotic BCC. Presence of well-formed horn cysts, papillary-mesenchymal bodies, lack of high-grade atypia and mitoses and absent epidermal connection favor the diagnosis of trichoepithelioma. Artifactual retraction is uncommon in trichoepithelioma [1]. Staining of the outermost epithelial layer with bcl-2 has been seen in trichoepitheliomas. In contrast, basal cell carcinomas frequently stain diffusely [10].

Hidradenoma shows differentiation towards intraepidermal and intradermal eccrine structures ranging from the poral epithelium to the secretory segment [1]. Hidradenomas occur as a solitary nodule at all ages and have variable histomorphological patterns, reflected by the various terms used to describe this entity. Occasionally some features show overlapping with those of poromas. Histological variants include epidermoid, oncocytic and pigmented. Clear cell change and squamous metaplasia may be prominent. However, squamoid change does not

appear to denote a worse prognosis [26]. These can occasionally redevelop even after excision [27].

Apocrine hidrocystoma is regarded as cystic adenomatous proliferation of apocrine glands rather than retention cyst. It can be differentiated from eccrine hidrocystoma by presence of decapitation and PAS positivity [1,27]. The largest series of SCP (100 cases) have been reported by Helwig and Hackney [28]. Clinically usually presents as raised warty plaque [27]. Most lesions exhibit both apocrine and eccrine differentiation. Positive immunoreactivity with gross cystic disease fluid protein-15 and 24 (GCDFP-15 and 24) is seen, indicating tumor of apocrine origin but electron microscopic features show evidence of eccrine origin.¹ Recent studies show loss of heterozygosity at chromosome 9q22 in cases of SCP [10]. Chondroid Syringoma is so named due to a mixture of proliferating ductular eccrine epithelium and a characteristic chondroid stromal matrix. It occurs mainly on the head and neck. Ultrastructurally, cells within its chondroid matrix show features of both epithelial and mesenchymal cells raising the possibility that both cell lines may be the neoplastic progeny of pluripotent cells which is expressed early in the development of this tumor [15].

Usually Hidradenocarcinomas arise *de novo* and also arise in pre-existing hidradenoma. Microscopic distinction between the benign and malignant forms is usually made based on the extent of invasion, asymmetry at scanning magnification, brisk mitotic activity and degree of nuclear atypia [29]. The tumor cells stain positively for LMWK, and the ductal structures/luminal surfaces are highlighted by EMA and CEA [26]. Apocrine carcinoma is a rare and highly aggressive adenocarcinoma, mostly occurs in areas rich in apocrine glands, especially the axilla. The tumor cells are positive for LMWK, and the luminal cells are reactive for CEA, EMA and GCDFP-15. Positivity to androgen receptors and GCDFP-15 is considered helpful but negativity of these tumors to ER and PR is more relevant. The use of calponin and SMA as markers for the peripheral myoepithelium is helpful in delineating invasion [26].

Ocular sebaceous carcinomas are aggressive tumors and most commonly arise from meibomian and zeiss glands. Usual clinical diagnosis is chalazion. Presence of lymphovascular invasion, orbital extension and large size indicate poor prognosis. Tumor is positive for EMA and androgen receptor [27].

Malignant SATs often resemble their benign counterparts and attention has to be paid to features

such as large size, rapid increase in size, infiltrative growth pattern, nuclear atypia, brisk mitosis and necrosis [1]. Some primary malignant SATs like hidradenocarcinomas mimic metastatic moderately to poorly differentiated adenocarcinomas. In such cases detailed histomorphological features and complete clinical diagnostic workup are prerequisite to exclude metastasis to the skin. Two such tumors were present in our study. IHC may play an important role in distinguishing primary cutaneous adnexal carcinoma from metastatic carcinoma. P63 and CK 5/6 positivity favors a primary cutaneous adnexal carcinoma over a metastatic carcinoma [12].

The SATs encompassed a wide variety of clinical presentation, presenting as nodules, swelling and ulcerative growth with histologically distinct features. The most common clinical diagnosis for benign and malignant SATs were sebaceous cyst and squamous cell carcinoma respectively. The clinical appearance of appendageal tumor was mostly non-specific and diagnosis of SATs was not suspected clinically in most of the cases. Hence, the discrepancy between clinical and histopathological diagnosis was encountered. However, the location, single/multiple lesions, and their distribution can provide important diagnostic clues. In the present study, the concordance between clinical and histopathological diagnosis was found to be 6.45% approximately. Kamyab-Hesari et al. and Pujani M et al. found a higher (64% and 50%) concordance between clinical and histopathological diagnosis [12]. Sometimes more than one type of appendage give rise to combined tumors with varying degrees of maturation. However in our study, such combined tumors were not detected. Many a times it is only of academic interest to give a specific name for SATs because the surgeons pay more attention to the nature of tumor and its prognosis once it has been removed. Cytochemical stains such as PAS, mucicarmine, alcian blue, and reticulin may aid in establishing the diagnosis. The need for immunohistochemistry varies from case to case depending on histogenesis of SATs.

Conclusion

SATs are diverse group of neoplasms, with a nonspecific clinical appearance and their diagnosis primarily rely only on histopathology. However the location and distribution of lesions may provide diagnostic clues. Occasionally these tumors are difficult to classify and prognosticate. In our study the frequency of benign skin adnexal tumors was

more as compared to the malignant ones. SATs can occur anywhere in the body; however head and neck region constituted the most common site in our study. Most of the SATs occurred in younger age group but still showed a wide age variation. Most SATs are benign and local complete surgical excision is curative. Because these tumors are not frequently encountered, their diagnosis may be challenging even to an experienced pathologist. Dermatopathologists should employ a reliable criteria for accurate and reproducible diagnosis of SATs and make a foundation to determine the specific line of differentiation.

References

- Klein W, Chan E, Seykora JT. Tumors of the epidermal appendages. In: Elder DE, Elenitsas R, Johnson BL Jr, Murphy GF, editors. *Lever's Histopathology of the Skin*. 9th ed. Philadelphia: Lippincott Williams and Wilkins; 2005.pp.867-926.
- McCalmont TH. A call for logic in the classification of adnexal neoplasms. *The American Journal of Dermatopathology*. 1996;18(2):111-2.
- Headington JT. Tumours of the hair follicle. *American journal of pathology*. 1976;85(2):480-501.
- Saha A, Das NK, Gharami RC, Chowdhury SN, Datta PK. A clinico-histopathological study of appendageal skin tumors, affecting head and neck region in patients attending the dermatology OPD of a tertiary care centre in Eastern India. *Indian J Dermatol*. 2011;56:33-6.
- Alsaad KO, Obaidat NA, Ghazarian D. Skin adnexal neoplasms. Part 1. An approach to tumors of the pilosebaceous unit. *Journal of Clinical Pathology*. 2007;60(2):129-44.
- Vani D, Ashwini NS, Sandhya M, Dayananda TR, Bharathi.M. A 5 year histopathological study of skin adnexal tumors at a tertiary care hospital. *Journal of Dental and Medical Sciences*. 2015;14(4):1-5.
- Reddy MK, Veliath AJ, Nagarajan S and Aurora AL. A clinicopathological study of adnexal tumors of skin. *Indian journal of medical research*. 1982;75: 882-9.
- Solanki RL, Anand VK, Gaur SK, Arora HL and Gupta R. Neoplasms of hair follicle. *Indian journal of Dermatology Verereol Leprology*. 1989;55:33-7.
- Solanki RL, Anand VK. Neoplasms of sweat gland. *Indian J of Dermatol Venerol Leprol*. 1989;55:108-12.
- Nair PS. A clinicopathological study of skin appendageal tumours. *Indian J of Dermatol Venerol Leprol*. 2008;74:550.
- Jindal U and Patel R. Study of adnexal tumors of skin: A three year study of 25 cases. *Internet Journal of Pathology*. 2012;13(3).
- Pujani M, Madaan GB, Jairajpuri ZS, Jetley S, Hassan MJ and Khan S. Adnexal tumors of skin: An experience at tertiary care centre at Delhi. *Ann Med Health Sci Res*. 2016;6(5):280-5.
- Radhika K, Phaneendra BV, Rukmangadha N and Reddy MK. A biopsy confirmed skin adnexal tumours: experience at a tertiary care teaching hospital. *Journal of Scientific Research*. 2013;2:132-8.
- Sharma A, Paricharak DG, Nigam JS, Rewri S, Soni PB, Omhare A, et al. Histopathological study of skin adnexal tumours – Institutional study in South India. *Journal of Skin Cancer*. 2014;4:1-4.
- Yaqoob N. Ahmad Z. Muzaffar S, Gill MS, Soomro IN, Hasan SH. Spectrum of Cutaneous Appendage Tumors at Aga Khan University Hospital; *Journal of Pakistan Medical Association*. 2003;53(9):427-31.
- Samaila MO. Adnexal skin tumors in Zaria, Nigeria. *Annals of African Medicine*. 2008;7(1):6-10.
- Kanth GK, Vijayasree V, Madhavi M, Rao ST and Narayana SV. A histopathological spectrum of skin adnexal tumors. *Journal of Dental and Medical Sciences*. 2015;14(8):90-2.
- Ye J, Nappi O, Swanson PE, Patterson JW and Wick MR. Proliferating pilar tumors: A clinicopathologic study of 76 cases with a proposal for definition of benign and malignant variants. *Am J Clin Pathol*. 2004;122:566-74.
- Kumar P, Chatura KR, Haravi RM and Chandrasekhar HR. Proliferating trichilemmal cyst mimicking squamous-cell carcinoma. *Indian J Dermatol Verereol Leprol*. 2000;66:149-50.
- Ramaswamy AS, Manjunatha HK, Sunilkumar B and Arunkumar SP. Morphological Spectrum of Pilar Cysts. *N Am J Med Sci*. 2013 Feb;5(2):124-8.
- Gulati HK, Deshmukh SD, Anand M, Morale V, Pande DP, Jadhav SE. Low-grade malignant proliferating pilar tumor simulating a squamous-cell carcinoma in an elderly female: A case report and immunohistochemical study. *Int J Trichol*. 2011 Dec 18;3:98-101.
- Headington JT. Tumours of the hair follicle. *American journal of pathology*. 1976;85(2):480-501.
- Rajalakshmi T, Correa M and Simi CM. Pilomatricoma: A tumor with hidden depths. *Indian J of Dermatol Venerol Leprol*. 2010;76(5):543-6.
- Abdel- Gawad MM, Zaghoul AB and Yousef AS. Giant peduncular pilomatricoma (calcifying epithelioma of Malherbe): An interesting case. *The Gulf Journal of Dermatology and Venereology*. 2011;18(2):50-3.
- Guinot-Moya R, Valmaseda-Castellon E, Berini-Aytes L, Gay-Escoda C. Pilomatrixoma. Review of 205 cases. *Med Oral Patol Oral Cir Bucal*. 2011;16(4):552-5.
- Obaidat NA, Alsaad KO, and Ghazarian D. Skin adnexal neoplasms – Part 2: An approach to tumors of cutaneous sweat glands. *J Clin Pathol*. 2007;60(2):

- 145-59.
27. David W. Tumors of cutaneous appendages. In: Weedon David's Skin Pathology. 3rd ed. Churchill Livingstone; 2002.pp.859-916.
28. Mammino JJ and Vidmar DA. Syringocystadenoma papilliferum. International Journal of Dermatology 1991;30(11):835-6.
29. Gauerke S, Driscoll JJ. Hidradenoma carcinomas. Arch Pathol Lab Med. 2010;134:81-5.
-