

Clonidine as an Adjuvant to Ropivacaine in Subclavian Perivascular Approach to Brachial Plexus Block

Vaishali Waindeskar¹, Anuj Jain², Deepti Agrawal³, Charan Raj⁴

Author Affiliation: ¹Professor, ²Assistant Lecturer, ³Senior Resident, ⁴Junior Resident, Dept. of Anesthesia, All India Institute of Medical Sciences (AIIMS), Bhopal, Madhya Pradesh 462020, India.

Corresponding Author: Anuj Jain, Assistant Lecturer, Dept. of Anesthesia, All India Institute of Medical Sciences (AIIMS), Bhopal, Madhya Pradesh 462020, India.

E-mail: vaishaliwaind@gmail.com

Received on 08.08.2019,

Accepted on 09.11.2019

How to cite this article:

Vaishali Waindeskar, Anuj Jain, Deepti Agrawal *et al.* Clonidine as an Adjuvant to Ropivacaine in Subclavian Perivascular Approach to Brachial Plexus Block. Indian J Trauma Emerg Pediatr. 2019;11(4):105-108.

Abstract

Background and Aim: To evaluate the effectiveness of clonidine as adjuvant to ropivacaine in subclavian perivascular approach to brachial plexus block for prolonging the duration of sensory and motor blockade and analgesia also.

Methods: Patients were randomly divided into two groups. Patients in Group R (n = 30) were administered 25 ml of 0.75% Ropivacaine + 1 ml normal saline and Group RC (n = 30) were given 25 ml of 0.75% Ropivacaine + 1 ml 75 microgram of clonidine.

Results: The onset and the duration of sensory and motor block was significantly faster and prolonged in Group RC compared to Group R. Hemodynamic variables did not differ between groups in the post-operative period.

Conclusion: Clonidine 75 microgram in combination with 0.75% Ropivacaine was found to be hastening the onset of sensory and motor block and improved postoperative analgesia in brachial plexus block.

Keywords: Ropivacaine; Clonidine; Brachial plexus Block.

Introduction

Peripheral nerve blocks are gaining widespread popularity for perioperative pain management

because of their distinct advantages over general and central neuraxial anesthesia. Brachial plexus block is sole anesthesia for upper limb surgeries. In 1911 Kullen kampff introduced the classic supraclavicular approach of brachial plexus block. Winnie and Collins introduced the subclavian perivascular approach to brachial plexus block. Several techniques have been used to prolong the duration of regional anesthesia. Many experimental and clinical studies have shown that alpha-2 adrenergic agonists like clonidine were able to prolong sensory and motor blockade. This study involves the addition of an alpha-2 adrenergic agonist. Clonidine to ropivacaine solution in subclavian perivascular approach to brachial plexus block and the effects are evaluated and compared with ropivacaine alone.

Materials and Methods

A prospective double-blinded randomized control study conducted on 60 patients of ASA Grade I or II of either sex and age more than 19 undergoing upper limb surgery, in orthopedic theaters, under supraclavicular brachial plexus block performed by subclavian perivascular approach with nerve stimulator were included. Excluded from the study were patient refusal, local infections at the site of block,

history of drug allergy, coagulation abnormalities, significant systemic disorders, Alcohol/drug abuse pregnancy/lactating women, contraindicated to clonidine, Chronic analgesic therapy, peripheral neuropathy and very obese. Patients were preoperatively assessed and the procedure explained and written informed consent was obtained. They were randomly divided into two groups of 30 patients namely: Group R and Group RC. No premedication was given to the patients. Intravenous access was obtained. Anesthesia machine checked, resuscitative equipment and drugs were kept ready. Supraclavicular block was performed by subclavian perivascular approach with nerve stimulator. In Group R: Patients received supraclavicular block with 25 ml of 0.75% Ropivacaine + 1 ml normal saline. In Group RC: Patients received supraclavicular block with 25 ml of 0.75% Ropivacaine + 1 ml 75 microgram of clonidine. Care was taken so that the toxic doses of the local anesthetics were not exceeded according to the weight of the patients.

Parameters Observed are

1. Onset of Sensory block

Onset of Sensory block was taken as abolishment of pin prick pain over the distribution of ulnar and median and was assessed every minute after the performance of the block.

2. Onset of motor block

Onset of motor blockade was assessed every 3 minute after the block using four point scales

- 0 – Normal power
- 1 – Weakness but able to move arm
- 2 – Not able to move arm but the fingers
- 3 – Complete motor blockade

Attaining a score of 2 was considered as the onset of motor block.

3. Duration of surgery: Time taken by the surgeon to do the surgical procedure.

4. Duration of motor block: When (3) in the four point scale changes to (2) the motor blockade is said to reverse. The duration of motor block is noted from the time from scale (3) to scale (0).

5. Duration of sensory block: Taken from the time of onset of block to first complaint of pain sensation (vas score 1).

6. Duration of analgesia: Taken from the time of the onset of block to appearance of pain requiring first supplement analgesia (vas score more than 5).

7. Vital parameters: Pulse rate, blood pressure, saturation were monitored at 2nd, 5th, 10th, 15th minutes and then every 15 minutes for 2 hours then hourly for 6 hours then up to 24 hours.

8. Sedation score: Ramsay sedation score was employed.

- Anxious and Alert
- Conscious and Oriented
- Sedated, responding to verbal stimulus
- Responding to mild physical stimulus
- Responding to moderate and severe physical stimulus
- Not arousable.

9. Side effects: These noted are hypotension, bradycardia.

10. The pain was assessed using visual analogue scale having 10 cm length numbered from 0 to 10. Patient was explained about the visual analogue scale as: 0 – No pain and 10 the worst possible pain and was asked the score in visual analogue scale. The patients were observed every 30 minutes after the surgery is over till the motor block reverses and thereafter hourly for 6 hours; 2 hourly for next 6 hours and than 24 hours. Time at which VAS score is greater than 5 is noted and patient was given intramuscular NSAID.

11. Patients in whom the block was unsuccessful due to total failure of missed dermatomes which needed intravenous supplementation or general anesthesia were excluded from the study.

Results

The information collected regarding all the selected cases were recorded in a Master Chart. Data analysis was done with the help of computer using Epidemiological Information package (2008). Using this software range, frequencies, percentages, means, standard deviations, chi square and *p* values were calculated. Krukul Wallischi-square test was used to test the significance of difference between quantitative variables and Yate's test for qualitative variables. A *p* value less than 0.05 is taken to denote significant result.

Table 1: Demographic Data of Patients in Two Groups

Parameters Groups	R	RC
Number of patients	30	30
Average Age (years)	38.8 ± 13.6 yr	36.4 ± 11.4 yr
Weight (kg)	61.7 ± 4.9 kg	61.9 ± 4.6 kg
Gender (Male : Female)	17 : 13	24 : 06
ASA (1:2)	23:7	22:8
Duration of surgery in hours	2.1 ± 1.6 hr	1.9 ± 1.5 hr

There were no differences between the Clonidine and the control groups regarding age, sex, weight, ASA and duration of surgery in hours [Table 1].

It was observed in Table 2 that the sensory onset time was faster in Group RC (4.6 ± 0.5 min) than in Group R (8.9 ± 0.5 mins). Similarly, the onset of motor block in Groups R, RC was (10.5 ± 0.5 min and 7.5 ± 0.8 min) respectively. The mean duration of motor block was (7.2 ± 0.6 hours and 9.4 ± 0.6 hours) in Groups R and RC respectively. The mean duration of sensory block in hours was maximum in Group RC (11.3 ± 0.7 hours) compared to Group R (8.9 ± 0.6 hours). The mean duration of analgesia was (14.083 ± 0.617 hours) in Group RC followed by Group R (9.41 ± 0.602 hours) respectively.

Table 2: Sensory and Motor Block Onset, Duration Time and Duration of Analgesia in both Groups

	Group R Mean \pm SD	Group RC Mean \pm SD
<i>Onset of Sensory</i>		
Block (in minutes)	8.9 ± 0.5 min	4.6 ± 0.5 min
<i>Onset of Motor</i>		
Block (in minutes)	10.5 ± 0.5 min	7.5 ± 0.8 min
<i>Duration of Sensory</i>		
Block (in hours)	8.9 ± 0.6 hours	11.3 ± 0.7 hours
<i>Duration of Motor</i>		
Block (in hours)	7.2 ± 0.6 hours	9.4 ± 0.6 hours
<i>Duration of</i>		
Analgesia (in hours)	9.41 ± 0.74 hours	13.18 ± 0.76 hours
Sedation Score	2.0 ± 0.0	2.4 ± 0.5

Patients remained hemodynamically stable and also without any complications throughout perioperative period.

Discussion

Ropivacaine is structurally related to bupivacaine. Ropivacaine with its efficacy, reduced potential for cardiotoxicity and central nervous system toxicity, appears to be an important option for regional anesthesia and management of postoperative pain. Alpha-2 agonists like clonidine was introduced in the early 1960s as a nasal decongestant. There have been four proposed mechanisms for the action of clonidine in peripheral nerve blocks. These mechanisms are centrally mediated analgesia, alpha-2-adrenoreceptor mediated vasoconstriction, attenuation of inflammatory response and direct action on peripheral nerve.¹ In our study we used 75 microgram of clonidine after seeing various studies Saied *et al.*², Singelyn *et al.*³ and Eledjam *et al.*⁴

showing 75 microgram as effective dose. In our study we found that the onset and duration of sensory and motor block was significantly faster and prolonged respectively in patients who received a combination of clonidine and ropivacaine. Our results also showed that sensory block tended to last longer as compared to motor block. Addition of clonidine to local anesthetic ropivacaine solution showed prolonged analgesia significantly when compared to ropivacaine alone. Perineurally injected clonidine is thought to exert an analgesic effect through systemic absorption.⁵ These results correlate with studies conducted by Saied, Steyn *et al.*². Clonidine may produce sedation by acting on locus ceruleus of brainstem via action on alpha-2 adrenergic receptor.⁶ In Group RC since the sedation score was not more than 3, the respiratory function was not compromised. In this study, no significant difference was observed with respect to the pulse rate, systolic and diastolic blood pressure and saturation after administration of the block in Group R and Group RC. This could be attributed to the lower dose of clonidine used in our study compared to 300 microgram. In both groups there is no incidence of hypotension and bradycardia. This was consistent with the observation by EL Saied *et al.*², Casati A *et al.*⁷ and Eledjam JJ *et al.*⁴

Conclusion

The addition of the clonidine 75 microgram can be considered as a safe additive to local anesthetic solution in subclavian perivascular approach of brachial plexus block shows early onset and prolongs the duration of sensory and motor blockade and also duration of analgesia without any adverse effects.

References

1. Kalyani NP and Noopur DS *et al.* Clonidine as an adjuvant to ropivacaine-induced supraclavicular brachial plexus block for upper limb surgeries. *Journal of Anesthesiology Clinical Pharmacology*. 2015 Jul-Sep;31(3):365-69.
2. Saied EL, Steyn MP. *et al.* Clonidine prolongs the effect of Ropivacaine for Axillary brachial plexus blockade. *Canadian journal of anesthesia*. 2000;47:962-67.
3. Singelyn FJ, Dangoisse M. Adding clonidine to mepivacaine prolongs the duration of anesthesia and analgesia after axillary brachial plexus block. *Reg Anesth*. 1992 May-Jun;17(3):148-50.

4. Eledjam JJ *et al.* Brachial plexus block with bupivacaine; effects of added alpha adrenergic agonists: comparison between clonidine and epinephrine. *CJA*. 1991;38(7):870-5.
 5. Popping DM, Elian *et al.* Clonidine as an adjuvant to local anaesthetics for peripheral nerve and plexus block: Meta analysis of randomized trials. *Anesthesiology*. 2009 Aug;111(2):406-15.
 6. Rustagi P *et al.* A double blinded evaluation of adding clonidine. *Minerva Anaesthesiology* 2001 May;67(5):407-12.
 7. Casati A *et al.* Improving postoperative analgesia after axillary brachial plexus anesthesia with 0.75% ropivacaine. *Minerva Anesthesiol*. 2001 May;67(5):407-12.
-
-