

Evaluation of Pseudo Exfoliation Syndrome and Its Associated Intra Operative and Post Operative Complications in Cataract Surgery

Meena Lakshman*, Anand Reddi**

*Assistant Professor, Department of Ophthalmology, KMCT Medical College Hospital, Kozhikode, Kerala -673602. **Assistant Professor, Department of Ophthalmology, Kerala Medical College Hospital, Mangode, Palakkad Dt, Kerala -679503.

Abstract

Background: The association of pseudoexfoliation with blindness and aging has public health implication for India; this is especially so considering the burden of cataract with aging and the association of pseudoexfoliation with cataract as well as complications of cataract surgery. *Aim:* To clinically evaluate the cases of pseudoexfoliation syndrome and its associated intra operative and post operative complications during cataract surgery. *Methods:* A descriptive study based on probability sampling of patients having pseudoexfoliation syndrome admitted for cataract surgery in a tertiary care hospital was carried out spread over 3 years. A total of 220 eyes of 125 cataract patients aged 50 years or more were included in the study. The data were analyzed through SPSS version 15.0. *Results:* A total of 125 patients having Pseudoexfoliation were included in the study. Vitreal prolapse in 21 (10.5%) and posterior capsular rupture in 18 (9%) patients were the most common complications seen in Pseudoexfoliation. Damage to sphincter pupillae in 16 (8%), iridodialysis in 2 (1%), decentration of IOL in 8 (4%) and hyphaema in 2 (1%) patients was seen. Zonular dialysis in 8 (4%), retained lens matter in 12 (6%) and lens dislocation was seen in 6 (3%) patients. *Conclusion:* Patients with pseudoexfoliation are at increased risk for development of complications. Ophthalmologists should stress to increase awareness among general public for the proper diagnosis and convince patients for proper and regular follow up visits to the hospital.

Keywords: Pseudoexfoliation; Cataract; Complications; Intraocular pressure.

Introduction

Pseudoexfoliation (PXF) is a senile condition, more common in females, familial and seems to be genetically inherited disease. A high risk of developing pseudoexfoliation syndrome and pseudoexfoliation glaucoma is confirmed by mutations in LOXL1 gene at locus 15q22, coding for elastic fibre components of the extracellular matrix. A grey-white fibrillary extracellular material composed of a protein core surrounded by glycosaminoglycans is produced by abnormal basement membranes of aging epithelial cells in the trabeculum, equatorial lens capsule, iris and ciliary body. The material is then deposited on the anterior lens capsule, zonules, ciliary body, iris trabeculum, anterior vitreous face and conjunctiva [1].

Ultra structural studies performed on eyes during autopsy suggest that pseudoexfoliation syndrome is a systemic disorder. Pseudoexfoliation material has been found in a number of organs, which includes skin, lungs, gall bladder, liver, myocardium, kidney, bladder and meninges [2].

Pseudoexfoliation is rarely seen before the age of 40, and its prevalence increases markedly with age. Although it occurs virtually in every area of the world, a considerable racial variation exists. Framingham study showed that the prevalence of

Reprint Request: Meena Lakshman, Assistant Professor, Department of Ophthalmology, KMCT Medical College and Hospital, Kozhikode, Kerala -673602.
E-mail: thesisum@gmail.com

Received on 12.12.2016, Accepted on 13.12.2016

pseudoexfoliation was 1.8 %. In another study of subjects over 60 yrs in various ethnicities, prevalence rates ranging from 0 % in Greenland Eskimos to 21 % in Icelanders were noted [3].

Pseudoexfoliation is a known risk factor for developing cataracts. Cataract surgery in eyes with pseudoexfoliation has higher incidence of operative complications like posterior capsular rupture, zonular dialysis, vitreous loss and intra ocular bleeding. Post-operatively, these patients are at greater risk of developing immediate elevation of IOP and inflammation. Posterior capsular opacification and intraocular lens decentration are more common in patients with pseudoexfoliation in post-operative period [4].

The association of pseudoexfoliation with blindness and aging has public health implication for India; this is especially so considering the burden of cataract with aging and the association of pseudoexfoliation with cataract as well as complications of cataract surgery [5].

Our hospital being a tertiary care hospital, we get fair number of cases of pseudoexfoliation syndrome, hence in the present study every effort is made to clinically evaluate pseudoexfoliation syndrome and its associated intra operative and post operative complication of cataract surgery. In this background it was decided to take up this study.

Method

This study was carried out from January 2013 to December 2015 at Departments of Ophthalmology, KMCT Medical College and Hospital, Calicut. Convenience (non probability sampling) was used for this prospective and descriptive study.

Patients with developmental cataract, traumatic cataract, complicated cataract, high myopia, any other ocular disease and previous surgery were excluded. Patients were admitted to the Ophthalmology wards of the hospitals.

Written consent of all the patients included in the study was taken after fully explaining the procedure and purpose of the study to the patients. A detailed proforma was devised containing all essential details for each individual. A complete ophthalmic history was taken. The patients were asked about their name, age, sex, occupation and address. The questions and inquiries about the complaints were made which included dimness of vision, headache, pain in the eyes, watering of the eyes, discharge from the eyes, sticking of the lids, photophobia, history of injury

and past history.

A thorough examination including visual acuity, anterior segment, posterior segment and measurement of intraocular pressure (IOL) was performed before the cataract surgery. Intraocular pressure was recorded with applanation tonometer.

The diameter of pupil of each patient was measured. Intra-operative maximum pupillary dilatation was obtained and its size measured. This pupil size was graded as poor, fair and satisfactory/good. Poor pupillary dilatation meant 2–4 mm, fair pupillary dilatation meant 5–6 mm, satisfactory/good pupillary dilatation meant 7–9 mm.

Standard extracapsular cataract extraction with PMMA intraocular lens implantation was done. Topical antibiotic ointment was applied. Eye was padded and patients were put on oral analgesics. The patients were examined with slit lamp on 1st post operative day. The follow up of the patients was carried out on 8th day, 4 weeks, 8 weeks and finally 12 weeks. All the patients received oral analgesics and topical Polymyxin, Neomycin, Dexamethasone and Ofloxacin/Ciprofloxacin. The data were entered and analysed using SPSS version 15.0. Analysis was done for quantitative and qualitative measures.

Results

Two hundred (220) cataract patients with pseudoexfoliation syndrome were included in this study. Patients lost during follow up were excluded from the study. The ages of these 220 patients ranged from 50 years to 87 years with 12% patients in age group of 80–89, 35% patients in the age group of 70–79, 37% patients in age group of 60–69 years and 16% patients in age group of 50–59 years. The distribution by gender shows that 84% of patients were males, while the remaining 16% of the patients were females; the male and female ratio was 5.25:1. Distribution of frequency and percentage of right and left eyes shows that 12% were right, 10% were left and 78% were bilateral.

Preoperative features show that 22% patients had zonular fragility, 6% had iridodonesis and 14% had phacodonesis. In 6% patients, pigment dispersion was seen, out of these 4% was present on lens and 2% was present on cornea. Subluxation of lens was seen in 4% patients. Mean IOP was 19.6 ± 2.30 mmHg. Intraoperative pupillary diameter measurements showed that poor pupillary dilatation was seen in 48% patients, fair pupillary dilatation was seen in 42% patients while satisfactory/good pupillary

dilatation was seen in 10% patients.

Different procedures were conducted during cataract surgery. Peripheral iridectomy was done in all the cases. Sphincterotomy was done in 46% patients with poor pupillary dilatation. Fifty-four (54%) patients received posterior chamber IOL and 92 (46%) had planned cataract extraction. Table 1

shows distribution of surgical complications during cataract surgery. A total of 93 (46.5%) complications were recorded. Routine follow up visits were done for 12 weeks in all the patients. On each follow up, visual acuity and slit lamp examination was done. Final visual acuity was recorded after 12 weeks.

Table 1: Surgical complications during Cataract surgery(n=93)

Surgical complications	Pseudoexfoliation patients(n=220)
Damage to sphincter pupillae	16(8%)
Iridodialysis	2(1%)
Lens dislocation	6(3%)
Posterior capsular rupture	18(9%)
Vitreous prolapse	21(10.5%)
Retained lens material	12(6%)
Decentration of IOL	8(4%)
Zonular dialysis	8(4%)
Hyphaema	2(1%)
Total:	93(46.5%)

Discussion

This study explored the peroperative outcomes and complications of cataract patients with pseudoexfoliation in tertiary care setting. Pseudoexfoliation was more commonly seen in males as compared to females in the present study, i.e., 84% in males and 16% in females. This ratio is approximately 5:1. More frequent occurrence of pseudoexfoliation in males can be explained by two factors. The first factor is that most of the patients undergoing cataract surgery are males. Secondly, males are more commonly involved in outdoor activities while most of the females are restricted to homes as traditional housewives. This particular aspect of restriction to houses partly explains environmental factors, as there are fewer risks for development of pseudoexfoliation. This is consistent with the finding of other studies [6,7]. Pseudoexfoliation can be unilateral or bilateral. Our study describes unilateral cases to be 22%.

Pseudoexfoliation is associated with constricted pupil. Adequate pupillary dilatation is necessary for standard extracapsular extraction. Pupillary dilatation is obtained by topical tropicamide. In the present study, pupil size was recorded after installation of tropicamide at 10 minutes interval for half an hour. Poor pupillary dilatation was seen in 48% of the patients, adequate in 42% of the patients and good in 10% of the patients with pseudoexfoliation. In another study, 68.75% of pseudoexfoliation patients had poor to moderate pupillary dilatation [7]. These results indicate that good/adequate pupil dilatation for standard

extracapsular cataract extraction is more difficult to be obtained in patients with pseudoexfoliation.

Constricted pupil exposes the patient to more complications [8]. To obtain adequate dilatation, different methods are in use. Bimanual stretching is one of the least time consuming methods [9] and was used in our study. This method is more convenient and cheaper but may lead to iris sphincter damage. Sphincter damage was seen in 8% of the patients. This damage leads to anisocoria postoperatively but is of help to obtain pupil dilatation thus making anterior capsulotomy more convenient and of appropriate size. Proper anterior capsulotomy and adequate dilatation of pupil help in uneventful surgery. The better option is sphincterectomy in patients who are susceptible to sphincter damage.

Sphincterectomy thus provides more controlled enlargement of pupil and at the same time better site can be selected.

Iridodialysis occurs intraoperatively as a result of the manipulation of intraocular tissues. It is one of the established, although rare, complications of cataract surgery. The patients with pseudoexfoliation syndrome are more prone to have complications as compared to patients without pseudoexfoliation [10]. In this study, only two patients had this complication. The main reason is inappropriate handling of intraocular tissues rather than pseudoexfoliation. These patients had only mild iridodialysis, which was optical as well as cosmetically insignificant. That is why no significant surgical intervention was done.

Pseudoexfoliation is a source of severe complications in cataract surgery [11]. The increased

frequency of intraoperative complications stem from zonular weakness rather than capsular tears. Zonular weakness poses different problems that are zonular dialysis, posterior dislocation of crystalline lens material [12] and delayed spontaneous dislocation of the intraocular lens and capsule [13]. In the present study, 8 patients (4%) had zonular dialysis. Experienced surgeons do have complications but their frequency and extent of damage is far less as compared to juniors or residents.

Posterior capsular rupture impacts the patients in terms of additional surgical procedures, additional topical and oral medications, and number and duration of follow up reviews [14]. In the present study, 18 patients (9%) had posterior capsular rupture. The risk factors for the development of this complication are quite extensively discussed, keeping in mind its effects on visual outcome. Even racial differences although not significant have been considered [15].

Our study describes vitreous loss rate as 10.5%. Measures were taken to clear the vitreous from the anterior chamber and wound edge and no postoperative complications occurred. Intraocular lens was not implanted in any patient who had posterior capsular rupture, vitreous loss or zonular dialysis. Posterior capsular tears, zonular dialysis and vitreous loss are reported to be more frequent in pseudoexfoliation syndrome than in otherwise normal subjects and this study confirms the findings [16,17]. Retained lens material does not necessarily require surgical intervention but may incite a significant postoperative inflammation that may be difficult to distinguish from microbial endophthalmitis. Present study describes 12 patients who had retained lens material. All of them had retained little cortical material, which is better tolerated and was having no optical effects due to inferior position in anterior chamber; that is why, repeat surgery was not done. Treatment with cycloplegics and topical steroids allows for the gradual dissolution of the retained lens material. Patients with pseudoexfoliation may be at risk for delayed spontaneous dislocation of intraocular lens within the capsular bag after uncomplicated surgery. There may be tilting of the intraocular lens in less severe cases. Keeping in mind this complication, it is better to re-evaluate surgical consideration for cataract removal. Better option is to implant the intraocular lens in the ciliary sulcus. Hyphema in the immediate postoperative period usually originates from the incision or iris. Only 2 patients with pseudoexfoliation had this complication. It was mild and not associated with vitreous loss. No intervention was

done and it did resolve spontaneously.

Conclusion

Cataract surgery, like any surgical procedure, has associated complications. Patients with pseudoexfoliation are at increased risk for development of complications. Early diagnosis, detailed examination, knowledge of the complications, ability to manage these complications, use of viscoelastic material and surgery by experienced surgeon are keys of success.

References

- 1 Kanski JJ, Bowling Brad. Clinical ophthalmology A systemic approach. 7th ed. China: Elsevier Saunders; 2011.p.355-56.
- 2 Lattners W J, Samuelson T W. Pseudoexfoliation glaucoma. In: Yanoff M, Duker J S. Editors: Ophthalmology. 2nd ed. Vol.2. New Delhi: Mosby; 2006.p.1499-1503.
- 3 Shazly A T, Farrag A N, Kamel A, Al-Hussaini A K. Prevalence of Pseudo exfoliation Syndrome and Pseudo exfoliation Glaucoma in upper Egypt. BMC Ophthalmology. 27 June 2011(6 January 2011); 11:(5). Available at <http://www.biocentral.com/1471-2415/11/18>.
- 4 Naseem A, Khan S, Khan M N, Muhammad S. Cataract Surgery in Patients with Pseudo exfoliation. Pak J Ophthalmol. 2007; 23(3):155-160.
- 5 Thomas et al. Pseudo exfoliation in southern India. IOVS. 2005; 46(4):1170-1175.
- 6 Naeem S. Incidence, age of presentation and lenticular changes in exfoliation syndrome. [Dissertation]. Karachi: College of Physicians & Surgeons; 1997.p.149.
- 7 Naseem A. Cataract surgery in patients with pseudoexfoliation. [Dissertation]. Karachi: College of Physicians & Surgeons; 2002.p.111.
- 8 Khanzada AM. Exfoliation syndrome in Pakistan. Pak J Ophthalmol 1986; 2:7-9.
- 9 Katsimpris JM, Petropoulos IK, Apostolakis K, Feretis D. Comparing phacoemulsification and extracapsular cataract extraction in eyes with pseudoexfoliation syndrome, small pupil, and phacodonesis. Klin Monatsbl Augenheilkd 2004; 221(5):328-33.
- 10 Akman A, Yilmaz G, Oto S, Akova YA. Comparison of various pupil dilatation methods for phacoemulsification in eyes with a small pupil secondary to pseudoexfoliation. Ophthalmology 2004; 111:1693-8.
- 11 Parmar P, Salman A. Anterior "lenticonus" in

- exfoliation syndrome. *Indian J Ophthalmol.* 2005; 53 (3):193-4.
12. Potocky M. The pseudoexfoliative syndrome and its significance in cataract surgery *Cesk Slov Oftalmol.* 2003; 59(6):422-32. [Article in Slovak].
 13. Dickinson PJ, Greven CM. Avoiding and managing the dislocated crystalline lens. *Compr Ophthalmol Update.* 2006; 7(3):113-27.
 14. Hohn S, Spraul CW, Buchwald HJ, Lang GK. Spontaneous dislocation of intraocular lens with capsule as a late complication of cataract surgery in patients with pseudoexfoliation syndrome. *Klin Monatsbl Augenheilkd.* 2004; 221(4):273-6. [Article in German].
 15. Ang GS, Whyte IF. Effect and outcomes of posterior capsule rupture in a district general hospital setting. *J Cataract Refract Surg.* 2006; 32(4):623-7.
 16. Chan FM, Mathur R, Ku JJ, Chen C, Chan SP, Yong VS, et al. Rates of posterior capsule rupture during cataract surgery among different races in singapore. *Ann Acad Med Singapore.* 2006; 35(10):698-700.
 17. Susic N, Brajkovic J. The prevalence of pseudoexfoliation syndrome in patients admitted for cataract surgery to the department of ophthalmology, Sibenik General Hospital. *Acta Med Croatica.* 2006; 60(2):121-4.
-

Special Note!

Please note that our all Customers, Advertisers, Authors, Editorial Board Members and Editor-in-chief are advised to pay any type of charges against Article Processing, Editorial Board Membership Fees, Postage & Handling Charges of author copy, Purchase of Subscription, Single issue Purchase and Advertisement in any Journal directly to Red Flower Publication Pvt. Ltd. Nobody is authorized to collect the payment on behalf of Red Flower Publication Pvt. Ltd. and company is not responsible of respective services ordered for.