

Screening for Gastrointestinal parasites in Murrah calves of Buffalo Research Station of West Godavari Region of Andhra Pradesh

G Deepika Kumari¹, K Ananda Rao²

Authors Affiliation

¹Assistant Professor, Scientist, ²Principal, Scientist and Head, Buffalo Research Station, Venkatarammanagudem, Andhra Pradesh 534211, India.

Corresponding Affiliation

G Deepika Kumari, Assistant Professor, Scientist, Buffalo Research Station, Venkatarammanagudem, Andhra Pradesh 534211, India.

Email: deepu.angrau@gmail.com

Abstract

The present study was aimed to screen the presence of gastrointestinal endoparasites in Murrah buffaloes calves reared in the Buffalo Research Station (BRS), Venkatarammannagudem of Andhra Pradesh, India. A total of 44 dung samples were collected during the month of January 2021, from calves of different age groups ranging from 15 days to one year. Microscopic screening of the dung samples revealed presence of Buxtonella sulcata cysts and trophozoite forms in higher number and few helminthic infections of Toxocara vitulorum and Strongyle eggs. An overall incidence of the gastrointestinal parasites was found to be 43.18%, among which the incidence of Buxtonella sulcata was higher (36.36%) when compared to other endoparasites (6.81%). From the above study, it could be summarised that Buxtonella sulcata is a common inhabitant of gastrointestinal tract of buffaloes and responsible for diarrhoea in association with other helminths or mixed infections.

Keywords: Bubalus bubalis; Endoparasites; Buxtonella sulcata; Andhra Pradesh.

How to cite this article:

G Deepika Kumari, K. Ananda Rao/Screening for Gastrointestinal parasites in Murrah calves of Buffalo Research Station of West Godavari Region of Andhra Pradesh 2021;7(1):19-22.

Introduction

Andhra Pradesh has a wide coastal region and is the best potential source for the extensive maintenance of buffaloes. Andhra Pradesh holds nearly 10.6 million buffaloes and positioned 4th in India for buffalo milk production constituting nearly 7.4 million metric tonnes (Sivaji et al., 2018). Presently the Buffalo research station, recorded a total milk production of 99,221.5 kg from the milch animals during the year 2020-21. Murrah buffaloes are good high milch yielding animals with nearly 1360 to 2270 kg per lactation (Suresh, 2013). Parasites of helminthic origin cause hazardous economic losses to the farmers so proper safe guarding of the health of the buffaloes from the

various factors especially the gastrointestinal parasites should be done regularly (Saha et al., 2013). Gastrointestinal parasites mainly include Buxtonella sulcata Fasciola gigantica, Strongyloides Paramphistomum cervi, Schistosoma nasale, S. indicum, S. spindale, Toxocara vitulorum, papillosus, Strongyle group of worms, Eimeria sp. etc. and play a vital role in the of degradation the health of the domestic animals finally leading to severe economic losses (Grisi et al., 2014). Buxtonella sulcata a general inhabitant of gastrointestinal tract of cattle appears morphologically similar to Balantidium coli present in pigs. The severity of the endoparasitic infection in

the calves leads to gastrointestinal disturbances like white scours, indigestion, malabsorption of minerals and vitamins in the gastrointestinal tract, reduced appetite, and discoloration in the haircoat. (Hamid et al., 2016). The presence of endoparasites in a farm may become a predisposing factor for the spread of infection among the healthy animals. Individual animal's health performance is very important in assessing the farm status. The present study was aimed for a routine screening of the dung samples to detect for the presence of any gastrointestinal infections in buffalo calves.

Materials and Methods

Study Area: Buffalo research station is located in Venkataramannagudem of Tadepalligudem mandal *, West Godavari District, Andhra Pradesh, India. The geographical coordinates of Venkataramannagudem are 17.4° 48' 52.2864" N and 78.48° 31' 35.6048" E.

Collection and microscopic examination of dung samples.

A total of forty four samples (n=44) were collected for the present study as given in table 1.

Table 1: Collection of Dung samples.

Category	Calves (Below 3 months)	Calves (3 to 6 months)	Calves (6 months to 1 year)	Total no. of samples
Number of calves	13	9	22	44

The dung samples were collected from the rectum in the early hours of dawn in sterile plastic containers and stored at a refrigeration temperature of 4°C until the samples were processed for microscopic examination using binocular microscope. Both direct smear and sedimentation methods (Soulsby, 1982) were followed for the identification of parasites eggs, cysts and trophozoite forms.

Direct smear examination was performed by placing a little amount of dung sample on a clean glass slide and equal quantity of water was added and mixed thoroughly with wooden stick and a thin smear was prepared and observed under 10X and 40X objective for better clarity under microscope. Sedimentation method required nearly 1-2 gm of fresh dung sample emulsified in 10 ml of water and later strained and centrifuged at 2000 rpm for 3-5 minutes.

A drop of sediment was placed at centre of slide and cover slip was placed with care avoiding air bubbles and observed under low power and high power objectives. The microscopic measurements were made for the parasitic eggs, protozoan cysts and trophozoites using the ocular micrometer.

Results and Discussion

On thorough examination of the dung samples employing both direct smear and sedimentation techniques, the total endoparasitic infections in the Murrah calf buffaloes noted the highest presence of *Buxtonella sulcata* (36.36 %) (fig. 1), followed by Strongyle eggs (4.54 %) (fig. 2) and *Toxocara vitulorum* eggs (2.27 %) (fig. 3). The results of the detected endoparasites were given based on animals grouped as age wise in Table 2.

Table 2: Age wise incidence of endoparasites in Murrah calves.

Age	No. of samples screened	Parasites recorded	No. of positive samples	Incidence percentage
Calves (Below 3 months)	13	<i>Buxtonella sulcata</i> cysts & Trophozoites	6	46.15 %
		Strongyle eggs	1	7.69 %
		<i>Toxocara vitulorum</i> eggs	1	7.69 %
Calves (3 -6 months)	9	<i>Buxtonella sulcata</i> cysts	4	44.44 %
Calves (6-1year)	22	<i>Buxtonella sulcata</i> cysts	6	27.27 %
		Strongyle eggs	1	4.54 %

In the diarrhoeic buffalo calf dung samples, *Buxtonella sulcata* cysts and *Toxocara vitulorum* were identified. Results of the present study revealed that *Buxtonella sulcata* cysts were commonly observed in the dung samples of healthy buffaloes but not as heavy infestation. Earlier reports of *Buxtonella sulcata* incidence in India, Karnataka (Mamatha and Placid, 2006), Jammu (Ganai et al.2015) and abroad, Iraq (Aayiz, 2005), Nepal (Adhikari et al. 2015) recorded a variation of 21 to 71% (Biswas et al., 2014).

Different Gastrointestinal endoparasites found in the feces of Murrah buffalo calves.






Fig. 1: Buxtonella sulcata cyst and Buxtonella sulcata Trophozoite.



Fig. 2: Strongyle eggs.




Fig. 3: Toxocara vitulorum eggs.

Buxtonella sulcata, is a ciliated protozoan is a common inhabitant of the gastrointestinal tract of buffaloes and is usually non-pathogenic in nature. *Buxtonella sulcata* acts as a commensal in the intestines of the calves and aids in the digestion of plant matter (Tomczuk et al., 2005; Kumar et al., 2017). It was also reported by few authors the pathogenic nature of *Buxtonella sulcata* (Roy et al., 2011) in causing chronic diarrhoea. Along with *Buxtonella sulcata*, lesser percentage of other parasitic eggs like *Toxocara vitulorum* and strongyles were detected but the infection is very meagre. The reason for less parasitic infection in the farm might be due to clean animal husbandry and hygienic managerial practices maintained in the farm and also the environmental conditions. Kumar et al. 2017 also reported less incidence of parasitic infection internally due to better hygienic practices of Jaffarbadi buffaloes maintained in Gujarat farm. The incidence of *Buxtonella sulcata* in Murrah buffaloes with respect to age was slightly higher in calves when compared to adult buffaloes. The findings were in accordance to Ganai et al. 2015 who reported that *Buxtonella sulcata* was common in young animals but contradictory to Kumar et al. 2017 who reported that the incidence was higher in older animals. The varying reasons could be due to the climatic conditions prevailing in the geographical region of farm, besides environmental conditions, care and managerial practices followed in the maintenance of farm also play a key role for the growth and incidence of the parasites.

The infected calves with *Toxocara vitulorum* were treated with Levamisole hydrochloride @7.5kg/body weight and those affected with *Strongyloides* were treated with Fenbendazole were treated with @ 5kg/body weight. As *Buxtonella sulcata* is a common inhabitant general deworming with albendazole was carried out for all the calves in the herd and thorough washing of the water troughs maintained in the farm was done regularly.

In the present study, from the observations it can be summarized that the incidence of ciliated protozoa *Buxtonella sulcata* was revealed but not to level of causing chronic diarrhoea except in case of mixed infection with causing diarrhoea in calves. Other helminthic infections like *Toxocara vitulorum* and strongyle eggs were very few and thus it could be concluded that the care and managerial practices carried out in the organised farm were effective in safeguarding and controlling the endoparasites in Murrah buffaloes.

Acknowledgements

The authors are grateful to Sri Venkateswara Veterinary University, Tirupati for providing the facilities for carrying out the work.

References

1. Aayiz N. Diagnostic study for cow infection with *Buxtonella sulcata* in Iraq. *Al qudissiyha Journal of Veterinary Science*. 2005; 4:53-56.
2. Adhikari BB, Rana HB, Sultana KMI, Devakota B, Nakao T, Kobayashi K, Sato H, Dhakal IP et al. Prevalence of *Buxtonella sulcata* in water buffaloes and cows in Chitwan Valley, Southern Nepal. *Jpn. J.Vet.Parasitol*. 2013;12: 55-60
3. Biswas H, Dey AR, Begum N, Das PM et al. Epidemiological aspects of gastrointestinal parasites in buffalo in Bhola, Bangladesh. *Indian J Anim Sci*, 2014; 84:3.
4. Ganai A, Parveen S, Kaur D, Katoch R, Yadav A, Godara R, Ahamed I et al. Incidence of *Buxtonella sulcata* in bovines in RS Pura, Jammu. *Journal of parasitic diseases*, 2015. 39(3), 446-447.
5. Grisi LRC, Leite JRD, Martins S, Barros ATMD, Andreotti R, Cañado PHD, Villela HS et al. Reassessment of the potential economic impact of cattle parasites in Brazil. *Revista Brasileira de Parasitologia Veterinária*, 2014, 23(2), 150-156.
6. Hamid PH, Kristianingrum YP, Prastowo S et al. Gastrointestinal parasites of cattle in Central Java. *American Journal of Veterinary and Animal Sciences* 2016.
7. Kumar BB, Maharana R, Prashad A, Joseph JP, Patel BR. et al. Incidence of *Buxtonella sulcata* in Jaffarbadi buffaloes of south-western Gujarat, India. *Buffalo Bulletin*, 2017; 36(4), 623-628.
8. Mamatha GS, Placid EDS. Gastrointestinal parasitism in cattle and buffalo in and around Bangalore. *J.Vet.Parasitol*. 2006; 20: 163-165.
9. Roy BC, Mondal MMH, Talukde MH, Majumder S et al. Prevalence of *Balantidium coli* in Buffaloes at different areas of Mymen singh. *Journal of the Bangladesh Agricultural University*, 2011; 9(452-2016-35702), 67-72.
10. Saha SS, Bhowmik DR, Chowdhary MMR et al. Prevalence of gastrointestinal helminthes in buffaloes in Barisal district of Bangladesh. *Bangladesh Journal of Veterinary Medicine* 2013;11(2): 131-135.
11. Sivaji DV, Natchimuth K, Ramkumar S et al. Sustainability of buffalo farming in milk shed areas of Andhra Pradesh. *Bull. Env. Pharmacol Life Sci*, 2018; Vol.7(7) 37-41.
12. Soulsby E.J.L. . Helminths, arthropods and protozoa of domesticated animals, 1982, 7th Edn.ELBS and Bailliere Tindal, London.
13. Suresh Babu D.. Production Performance of Murrah Buffaloes under organized dairy farm production system in west Godavari district of Andhra Pradesh. *Indian Journal of Applied Research*. 2013;Vol.3 Issue 8.
14. Tomczuk KL, Kurek M, Stec Studzinska, Mochol J et al. Incidence and clinical ciliate *Buxtonella sulcata* infection in cattle. *B.Vet.I Pulawy* 2005; 49: 29-33.



SUBSCRIPTION FORM

I want to renew/subscribe international class journal "Indian Journal of Agriculture Business" of Red Flower Publication Pvt. Ltd.

Subscription Rates:

- Institutional: INR 6000 / USD 469

Name and complete address (in capitals): _____

Payment detail:

Online payment link: <http://rfppl.co.in/payment.php?mid=15>

Cheque/DD: Please send the US dollar check from outside India and INR check from India made payable to 'Red Flower Publication Private Limited'. Drawn on Delhi branch.

Wire transfer/NEFT/RTGS:

Complete Bank Account No. 604320110000467

Beneficiary Name: Red Flower Publication Pvt. Ltd.

Bank & Branch Name: Bank of India; Mayur Vihar

MICR Code: 110013045

Branch Code: 6043

IFSC Code: BKID0006043 (used for RTGS and NEFT transactions)

Swift Code: BKIDINBBDOS

Term and condition for supply of journals

1. Advance payment required by Demand Draft payable to **Red Flower Publication Pvt. Ltd.** payable at **Delhi**.
2. Cancellation not allowed except for duplicate payment.
3. Agents allowed 12.5% discount.
4. Claim must be made within six months from issue date.

Mail all orders to

Subscription and Marketing Manager

Red Flower Publication Pvt. Ltd.

48/41-42, DSIDC, Pocket-II

Mayur Vihar Phase-I

Delhi - 110 091(India).

Phone: 91-11-79695648, 22754205, 22756995, Cell: +91-9821671871

E-mail: sales@rfppl.co.in

REDKART.NET

(A product of Red Flower Publication (P) Limited)
(Publications available for purchase: Journals, Books, Articles and Single issues)
(Date range: 1967 to till date)

The Red Kart is an e-commerce and is a product of Red Flower Publication (P) Limited. It covers a broad range of journals, Books, Articles, Single issues (Print & Online-PDF) in English and Hindi languages. All these publications are in stock for immediate shipping and online access in case of online.

Benefits of shopping online are better than conventional way of buying.

1. Convenience.
2. Better prices.
3. More variety.
4. Fewer expenses.
5. No crowds.
6. Less compulsive shopping.
7. Buying old or unused items at lower prices.
8. Discreet purchases are easier.

URL: www.redkart.net