

Therapeutic Efficacies of Dexmedetomidine and Tramadol on Post Subarachnoid Block Shivering: A Prospective Study

Vishal Joshi¹, Vijay Goswami²

^{1,2}Assistant Professor, Department of Anaesthesia, Gujarat Adani Institute of Medical Science, Bhuj, Gujarat 370001, India.

Abstract

Background and Aim: Shivering is a physiological response to core hypothermia in an attempt to raise the metabolic heat production. Non-pharmacological methods using equipment to maintain normal temperature of the body are effective but expensive and lack practicality, while the pharmacological methods are simple, cost-effective and easy to implement. Present study aimed to compare therapeutic efficacy of Dexmedetomidine and Tramadol on post subarachnoid block shivering. **Material and Methods:** This prospective study was conducted in the department of Anaesthesiology at Gujarat Adani Institute of Medical Science, Bhuj, Kutch, Gujarat for the period of one year. Seventy ASA Grades-I & II patients of either gender, aged 17 to 61 years scheduled for elective surgeries under subarachnoid block were randomly allocated to two groups (A and B) after obtaining written informed consent. **Results:** Time interval from injecting the study drug to cessation of shivering was quite less with Dexmedetomidine (39.90±5.98) than with Tramadol (209.14±25.78 seconds). Insignificant difference was noted between the two groups in relation to dizziness. Incidence of sedation (Grade 2) was 17/30 in group D while it was 4/30 in group T and the difference was highly significant (p= 0.0012). **Conclusion:** Both Dexmedetomidine and Tramadol can be used for treating post sub-arachnoid block shivering but Dexmedetomidine is more efficacious than Tramadol but should be used cautiously in hemodynamically unstable patient.

Keywords: Dexmedetomidine; Kutch; Shivering; Tramadol.

How to cite this article:

Vishal Joshi, Vijay Goswami. Therapeutic Efficacies of Dexmedetomidine and Tramadol on Post Subarachnoid Block Shivering-A Prospective Study. Indian J Anesth Analg. 2019;6(1):151-54.

Introduction

Shivering, a common post-anaesthesia occurrence is defined as an involuntary, repetitive activity of skeletal muscles. The incidence of shivering has been found to be quite high, approximately 40-50% in different studies [1]. Shivering is a potentially serious complication, resulting in increased metabolic rate; oxygen consumption (up to 100-600%); carbon

dioxide (CO₂) production; ventilation and cardiac output; adverse postoperative outcomes, such as increased surgical bleeding; and morbid cardiac events. Therefore, shivering may cause problems in patients with low cardiopulmonary reserves [2].

Shivering is a physiological response to core hypothermia in an attempt to raise the metabolic heat production. Main causes of intra/postoperative shivering are temperature loss,

Corresponding Author: Vijay Goswami, Assistant Professor, Department of Anaesthesia, Gujarat Adani Institute of Medical Science, Bhuj, Gujarat 370001, India.

E-mail: researchguide86@gmail.com

Received on 11.10.2018, **Accepted on** 14.11.2018

increased sympathetic tone, pain, and systemic release of pyrogens [3,4,5]. Spinal anaesthesia significantly impairs thermoregulation by inhibiting tonic vasoconstriction. It also causes redistribution of core heat from trunk (below the block level) to peripheral tissues. Postoperative shivering can be controlled both pharmacologically and non-pharmacologically.

Non-pharmacological methods using equipment to maintain normal temperature of the body are effective but expensive and lack practicality, while the pharmacological methods using drugs like pethedine, tramadol, clonidine, doxapram, katenserin, nefopam, etc., are simple, cost-effective and easy to implement. During the last decade, Tramadol has become favored and commonly used drug for post-spinal anaesthesia shivering. However, it has many adverse effects like nausea, vomiting, dizziness etc. which cause further discomfort to the patient [6,7].

Dexmedetomidine is a highly selective α_2 -adrenoceptor agonist with potent effects on the central nervous system [8,9]. The stable haemodynamics and the decreased oxygen demand make it a very useful pharmacologic agent [10,11,12]. However, adverse effects such as bradycardia and hypotension limits its use.

Few studies have compared the antishivering effect of above mentioned drugs. Hence, this study aimed to compare therapeutic efficacy of Dexmedetomidine and Tramadol on post subarachnoid block shivering.

Material and Methods

This prospective study was conducted in the department of Anaesthesiology at Gujarat Adani Institute of Medical Science, Bhuj, Kutch, Gujarat for the period of one year. Seventy ASA Grade-I & II patients of either gender, aged 17 to 61 years scheduled for elective surgeries under subarachnoid block were randomly allocated to two groups (A and B) after obtaining written informed consent.

Group A (n=35) received Dexmedetomidine 0.5 μ g/kg IV

Group B (n=35) received Tramadol 0.5 mg/kg IV

Patients with known hypersensitivity to Dexmedetomidine or Tramadol, ASA > 2 class were excluded from the study. The patients were randomly allotted to one of the two groups using a random list. Baseline pulse rate, non-invasive blood pressure (NIBP), oxygen saturation (SpO₂)

and body temperature (axillary) were recorded 2 before the commencement of surgery and thereafter at every 5 minutes from the spinal block (SAB), for 1st hour; and every 15 minutes, for the rest of the observation period.

Subarachnoid block was given with inj. Bupivacaine 0.5% (10-15 mg) at L3-4 or L4-5 interspace using 26 gauge spinal needle and sensory block up to T6-7 dermatome was achieved. All operation theatres in which the operations were performed maintained an ambient temperature of 21-24°C. No means of active re-warming were used. Intravenous fluids and anaesthetic drugs were administered at room temperature. Grade of shivering was determined by Wrench classification [13] as:

Grade 0: No shivering

Grade 1: One or more of the following: Piloerection, Peripheral vasoconstriction, peripheral cyanosis with, but without visible muscle activity

Grade 2: Visible muscle activity confined to one muscle group

Grade 3: Visible muscle activity in more than one muscle group

Grade 4: Gross muscle activity involving the whole body

Patients who developed either grade 3 or grade 4 shivering were included in the study. Either of the two drugs was given as slow IV bolus injection. Time at which shivering started after spinal anaesthesia, severity of the shivering, time to disappearance of shivering and response rate were recorded. If shivering did not subside by 15 minutes, the treatment was considered to be ineffective. Recurrence of shivering was also noticed until the patient left the operation theatre. Patients who did not respond or in whom recurrence of shivering occurred were treated with additional dose of study drug. Side effects like nausea, vomiting, bradycardia (<50/min), hypotension (>20% fall from baseline), dizziness; and sedation score were recorded. Sedation score was assessed with a four-point scale as per Filos [14].

1: Awake and alert

2: Drowsy, responsive to verbal stimuli

3: Drowsy, arousable to physical stimuli

4: Unarousable

Bradycardia, hypotension and vomiting were treated with atropine, mephentermine and ondansetron respectively, in titrated doses when required.

Statistical Analysis

Qualitative data will be expressed as percentages and proportions. Quantitative data will be expressed as mean and standard deviation. The differences between two groups with respect to continuous variables will be analysed using t-test while categorical variables will be analysed using chi-square test. All the statistical tests will be performed in SPSS version 15 software. p value <0.05 will be considered as statistically significant while p value <0.01 will be considered as statistically highly significant.

Results

Dexmedetomidine is as effective as Tramadol in treating post subarachnoid block shivering. Time interval from injecting the study drug to cessation of shivering was quite less with Dexmedetomidine (39.90±5.98) than with Tramadol (209.14±25.78 seconds) [Table 1]. The response rate was 100% in both groups. Incidence of recurrence of shivering post-operatively was 4/30 in group T while 1/30 in group D. Side effects hypotension and bradycardia in group D while nausea and vomiting in group T were significantly high.

Insignificant difference was noted between the two groups in relation to dizziness. Incidence of sedation (Grade 2) was 17/30 in group D while it was 4/30 in group T and the difference was highly significant (p= 0.0012).

Table 1: Time duration for recovery from shivering in A and B Group

Groups	Mean±SD Time Duration (Seconds)	P value
Group A (n=35)	39.90±5.98	0.01*
Group B (n=35)	209.14±25.78	

* indicates statistically significance at p≤0.05, test applied student t test

Discussion

Postoperative shivering is an unpleasant experience for the patient affecting the quality of recovery. Various researchers have compared many drugs with each other to find an ideal agent to prevent intra and postoperative shivering. Tsai YC, Chi KS et al. [15] did a comparison of tramadol, amitriptyline, and meperidine for postepidural anesthetic shivering in parturients. Bilotta F et al. [16] studied Nefopam and Tramadol for the prevention of shivering during neuraxial

anesthesia.

We chose Dexmedetomidine and Tramadol in our study because of their anti-shivering property and compared their efficacy in treating post sub-arachnoid block shivering as well as side effects. The same anti-shivering property was studied in various researches done by Kim YS et al. [17], Bajwa SJ et al. [18], and Blaine Easley R, Brady KM et al. [19] Kim YS et al. [17] searched for an optimal dose of prophylactic dexmedetomidine for preventing postoperative shivering. Bajwa SJ et al. [11,18] found reduction in the incidence of shivering with peri-operative dexmedetomidine. Elvan EG, Oc B, Uzun S et al. [9] used dexmedetomidine in postoperative shivering in patients undergoing elective abdominal hysterectomy. Blaine Easley R, Brady KM et al. [19] studied dexmedetomidine for the treatment of post-anesthesia shivering in children. Tramadol is an opioid analgesic. The anti-shivering action of tramadol is probably mediated via its opioid or serotonergic and noradrenergic activity or both. In the present study, factors that influence the occurrence of shivering, like temperature of IV fluids etc, were not tightly controlled, but this should not affect the validity of our study because the present study is focused on response to treatment used rather than incidence of shivering; and by randomization, both groups were subjected to similar degrees of influence of these factors.

In present study, we found that Dexmedetomidine is as effective as Tramadol in treating post-spinal anaesthesia shivering. In addition, the time interval from the commencement of treatment to cessation of shivering is quite less with dexmedetomidine than with tramadol. Result of our study is coinciding with another similar comparative study of Dexmedetomidine and Tramadol for post-spinal anaesthesia shivering done by Geeta Mittal et al. [10].

Conclusion

Both Dexmedetomidine and Tramadol can be used for treating post sub-arachnoid block shivering but Dexmedetomidine is more efficacious than Tramadol but should be used cautiously in hemodynamically unstable patient.

References

1. De Whitte, Sessler DI. Perioperative shivering: Physiology and Pharmacology. *Anaesthesiology* 2002;96:467-84.

2. Katyal S, Tewari A. Shivering: Anaesthetic considerations. *J Anaesth Clin Pharmacol* 2002; 18:363-76.
3. Bhattacharya P, Bhattacharya L. Post anaesthetic shivering (PAS): A review. *Indian J Anaesth.* 2003; 47:88-93.
4. Kranke P, Eberhart LH, Roewer N, Tramèr MR. Pharmacological treatment of postoperative shivering: A quantitative systematic review of randomized controlled trials. *Anesth Analg.* 2002; 94:453-60.
5. Sessler DI. Temperature regulation and monitoring. In: Millar RD, editor. 7th ed. *Textbook of Anaesthesia*. New York: Churchill Livingstone Inc.; 2010. pp.1533-56
6. Mathews S, Al Mulla A, Varghese PK, Radim K, Mumtaz S. Postanaesthetic shivering – A new look at tramadol. *Anaesthesia.* 2002;57:394-8
7. Shukla U, Malhotra K, Prabhakar T. A comparative study of the effect of clonidine and tramadol on post spinal anaesthesia shivering. *Indian J Anaesth* 2011;55:242-6.
8. Doze VA, Chen BX, Maze M. Dexmedetomidine produces a hypnotic-anesthetic action in rats via activation of central alpha-2 adrenoceptors. *Anesthesiology.* 1989;71:75-9. doi: 10.1097/00000542-198907000-00014
9. Elvan EG, Oc B, Uzun S, Karabulut E, Coskun F, Aypar U. Dexmedetomidine and postoperative shivering in patients undergoing elective abdominal hysterectomy. *Eur J Anaesthesiol.* 2008;255:357-64.
10. Mittal G, Gupta K, Katyal S, Kaushal S. Randomized double-blind comparative study of dexmedetomidine and tramadol for post-spinal anesthesia shivering. *Indian J Anaesth.* 2014;58:257-62.
11. Bajwa SJ, Gupta S, Kaur J, Singh A, Parmar S. Reduction in the incidence of shivering with perioperative dexmedetomidine: A randomized prospective study. *J Anaesthesiol Clin Pharmacol* 2012;28:86-91.
12. Usta B, Gozdemir M, Demircioglu RI, Muslu B, Sert H, Yaldiz A. Dexmedetomidine for the prevention of shivering during spinal anesthesia. *Clinics (Sao Paulo)* 2011;66:1187-91.
13. Wrench IJ, Singh P, Dennis AR, Mahajan RP, Crossley AW. The minimum effective doses of pethidine and doxapram in the treatment of post anaesthetic shivering. *Anaesthesia.* 1977;52:32-6.
14. Filos KS, Goudas LC, Patroni O, Polyzou V. Hemodynamic and analgesic profile after intrathecal clonidine in humans. A dose-response study. *Anesthesiology.* 1994;81:591-601
15. Tsai YC, Chi KS. A comparison of tramadol, amitriptyline and meperidine for postepidural anaesthetic shivering in parturients. *Anaesth Analg* 2002;93:1288-92.
16. Bilotta F, Pietropaoli P, Sanita R, Liberatori G, Rosa G. Nefopam and tramadol for the prevention of shivering during neuraxial anaesthesia. *Reg Anaesth Pain Med.* 2002;27:380-4.
17. Kim YS, Kim YI, Seo KH, Kang HR. Optimal dose of prophylactic dexmedetomidine for preventing postoperative shivering. *Int J Med Sci.* 2013 Aug 13; 10(10):1327-32
18. Bajwa SJ, Bajwa SK, Kaur J, Singh G, Arora V, Gupta S, et al. Dexmedetomidine and clonidine in epidural anaesthesia: A comparative evaluation. *Indian J Anaesth.* 2011;55:116-21
19. Blaine Easley R, Brady KM, Tobias JD. Dexmedetomidine for the treatment of post anesthesia shivering in children. *Paediatr Anaesth* 2007;17:341-6.