

## Characterization of Blood cells and Comparative Analyses of Hematological Profile of Different Breeds of Dairy Cows in Odisha

Ipsita Dash

### How to cite this article:

Ipsita Dash/Characterization of Blood cells and Comparative Analyses of Hematological Profile of Different Breeds of Dairy Cows in Odisha/Indian J Biol 2022; 9(2):41-49.

### Abstract

The pathophysiological conditions of organisms can be revealed by the study of haematological profile. Three breeds of dairy cows of 02-07 years age groups namely, non-descriptive(ND), Red Sindhi (RS) and crossbred Jersey (CBJ) cows, were used in this study. From each breed nine cows were taken. The aim of the present study was to investigate comparative analyses of haematological profile of different breeds of dairy cows of Odisha which is useful to provide a baseline data for comparison with other breeds of cows. For Hb and PCV highly significant difference ( $p<0.01$ ) was there among the three breeds of cows. For TRBC, MCV and MCH no significant difference was there among three breeds. For MCHC highly significant difference ( $p<0.01$ ) is there between ND and CBJ cows and between RS and CBJ cows. For WBC significant difference ( $p<0.05$ ) was there between ND and RS cows. For lymphocyte, neutrophil and basophil no significant difference is there among the different breeds of cows. For monocyte highly significant difference ( $p<0.01$ ) is there among the three breeds of cows. For eosinophil highly significant difference ( $p<0.01$ ) was there between ND and CBJ cows and between RS and CBJ cows.

**Keywords:** ND, RS, CBJ, cows

### INTRODUCTION

Cows belong to the sub-family Bovinae of the family Bovidae. They are raised for milk, cheese, other dairy products, also for meat such as beef and veal and materials such as leather hide. Red Sindhi cattle are the most popular of

all Zebu dairy breeds and originated in the Sindh province of Pakistan, they are widely kept for milk production across Pakistan, India, Bangladesh, Sri Lanka, and other countries. They have been used for crossbreeding with temperate (European) origin dairy breed in many countries to combine their tropical adaptations (heat tolerance, tick resistance, disease resistance, fertility at higher temperature etc.) with the higher milk production found in temperate regions. It has been crossed with Jerseys in many places, including India, the United States, Australia, Sri Lanka, etc. Jersey was bred on the British Channel Island of Jersey and adapts well to various climatic conditions. The cows are calm and docile in nature; bulls may be aggressive. With

**Author's Affiliation:** Scientific Officer, State Forensic Science Laboratory, Rasulgarh 751010, Odisha, India.

**Corresponding Author:** Ipsita Dash, Scientific Officer, State Forensic Science Laboratory, Rasulgarh 751010, Odisha, India.

**E-mail:** [ipsita.dash9@gmail.com](mailto:ipsita.dash9@gmail.com)

**Received date:** 26.07.2022

**Accepted date:** 17.08.2022

the advent of artificial insemination crossbreeding of cattle with European donor breeds are done for improvement of milk production.<sup>1</sup> Physiological equilibrium is maintained mainly by the blood in the body.<sup>2</sup> The haematological values during different physiological situations are essential for the diagnosis of various pathological and metabolic disorders, which can adversely affect the productive and reproductive performance of cows, leading to heavy economic losses.<sup>3</sup>

Three breeds of dairy cows of 02-07 years age groups namely, non-descriptive (ND), Red Sindhi (RS) and crossbred Jersey (CBJ) cows, were used in this study. The aim of the present study was to investigate comparative analyses of haematological profile of different breeds of dairy cows in Odisha which is useful to provide a baseline data for comparison with other breeds of cows.

## MATERIALS AND METHODS

### *Hematological Method*

After disinfecting of the sampling area, blood samples were taken from the jugular vein<sup>4,5</sup> of each individual cow by a skilled veterinary expert for routine haematological examination. Dry and sterilized needles and dry syringes were used for collection of blood.<sup>4</sup> Since EDTA is an excellent anticoagulant<sup>6</sup>, the needle of syringe was inserted through the purple cap of EDTA vial for each sample (K3 EDTA, 2ml \* 13×75mm, Mfg By: HXS Tech Co., Ltd. PRC. For: Peerless Biotech Pvt. Ltd., Chennai, Tamilnadu, India), were kept within an ice box and were taken to the laboratory for haematological examination. After that the sample blood were smeared on grease free microscopic slides.<sup>7</sup> After air drying, the smears were treated with methanol for 1-5 minutes.<sup>5</sup> Estimation of hemoglobin was done by Sahli's acid hematin method<sup>8</sup> with Sahli's haemometer (HiMedia GW 191-1NO, Plane haemometer (Square Type), HiMedia Laboratories Pvt. Ltd., Mumbai, Maharashtra, India). PCV was done by centrifugation of sample blood at 3,500 rpm for 15 minutes.

Total erythrocyte count (TEC) and total leukocyte count (TLC) were done by using the conventional method<sup>5</sup> by using Neubauer's counting chamber. Erythrocyte indices like, MCH, MCV and MCHC were studied according to the methods described by earlier worker.<sup>4</sup> Differential leukocyte counts (DLC) (Sastry, 1983) were assessed by staining the smeared slides with Giemsa stain prepared from Giemsa powder [Qualigens CAS No. 51811-

82-6 Product No. 39382, scientific India Pvt. Ltd., Mumbai, Maharashtra, India] following the standard hematological procedure<sup>9</sup> under 40X objective of light microscope and photomicrography of blood cells were done by CC130-1.3 mega pixel microscopic camera (Mfg. by Catalyst Biotech, Maharashtra, India) connected to microscope under 40X objective.

## STATISTICAL ANALYSES


Mean  $\pm$  SE were calculated for each parameter by using Microsoft Office Excel 2007. For comparison of means, statistical analyses were done by Paleontological statistics (PAST) version 2.17 [Natural History Museum, University of Oslo] for One-Way Analysis of variance (ANOVA) followed by Turkey's pair wise comparison tests. Differences were classified as significant at  $p < 0.05$  and highly significant at  $p < 0.01$ .

### *Photomicrography*

Photomicrography of blood cells were done by CC130-1.3 mega pixel microscopic camera (Mfg. by Catalyst Biotech, Maharashtra, India) connected to microscope under 40X objective.


## RESULTS

Small erythrocytes (microcytes) and large erythrocytes (macrocytes) were observed (Fig. 1).



**Fig. 1:** RBCs (Red blood cells) Microcyte (small arrow), macrocytes (large arrows)

Small lymphocyte (Fig. 2), large lymphocyte (Fig. 3), monocyte (Fig. 4), neutrophil (Fig. 5), eosinophil (Fig. 6) and basophil (Fig. 7) were observed in the blood smears of this study.



Erythrocyte parameters in different breeds of dairy cows of 02-07 years age are recorded (Table 1). Leukocyte parameters in different breeds of dairy cows of 02-07 years age are observed (Table 2).

**Table 1:** Erythrocyte parameters in different breeds of dairy cows of 02-07 years age.

Parameters	Breed		Breed		Breed		F-value
	Non descriptive (ND)		Red Sindhi (RS)		Crossbreed Jersey (CBJ)		
	Mean±S.E.	Range	Mean±S.E.	Range	Mean±S.E.	Range	
Hb(%)	10.47±0.50 <sup>a</sup>	8.45-11.85	12.51±0.54 <sup>a</sup>	10.32-13.95	14.03±0.16 <sup>a</sup>	13.32-14.35	23.42**
TRBC (millions/mm <sup>3</sup> )	7.23±0.95	4.66-11.13	7.01±0.24	6.29-8.06	7.42±0.12	7.05-8.32	1.04NS
PCV(%)	33.07±1.67 <sup>a</sup>	26.3-37.4	40.31±2.59 <sup>a</sup>	31.00-47.00	46.67±1.16 <sup>a</sup>	43.5-52.00	21.47**
MCV(fl)	50.62±5.62	34-76.18	59.81±5.09	39.39-71.86	62.93±1.11	59.18-68.60	2.29NS
MCH(pg)	16.02±1.75	10.69-23.81	18.17±1.29	12.9-21.15	18.44±0.24	17.22-19.57	0.89NS
MCHC(%)	31.69±0.12 <sup>a</sup>	31.17-32.15	30.76±0.54 <sup>b</sup>	29.20-33.29	29.32±0.40 <sup>ab</sup>	27.62-30.64	15.32**

Same lowercase superscript in the same row differs significantly at p<0.05 and p<0.01

\*\* Means highly significant at p<0.01 and NS means not significant

**Table 2:** Leukocyte parameters in different breeds of dairy cows of 02-07 years age.

Parameters	Breed		Breed		Breed		F-value
	Non descriptive (ND)		Red Sindhi (RS)		Crossbreed Jersey (CBJ)		
	Mean±S.E.	Range	Mean±S.E.	Range	Mean±S.E.	Range	
WBC(thousands/mm <sup>3</sup> )	17.92±2.17 <sup>a</sup>	10.95-26.5	12.06±0.95 <sup>a</sup>	8.75-15.95	13.61±0.73	10.60-15.80	3.03*
Lymphocyte(%)	64.87±5.51	38-89	64.55±6.37	36-86	64.57±4.11	54-89	0.0010NS
Monocyte(%)	11.62±3.80 <sup>a</sup>	2-30	3.44±0.47 <sup>a</sup>	2-6	2.42±0.64 <sup>a</sup>	0-5	3.10**
Neutrophil(%)	15.87±3.27	5-31	23.66±5.97	2-49	17.00±2.10	11-24	0.64NS
Eosinophil(%)	3.62±0.88 <sup>a</sup>	0-7	1.55±0.37 <sup>b</sup>	0-4	9.42±1.95 <sup>ab</sup>	7-17	8.996**
Basophil(%)	1.62±0.37	1-4	2.22±0.46	0-5	0.85±0.45	0-3	2.09 NS

Same lowercase superscript in the same row differs significantly at p<0.05 and p<0.01

\*Means significant at p<0.05, \*\* means highly significant at p<0.01 and NS means not significant

Among the three breeds the highest ( $14.03 \pm 0.16$ g/dl) and the lowest ( $10.47 \pm 0.50$ g/dl) mean haemoglobin concentrations were observed in CBJ and ND cows respectively (Fig.8). The highest mean of total erythrocyte count was observed in CBJ cows ( $7.42 \pm 0.12 \times 10^6$ ) and lowest ( $7.01 \pm 0.24 \times 10^6$ ) was observed in RS cows (Fig. 9). The highest average PCV value was observed in CBJ cows ( $46.67 \pm 1.16$ %) and lowest was observed in ND cows ( $33.07 \pm 1.67$ %) (Fig. 10). The highest average MCV ( $62.93 \pm 1.11$ fl) and MCH ( $18.44 \pm 0.24$ pg) were recorded in CBJ cows whereas lowest average MCV ( $50.62 \pm 5.62$ fl) and MCH ( $16.02 \pm 1.75$ pg) were noted in ND cows (Fig. 11 & 12). The highest mean MCHC was there in ND cows ( $31.69 \pm 0.12$ %) and lowest was in CBJ cows ( $29.32 \pm 0.40$ %) (Fig. 13). The highest average

leukocyte count ( $17.92 \pm 2.17 \times 10^3$ ) and lymphocytes ( $64.87 \pm 5.51$ %) were observed in ND cows and lowest average leukocyte count ( $12.06 \pm 0.95 \times 10^3$ ) and lymphocytes ( $64.55 \pm 6.37$ %) were observed in RS cows (Fig. 14 & 15). The highest average monocytes ( $11.62 \pm 3.80$ %) were found in ND cows and lowest ( $2.42 \pm 0.64$ %) were observed in CBJ cows (Fig. 16). The highest average neutrophils ( $23.66 \pm 5.97$ %) were noticed in RS cows and lowest ( $15.87 \pm 3.27$ %) were observed in ND cows (Fig. 17). The highest average eosinophils ( $9.42 \pm 1.95$ %) were found in CBJ cows and lowest average eosinophils ( $1.55 \pm 0.37$ %) were observed in RS cows (Fig. 18). The highest average basophils ( $2.22 \pm 0.46$ %) were noticed in RS cows and lowest ( $0.85 \pm 0.45$ %) were observed in CBJ cows (Fig. 19).




Fig. 8: Average haemoglobin in different breeds of cows.




Fig. 9: Average red blood cell counts in different breeds of cows.




Fig. 10: Average packed cell volume (PCV) in different breeds of cows.




Fig. 11: Average mean corpuscular volume (MCV) in different breeds of cows.




Fig. 12: Average mean corpuscular haemoglobin (MCH) in different breeds of cows.




Fig. 13: Average mean corpuscular haemoglobin concentration (MCHC) in different breeds of cows.




Fig. 14: Average total leucocyte count (TLC) in different breeds of cows.




Fig. 15: Average mean lymphocytes in different breeds of cows.




Fig. 16: Average mean monocytes in different breeds of cows.




Fig. 17: Average mean neutrophils in different breeds of cows.




Fig. 18: Average mean eosinophils in different breeds of cows.




Fig. 19: Average mean basophils in different breeds of cows.

For Hb and PCV highly significant difference ( $p < 0.01$ ) was there among the three breeds of cows. For RBC, MCV and MCH no significant difference was there among three breeds. For MCHC highly significant difference ( $p < 0.01$ ) is there between ND and CBJ cows and between RS and CBJ cows. For WBC significant difference ( $p < 0.05$ ) was there between ND and RS cows. For lymphocytes, neutrophils and basophils no significant difference is there among the different breeds of cows. For monocytes highly significant difference ( $p < 0.01$ ) is there among the three breeds of cows. For eosinophils highly significant difference ( $p < 0.01$ ) was there between ND and CBJ cows and between RS and CBJ cows.

## DISCUSSION

In veterinary medicine, examinations of hematological parameters present an effective tool in monitoring the health and nutritional status of animal.<sup>10</sup> The haematological values provide valuable baseline information and help in realistic evaluation of managerial practices, nutritional and physiological status of animals and diagnosis of health condition.<sup>11</sup> The erythrocyte parameters and total and differential leucocyte counts are affected by various physiological determinants.<sup>12</sup> The typical RBCs shape for multiple veterinary species is the disc or biconcave disc (discoid) resulting in a high surface area to volume ratio making the red blood cells deformable. Central pallor can be observed to various degrees in these species on examination of a peripheral blood smear.<sup>13-16,7</sup>

Long et al. (1952)<sup>17</sup> were working with Shorthorn, Angus and Herefords and Rusoff et al. (1954)<sup>18</sup> working with Jerseys, Guerneys and Holsteins found no breed differences in haemoglobin content of blood, the present study disagrees with these; whereas Mac Donald et al. (1956)<sup>19</sup> found that Angus males were higher in haemoglobin than Hereford males. Earlier worker<sup>20</sup> reported a decrease in haemoglobin content when calves grew older but no marked difference was observed in total number of red blood cells. Cornelius (1956)<sup>21</sup> determined blood haemoglobin values for dwarf beef cattle and concluded that a mean value of  $11.7 \pm 1.4$  g of haemoglobin per 100ml of blood was within the normal range for the bovine species. According to Mac Donald et al. (1956)<sup>19</sup> an average value of 12.0g of haemoglobin per 100ml of blood for Hereford and Angus beef calves on performance test at 500- and 800-pounds body weight. Late pregnancy and onset of lactation is a period when slight anaemia exists.<sup>22</sup> The Hb value of  $117.4 \pm 2.42$  and  $106.3 \pm 2.21$ , the RBC value of  $8.49 \pm 0.21$  and  $7.63 \pm 0.16$ , the haematocrit value of  $36.15 \pm 0.75$  and  $32.10 \pm 0.64$ , the MCV value of  $43.55 \pm 0.69$ ,  $42.79 \pm 0.61$ , the MCH value of  $14.42 \pm 0.43$ ,  $14.03 \pm 0.17$ , MCHC value of  $324.7 \pm 1.38$  and  $330.1 \pm 1.48$  and WBC value of  $11.88 \pm 0.36$  and  $10.57 \pm 0.29$  were studied respectively by previous researcher<sup>23</sup> for Muturu and Bunaji breeds of cattle in Nigeria.

A range of 4.9 to 9.98 million red blood cells per cu. mm of blood has been reported for dairy cattle<sup>24,25, 26,20,18,27,28</sup> and 4.5 to 9.8 million per cu.mm of blood for beef cattle.<sup>17,29</sup> Rusoff (1946)<sup>31</sup> explained the variable means for red blood counts on dairy cows

(Jerseys 6.55, Guernseys 7.49 and Holsteins 7.84 million per cu.mm). Long et al. (1952)<sup>17</sup> reported the composition of blood taken from Hereford, Angus and Shorthorn beef cattle. All the breeds had a normal variation from 4.5 to 6.0 million red blood cells per cu.mm. This seems low when compared with data on dairy cattle which is congruent with Dash et al., 2015.<sup>28</sup> The difference may be due to the indirect turbidity technique used in determining the red cell count. From complications of information of several investigators, Coffin (1953)<sup>26</sup> reported a red blood cell count ranging from 5.4 to 9.0 million per cu.mm for cattle. In my study the range of RBCs count ranging from 8.45-11.85 million in ND cows, 10.32-13.95 million in RS cows and 13.32-14.35 million in CBJ cows. Ferguson (1954)<sup>24</sup> observed a mean of 6.33 million per cu. mm with a range of 5.64 to 7.44 million per cu.mm for dairy cattle. Dukes (1955)<sup>28</sup> noted an overall mean of 6.3 million per cu.mm and a range of 4.90 to 9.79 per cu.mm calculated from 144 samples on six dairy cows. Cornelius (1956)<sup>21</sup> found, from data on 38 Hereford and Angus dwarf cattle ranging in age from 6 days to 14 months, a mean red cell count of  $9.8 \pm 1.6$  million per cu.mm. Arthaud et al. (1959)<sup>29</sup> reported both an increase in haemoglobin and in total number of red blood cells as Hereford, Shorthorn and Angus calves grew from 7 months old to 9 months old. Age<sup>30-32</sup>, sex<sup>33</sup>, breed, exercise<sup>34,35</sup>, pregnancy and lactation<sup>36-38</sup>, emotional state are variables to be considered when establishing references values in domestic animals, altitude polyglobulia (increased RBCs count at high altitude) is well documented and extensively studied.<sup>39,40</sup>

The total leukocyte count of 8,580 per cu.mm of blood is reported by.<sup>18</sup> Greatorrex (1954)<sup>20</sup> found the number of leukocytes per cu.mm of blood, in dairy calves from birth to a year of age, ranged between 4,500 and 15,000 with the majority ranging from 6,500 to 11,500 per cu.mm. Dash et al. (2015)<sup>28</sup> reported a mean value of  $26.00 \pm 0.26$  for 02 years ND cows, a mean value of  $16.68 \pm 0.23$  for 06 years ND cows and a mean value of  $11.08 \pm 0.07$  for 10 years ND cows, a mean value of  $11.73 \pm 0.27$  for 02 years RS cows, a mean value of  $8.93 \pm 0.10$  for 06 years RS cows and a mean value of  $15.51 \pm 0.21$  for 10 years RS cows, a mean value of  $14.33 \pm 0.14$  for 02 years CBJ cows, a mean value of  $10.78 \pm 0.11$  for 06 years CBJ cows and mean value of  $15.71 \pm 0.04$  for 10 years CBJ cows. Other studies in fatty liver cows, without being downer reported decreased<sup>41, 42</sup> or increased WBC.<sup>43</sup> Previous researcher<sup>44</sup> also reported that there was an increase in the number of leukocytes during gestation.

Coffin (1953)<sup>26</sup> opined normal ranges and means for

differential white cell counts of cattle. Neutrophils range from 1 to 15, with a mean of 30%; eosinophils range from 1 to 15, with a mean of 8%, basophils range from 0 to 1, with a mean of 0.5%; lymphocytes range from 40 to 70; with a mean of 52%; and monocytes range from 3 to 15, with a mean of 9%. Dash et al. (2015)<sup>28</sup> reported a mean value of  $72.33 \pm 3.17\%$ ,  $74.50 \pm 14.50\%$ ,  $51.00 \pm 7.00\%$  lymphocytes for 02 years, 06 years and 10 years ND cows respectively, a mean value of  $11.73 \pm 0.27\%$ ,  $8.93 \pm 0.10\%$  and  $15.51 \pm 0.21\%$  lymphocytes for 02 years, 06 years and 10 years Red Sindhi cows respectively, a mean value of  $14.33 \pm 0.14\%$ ,  $10.78 \pm 0.11\%$ ,  $15.71 \pm 0.04\%$  lymphocytes for 02 years, 06 years and 10 years Crossbred Jersey cows respectively.

The haematological parameters of cattle are influenced by many factors like breed, age, sex, seasonal variation, lactation, pregnancy, health and nutritional status of the animal.<sup>45</sup> Dash et al. (2015)<sup>28</sup> reported a mean value of  $5.00 \pm 1.52\%$ ,  $3.00 \pm 1.00\%$ ,  $24.00 \pm 3.05\%$  monocytes for 02 years, 06 years and 10 years non-descriptive cows respectively, a mean value of  $3.66 \pm 0.33\%$ ,  $2.33 \pm 0.33\%$  and  $4.33 \pm 1.20\%$  monocytes for 02 years, 06 years and 10 years Red Sindhi cows respectively, and a mean value of  $3.50 \pm 0.50\%$ ,  $2.66 \pm 1.20\%$  and  $1.00 \pm 1.00\%$  monocytes for 02 years, 06 years and 10 years Crossbred Jersey cows respectively.

Dash et al. (2015)<sup>28</sup> reported a mean value of  $14.00 \pm 2.501\%$ ,  $16.00 \pm 11.00\%$ ,  $17.66 \pm 6.88\%$  neutrophils for 02 years, 06 years and 10 years non-descriptive cows respectively, a mean value of  $8.00 \pm 4.16\%$ ,  $46.33 \pm 1.45\%$  and  $16.66 \pm 2.33\%$  of neutrophils for 02 years, 06 years and 10 years Red Sindhi cows respectively, and a mean value of  $22.00 \pm 1.00\%$ ,  $17.66 \pm 3.17\%$  and  $11.00 \pm 0.00\%$  of neutrophils for 02 years, 06 years and 10 years Crossbred Jersey cows respectively.

Dash et al. (2015)<sup>28</sup> reported a mean value of  $4.66 \pm 0.33\%$ ,  $4.00 \pm 1.00\%$  and  $2.33 \pm 2.33\%$  of eosinophils for 02 years, 06 years and 10 years non-descriptive cows respectively, a mean value of  $1.66 \pm 1.20\%$ ,  $2.00 \pm 0.00\%$  and  $1.00 \pm 0.00\%$  of eosinophils for 02 years, 06 years and 10 years Red Sindhi cows respectively, and a mean value of  $7.00 \pm 0.00\%$ ,  $10.33 \pm 4.80\%$  and  $10.50 \pm 1.50\%$  for 02 years, 06 years and 10 years Crossbred Jersey cows respectively.

Dash et al. (2015)<sup>28</sup> reported a mean value of  $1.33 \pm 0.33\%$ ,  $2.50 \pm 1.50\%$  and  $1.33 \pm 0.33\%$  of basophils for 02 years, 06 years and 10 years non-descriptive cows respectively, a mean value of  $3.00 \pm 1.15\%$ ,  $2.00 \pm 1.00\%$  and  $1.66 \pm 0.88\%$  of basophils for 02 years, 06 years and 10 years Red Sindhi cows respectively,

and a mean value of  $1.00 \pm 1.00\%$ ,  $1.00 \pm 1.00\%$  and  $0.50 \pm 0.50\%$  for 02 years, 06 years and 10 years Crossbred Jersey cows respectively.

Greatorex (1954)<sup>20</sup> has reported a decrease in the number of lymphocytes as calves grew from birth to one year of age and observed the number of monocytes, neutrophils and eosinophils to be variable from age to age while basophils were absent. Cornelius (1956)<sup>21</sup> found that all haematological values appeared normal with the exception of the differential leukocyte count. Seasonal of some diseases may influence the levels of certain blood indices. In spring more inflammatory processes occurs which may stimulate the neutrophils.<sup>46</sup> Changes in white blood cell counts are observed around parturition, for example, an increase in the numbers of circulating neutrophils.<sup>47,48</sup> It is accepted that fatty liver usually induces low leukocyte counts in the peripheral blood<sup>49,50</sup>; one author<sup>51</sup> stated that in calves the most disadvantageous influences are high temperature and high humidity and, in these conditions, the greatest changes occur in the white blood system in bone marrow and in peripheral blood. Pereira et al. (1987)<sup>44</sup> reported that there was an increase in the number of leukocytes.

A range in average haematocrit values in dairy cattle from 28 to 50% has been reported<sup>52, 53, 20</sup> and in beef cattle from 31 to 48%.<sup>29</sup> The packed cell volume (PCV) is one of the most valuable techniques for determination of the percentage of cellular components of blood in the clinical laboratory.<sup>54</sup> As per previous study haematocrit value of Muturu cattle was  $36.15 \pm 0.75$ <sup>23</sup> and 35.86 in West African cattle breeds as stated by Berthier et al. (2015)<sup>55</sup> and  $29.23 \pm 0.42$  reported by Mbanasor et al. (2003)<sup>56</sup> which is not in accordance of our study. Albritton (1952)<sup>52</sup> puts forth a range of 14.2 to 18.5 micro micrograms with a mean of 15.7 micro micrograms for corpuscular haemoglobin content of the blood of the adult cow. Normal ranges reported on cattle by<sup>26</sup> are 14.4 to 18.6 micro micrograms for mean corpuscular volume and 32% to 34% for mean corpuscular haemoglobin concentration. Mean corpuscular volume in dairy and beef cattle varied from 40.0 to 60.7 cubic microns [52, 26, 20]. Average of 32.0 and 37.2% for mean corpuscular haemoglobin were observed.<sup>27</sup> Greatorex (1954)<sup>20</sup> calculated the mean corpuscular volume on 233 dairy calves from birth to one year of age. The extreme values were 28 to 112 cubic microns whereas the mode was from 40 to 60 microns and individual values were evenly distributed throughout the first 12 months of life. Despite the fact wide variation of normal ranges and the anisocytosis that usually appear<sup>57</sup> make the

diagnostic value of MCHC for cows with fatty liver poor.

## CONCLUSION

The haematological results obtained in this study vary according to different breeds. Differences in results are may be due to environmental, genetic, non-genetic factors, managerial practices, body weight, age groups and due to breed difference. These factors may be considered when establishing reference values and this study is helpful for interpretation of haematological values with other breeds.

## REFERENCES

1. Singh, C.V., 2016. Cross-breeding in Cattle for milk production: Achievements, Challenges and Opportunities in India-A Review. *Adv. Dairy. Res.* 4,158.
2. Geneser, F., 1986. *Textbook of histology.* Denmark: Munksgaard, Copenhagen pp. 335-352.
3. Dutta, J.C., et al., 1988. Blood biochemical studies in anoestrus and normal cyclic cattle. *Indian. Vet. J.* 65, 239-241.
4. Sastry, G.A., 1983. *Vet. Clin. Pathol.* Delhi: CBS Publishers and Distributors, pp.1-18.
5. Brar, R.S., et al., 2002. *Veterinary Clinical Diagnosis by Laboratory Methods.* Ludhiana: Kalyani Publisher, pp. 10.
6. Harvey, J.W., 2001. *Atlas of Veterinary Hematology: Blood and Bone marrow of Domestic Animals.* USA, Philadelphia: WB Saunders Company, pp. 1-40.
7. Dash, I., 2019. *Comp. Clin. Pathol.* <http://doi.org/10.1007/s00580-019-03037-3>
8. Sahli, H., 1953. *Lehrbuch. D.Klin. Untersuchungen Method,* 5th ed. Leipsic. pp. 846.
9. Lillie, R.D. (Ed.), 1977. *H.J. Conn's Biological Stains.* Baltimore, USA: The Williams and Wilkins Company, pp. 606-607.
10. Klinkon, M., Zadnik, T., 1999. Dynamics of red and white blood picture in dairy cows during periparturient period. *Comp. Haematol. Intern.* 9, 156-161.
11. Osman, T.E.A., Busadah, K.A., 2003. Normal concentrations of twenty serum biochemical parameters of she-camels, cows and ewes in Saudi Arabia. *Pakistan J. Biol. Sci.* 6(14),1253-1256.
12. Klinkon, M., 1992. Red blood picture of cattle in Slovenia with regard to breed, sex, age,

- physiological state and rearing conditions. Dissertation, Veterinary Faculty, Ljubljana, Slovenia.
13. Lording, P.M., 2008. Erythrocytes. *Vet. Clin. N. Am. Equine Pract.*, 24, 225-237.
  14. Barger, A.M., 2010. Erythrocyte morphology, in: Weiss, D.J., Wardrop, K.J., (Eds.) *Schalm's Veterinary Hematology*. Wiley-Blackwell Publishing Ltd., Ames, USA, Iowa, pp. 144-151.
  15. Wood, D., Quiroz-Rocha, G.F., 2010. Normal hematology of cattle, in: Weiss, D.J., Wordrop, K.J. (Eds.) Iowa, USA: Wiley-Blackwell Publishing Ltd., Ames, pp. 829-835.
  16. Dash, I., Mohanty, P.K., 2015. Morphological features and influence of age and breed on the morphometry of red blood cells of female cattle. *I.J.S.R.* 4(2), 259-264.
  17. Long, R.A., et al., 1952. Blood composition of normal beef cattle. Stillwater, Oklahoma Agricultural and Mechanical College, Okla. *Agr. Exp.Stat. Tech. Bul. No 43*, pp. 16.
  18. Rusoff, L.L., et al., 1954. Blood studies of breeding dairy bulls. I. Hematocrit, hemoglobin, plasma calcium, plasma inorganic phosphorous, alkaline phosphatase values, erythrocyte count and leukocyte count. *J. Dairy. Sci.* 37, 30.
  19. Mac Donald, M.A., et al., 1956. Rate and efficiency of gains in beef Cattle. Corvallis, Oregon State College. *Ore. Agr. Exp. Stat.Tech. Bul. No. 36*, pp. 34.
  20. Greatorex, J.C., 1954. Studies on the hematology of calves from birth to one year of age. *Br. Vet. J.* 110, 120-138.
  21. Cornelius, C.E., et al., 1956. Chemical and haematological studies on blood of bovine dwarfs. *P.S.E.B.M.*, 92(3), 522-524.
  22. Stöber, M., Grunder, H.D., 1979. Circulation. in: Rosenberger, G. (Ed.). *Clinical examination of cattle (Translation of the second edition 1977)*. Verlag Paul Parey, Berlin and Hambur, pp. 128.
  23. Ode, S.A., et al., 2016. Haematology and some serum biochemistry of apparently healthy Muturu and Bunaji breeds of cattle in Benue State, Nigeria. *Comp. Clin. Pathol.* DOI 10.1007/s00580-016-2375-1
  24. Ferguson, L.C., 1945. On variation in the blood cells of healthy cattle. *J. Infect. Dis.* 76, 24-30.
  25. Reid, I.M., Roberts, C.J., 1983. Subclinical fatty liver in dairy cows, current research and future prospects. *Ir. Vet. J.* 37, 104-110.
  26. Coffin, D.L., 1953. *Manual of Veterinary clinical pathology*. Ithaca: Comstock, pp. 322.
  27. Dukes, H.H. 1955. *The physiology of domestic animals*. Comstock Publishing Association, Ithaca, New York, pp. 1-1020.
  28. Dash, I., et al., 2015. Haematological analysis of three breeds of cows. *I.J.B.* 2(1),31-39.
  29. Arthaud, R.L., et al., 1959. The relationship of certain blood constituents to rate and economy of gain in beef cattle. *J. Anim. Sci.* 18, 314.
  30. Harvey, J.W., et al., 1984. Hematology of the foals up to one year old. *Equine. Vet. J.* 16(4), 347-353.
  31. Brun-Hansen HE, Kampen AH, Lund A. Hematologic values in calves during the first six months of life, *Vet ClinPathol* 2006; 35(2): 182-187.
  32. Aoki, T., Ishii, H., 2012. Hematological and biochemical profiles in peripartum mares and neonatal foals (heavy Draft horse). *J. Equine. Vet. Sci.* 32, 170-176.
  33. Shaikat, A.H., et al., 2013. Hemato-biochemical profiles of indigenous goats (*Capra hircus*) at Chittagong, Bangladesh. *Vet. World.* 6(10), 789-793.
  34. Acena, M.C., et al., 1995. Modifications hematologiques at musculaires pendant la corrida chez le taureau de combat. *Rev. Méd. Vét.* 146 (4), 277-282.
  35. Zobra, R., et al., 2011. Physical, hematological and biochemical responses to acute intense exercise in polo horses. *J. Equine. Vet. Sci.* 31, 542-548.
  36. Masoni, F., et al., 1985. Paramètres Hématologiques de la chèvre laitière Variations physiologiques chez l' animal Sain autour de la mise-bas. *Rec. Méd. Vét.* 161(1), 41-49.
  37. Roy, S., et al., 2010. Hematological and biochemical profile during gestation period in Sahiwal cows. *Vet. World.* 3(1), 26-28.
  38. Mariella, J., et al., 2014. Hematologic and biochemical profiles in standardbred mares during peripartum. *Theriogenology.* 81(4), 526-534.
  39. Kolb, E., 1975. *Physiologie des animaux domestiques*. Edition Vigot Freres. France, Paris, pp. 1-918.
  40. George, J.W., et al., 2008. Changes in bovine hematology references intervals from 1965 to 2001. *Vet. Clin. Pathol.* 36, 312-313.
  41. Morrow, D.A., et al., 1979. Clinical investigation of a dairy herd with the fat cow syndrome. *J. Am. Vet. Med. Assoc.* 15, 161-167.
  42. Reid, J.T., et al., 1948. Simple versus complex concentrate mixtures for young breeding bulls. I. Growth, blood composition and cost. *J. Dairy Sci.* 31, 429.
  43. Padilla-Arenalles, S., et al., 2007. Coagulation tests and selected biochemical analytes in dairy cows with hepatic lipidosis. *Acta. Vet. Brno.* 76, 541-546.



44. Pereira, J.L., et al., 1987. Hematological variations related to gestation and age in the autochthonous bovine breed. *Blanca Cacerena. An. de Vet. de Mur.* 3, 93-97.
45. Sattar, A., Mirza, R.H., 2009. Hematological parameters in exotic cows during gestation and lactation under subtropical conditions. *Pakistan. Vet. J.* 29(3),129-132.
46. Schalm, O.O., 1962. Leukocyte responses to disease in various domestic animals. *J. Am.Vet. Med. Assoc.* 140, 557.
47. Guidry, A.J., et al., 1976. Effects of parturition and lactation on blood and milk cell concentrations, corticosteroids and neutrophil phagocytosis in the cow. *Am. J. Vet. Res.* 37, 1195-1200.
48. Kehrli, Jr M.E., et al., 1989. Alterations in bovine neutrophil function during the peripartum period. *Am. J. Vet. Res.* 50, 207-214.
49. Reid JT, Ward GM, Salsbury. Simple versus complex concentrates mixture for young breeding bulls. I. Growth, blood composition and cost. *J Dairy Sci* 1948; 31:429.
50. Franklin, S.T., et al., 1991. Effects of ketones, acetate, butyrate and glucose on bovine lymphocyte proliferation. *J. Dairy Sci.* 74, 2507-2514.
51. Deptula, W., Buczek, J., 1982. Wplyw warunków środowiska na niektóre mechanizmy odporności bydła. *Med. Wet.* 38, 51-54 [In Polish with English summary]
52. Albritton, E.C., (Ed.), 1952. Standard values in blood. Philadelphia: Saunders. pp.199.
53. Holmen, H.H., 1952. A negative correlation between size and number of the erythrocytes of cows, sheep, goats and horses. *J. Path. And. Bact.* 64, 3.
54. Coles, E.H., 1986. *Veterinary Clinical Pathology.* Philadelphia: WB Saunder, pp. 457.
55. Berthier, D., et al., 2015. A comparison of phenotypic traits related to trypanotolerance in five West African cattle breeds. *PLoS One* 10(5), 1-21. doi:10.1371/journal.pone.0126498
56. Mbanasor, U.U., et al., 2003. Haematology of normal and trypanosome infected Muturu cattle in Southeastern Nigeria. *Nig. J. Anim. Prod.* 30(2),236-241.
57. Weiss, D.J., Perman, V., 1992. Assessment of the hematopoietic system in ruminants. *Vet. Clin. N. Am.-Food AnimPract.*, 8, 411-428.

