

***In-Vitro* Callus Induction and Shoot Regeneration in *Justica adathoda* (L.): Potent Indian Medicinal Plant**

P. Muthuramalingam *, J. Vijayakumar**

Author's Affiliation: * Department of Biotechnology, Alagappa University, Karaikudi - 630004, Tamil Nadu, India, **Department of Biotechnology, Ayya Nadar Janakiammal College (Autonomous), Sivakasi - 626124, Tamil Nadu, India.

Abstract

Not provided.

Keywords

Justica adathoda; Tissue culture; Medicinal plant

Introduction

In vitro propagation is an important tool for rapid multiplication of medicinal plants [1] as well as for the extraction of active ingredients. *Justica adathoda* (L) commonly known as Adhathoda in ayurveda belongs to the family Acanthaceae. It is small clambering vine, leaves small, cordate, apex acuminate, corolla composed of fully fused petals, white colored, annual plant. Prefers western gatz, fences or low ground cover as a substrate. The leaves collected from wild are eaten as a cooked vegetable in Kenya and added to soup in Nigeria. Ayurveda has identified many medicinal properties of this plant and it is effectively used against dysentery [2,3] choosy leaves with alcohol are applied to open sores and pustules. A paste of leaves is applied on throat infection, respiratory problems and bronchial asthma. Flower are used as cleaning agents to improve difficult breathing, relive pain and to improve vision. It is being used as an antioxidant. The plants of this family are extensively investigated as a newer source of natural antioxidants and for other bioactive compounds of human benefits.[4] Due to its seasonal availability and endemic distribution the

present investigation was focused on to obtain callus from leaves, node and bud. The *in-vitro* multiplication may benefit as the perennial source for the isolation of bioactive compounds.

Materials and Methods

Plant Material

Healthy plant of *Justica adathoda* (L.) was collected from Western gatz, Suruli falls, Theni, Tamil nadu, India. The different parts of plants like leaves, node and bud were washed with tap water and then with detergent (Teepol) for 15 min followed by washing with distilled water.

Preparation of Explants

Leaf as Explants: The leaf explants were cut into small pieces and washed with running water. Then, the explants were surface-sterilized with 0.1% (w/v) mercuric chloride for 2-3 min, followed by 70% ethyl alcohol 2-3 min, washed 3-4 times with sterile double-distilled water and inoculated on agar-solidified MS (Murashige & Skoog) medium supplemented with different concentrations of 2,4-D, kinetin and BAP, either alone or in combination, with 3% (w/v) sucrose. The pH of the medium is adjusted to 5.8 before sterilization. Cultures were maintained at 27±1°C photoperiod.

Corresponding Author: J. Vijayakumar,
E-mail:pandianmuthuramalingam@gmail.com

Node as Explants: Nodal region was surface sterilized (similar to leaves) with 0.1% mercuric chloride and 70% ethyl alcohol and cultured on agar solidified MS medium supplemented with different various combinations and concentrations of Auxins 2,4 D (0.1-0.8 mg/L), NAA (0.2,0.4 and 0.8 mg/L) and cytokines kinetin (0.1-0.8 mg/L), or BAP (0.2 mg/L) with 3% (w/v) sucrose. The pH of the medium was adjusted to 5.8 before sterilization. Cultures were maintained at $27\pm 1^\circ\text{C}$ photoperiod.

Bud Explants: Bud explants were surface-sterilized with 0.1% (w/v) mercuric chloride for 2-3 min, followed by 70% ethyl alcohol 2-3 min, then washed 3-4 times with sterile double-distilled water and inoculated on agar-solidified MS medium supplemented with different combinations and concentrations of auxins 2,4 D (0.1-0.8 mg/L), NAA (0.2,0.4 and 0.8 mg/L) and cytokines kinetin (0.1-0.8 mg/L), or BAP (0.2 mg/L) with 3% (w/v) sucrose. The pH of the medium was adjusted to 5.8 before sterilization. Cultures were maintained at $27\pm 1^\circ\text{C}$ photoperiod. The experiment was terminated after an interval of 30 days. In another set of experiments where the shoot regeneration capacity was determined, for this node explants was inoculated in the tubes containing MS medium supplemented with kinetin and NAA in concentrations. At least 10 tubes were inoculated and incubated under optimal condition as defined above. After 30 days, the experiment was terminated and shoot generation capacity, its length and morphology were recorded.

Shoot Culture

Basal medium used for initial set of experiment for shoot proliferation consisted of MS salt 3% (w/v) sucrose, and 0.9% (w/v) agar. The pH of medium was adjusted to 5.7. The basal medium was supplemented with various combinations of auxins (NAA, 2,4-D) with cytokines (Kinetin, BAP) at different concentration. The cultures were incubated at $27 \pm 1^\circ\text{C}$ under 16 hr photoperiod with cool white fluorescent light.

Results and Discussion

Callus Induction: All the combinations of NAA, 2, 4-D with kinetin and BAP produced callus (Table 1&2). Optimum concentration of auxins and cytokines which initiated callus with high percentage of response was used for further study. Callus induction was different in different explant with different hormone concentrations. The highest response of callus formation in leaves ($94.3\pm 2.54\%$) was observed in MS medium supplemented with 0.8 mg/L 2,4-D plus 0.8mg/L kinetin which resulted in white soft callus. The combination of NAA ($95.6\pm 0.96\%$ 0.2mg/L) with 0.2 mg/L of BAP gave greenish white hard callus. Callus induction was observed after 7 days of culturing of leaf samples (Table 1, 2; Plate 1). The highest response in node explant ($84.3\pm 2.6\%$) was observed on MS medium supplemented with 0.4 mg/L 2,4-D with 0.4mg/L kinetin which resulted in white soft callus; while, $87.6\pm 2.53\%$ 0.2 mg/L of NAA with 0.2mg/L BAP resulted in greenish white hard callus. Callus induction was observed after 6 days in case of node (Table 1, 2; Plate1). These results are in support of earlier investigations carried out for callus induction in *Ipomoea obscura* (L). using node as explant (A. Mungole et al., 2009). The highest response of callusing was observed in bud explant on MS medium supplemented with 0.2 mg/L of 2, 4-D plus 0.2 mg/L of kinetin which resulted in white callus (82.33 ± 0.9). While, 0.8 mg/L of NAA with 0.8mg/L kinetin resulted in green callus ($96.6\pm 0.96\%$). Callus induction was reported in 6 days of culturing in medium with 2, 4-D plus Kinetin and 4 days in NAA plus Kinetin (Table 1, 2; Plate1). Growth of callus increased significantly and this covers the entire surface of the explant. In general, it was observed that NAA was the best source of auxin for callus induction along with kinetin (80– 100%) or with BAP (90-96%); then 2, 4- D with kinetin (40-90%) or with BAP (55-80%) (Table 1, 2). The callus produced by 2,4-D was white in contrast to dark green in NAA. MS

Table 1: Effect of different concentration of 2, 4- D with Kinetin and BAP on Callus induction

Auxins (mg/L)	Cytokines (mg/L)		% of callus response			Nature of Callus
	2,4-D	Kinetin	BAP	Leaf explants	Node explants	
0.1	0.1	--	90±1.66	70±1.6	46.6±2.5	White, Soft
0.2	0.2	--	86.6±2.5	84.3±2.6	82.33±0.9	White, Soft
0.4	0.4	--	85±1.6	80.3±1.5	NR	White, Soft
0.8	0.8	--	94.3±2.54	75±1.05	NR	White, Soft
0.1	--	0.2	56.6±0.96	79.6±1.5	NR	White, Soft

Media – MS +3% sucrose, NR = No Response, Mean value of three readings.

Table 2: Effect of concentration of NAA with Kinetin and BAP on Callus induction

Auxins (mg/L)	Cytokines (mg/L)		% of callus response			Nature of Callus
	2,4-D	Kinetin	BAP	Leaf explants	Node explants	
0.2	--	0.2	95.6±0.96	87.6±2.53	91.6±0.96	Greenish white, Hard
0.4	0.4	--	80±1.66	78.3±0.96	86.67±0.96	Greenish white, Hard
0.8	0.8	--	NR	93.3±0.96	96.6±0.96	Greenish white, Hard

Media: MS + 3% sucrose; NR= No response; Mean value of three readings.

Table 3: Effect of different concentration of Hormone(S) NAA(0.8 Mg/L) and Kinetin (0.8 Mg/L) on shoot regeneration and height attained from Node Explant of *Ipomoea obscura (L.)*

No. of. testtubes Inoculated	No. of. shoot per treatment	Shootlength in cm.	Shoot morphology
1	2	3.6	Thin Short
2	2	2.7	Thin Short
3	3	3.1	Thin Short
4	2	5.4	Green and long
5	4	3.8	Thin Short
6	2	7.5	Green and long
7	1	3.2	Thin Short
8	NR	NR	High callus induction
9	NR	NR	High callus induction
10	NR	NR	High callus induction

NR = No response

medium is frequently used for micropropagation of large number of plants.[5] It was reported that *T. bellerica* cultures grew better on MS medium in comparison to all other medium (Rathore et al.,2008). The medium for *in-vitro* multiplication of *Drosera* plants was MS medium. MS was reported superior medium for micropropagation of *Coptis teeta*. [6] Further studies were carried out for shoot regeneration capacity of the node. Shoot were initiated from the node explants by showing both direct and indirect organogenesis. The best result of shooting (7.5 cm; 4 shoots

per treatment) was observed in MS medium supplemented with the combination of Kinetin (0.8 mg/l) and NAA (0.8 mg/l) after 17-19 days (Table 3; Plate1).

Conclusion

Depletion of wild population can be prevented through such invitro cultivation for further commercial exploitation. Amongst all the

Fig A: White Callus**Fig B: Greenish White Colour Callus****Fig C: Initiation of Shoot****Fig D: Multiple Shoots**

explants tried for callus, the leaf was found to be the most suitable explant for callusing. NAA and BAP were found to be the most appropriate hormone concentration for callusing. Thus, the present investigation provides optimum parameters for callus and shoots induction. The multiple shooting of *Justica adathoda* (L.) was also established for single nodal region on MS medium supplemented with NAA (0.8 mg/L) and Kinetin (0.8 mg/L). Thus this study provides a standard protocol to initiate multiple shoot culture, optimization of media content and hormonal concentration that may provide desired source of pharmacologically active plant constituents through callus culture.

References

1. Atal CK and Kapur BM. Cultivation and Utilization of Medicinal plants. Jammu Tawi: Regional Research Laboratory CSIR; 1982b.
2. Christophe W, Pharm D, Humana press. Ethnopharmacology of medicinal plants. *Asia and the Pacific*. 2002.; 1(69): 91.
3. Eckart E. Solanaceae and Convolvulaceae – secondary metabolite. 2008; 1: 637.
4. Lakshmi B and Vimala V. Nutritive value of dehydrated green leafy vegetables. *J Food Sci Tech*. 2000; 37: 465-474
5. Feyissa T, Welander M and Nagesh L. Micropropagation of *Hagenia abyssinica*: A multipurpose tree. *Plant Cell Tissue Organ Cult*.

2005; 80: 119-127.

medicinal plant of Arunachal Pradesh, India. *Indian J Biotechnol.* 2007; 6: 280-282.

6. Pramod T, Tilok SR and Suman K.
Micropropagation of *Coptis teeta* wall-Threatened

Indian Journal of Trauma and Emergency Pediatrics

Handsome offer for Indian Journal of Emergency Pediatrics subscribers

Subscribe **Indian Journal of Trauma and Emergency Pediatrics** and get any one book or both books absolutely free worth Rs.400/-.

Offer and Subscription detail

Individual Subscriber

One year: Rs.1000/- (select any one book to receive absolutely free)

Life membership (valid for 10 years): Rs.5000/- (get both books absolutely free)

Books free for Subscribers of **Indian Journal of Trauma and Emergency Pediatrics**. Please select as per your interest. So, don't wait and order it now.

Please note the offer is valid till stock last.

CHILD INTELLIGENCE

By Dr. Rajesh Shukla

ISBN: 81-901846-1-X, Pb, vi+141 Pages

Rs.150/-, US\$50/-

Published by **World Information Syndicate**

PEDIATRICS COMPANION

By Dr. Rajesh Shukla

ISBN: 81-901846-0-1, Hb, VIII+392 Pages

Rs.250/-, US\$50

Published by **World Information Syndicate**

Order from

Red Flower Publication Pvt. Ltd.

48/41-42, DSIDC, Pocket-II, Mayur Vihar, Phase-I

Delhi - 110 091 (India)

Tel: 91-11-22754205, 45796900, Fax: 91-11-22754205

E-mail: redflowerpppl@gmail.com, redflowerpppl@vsnl.net

Website: www.rfppl.org