

Author Affiliation:

^{1,6}UG student, GCS Medical College and Hospital Ahmedabad, Gujarat 380025.

²Junior Resident,

³Associate professor, Zydus medical College and Hospital, Dahod, Gujarat 389151, India.

^{4,5}Intern Doctor, B. J. Medical College, Ahmedabad, Gujarat 380016, India

Coresponding Author:

Mukundkumar V, Associate professor in medicine, Zydus medical College and Hospital, Dahod, Gujarat 389151, India.

E-mail: drmukundvpatel123@yahoo.co.in

Isolated Tuberculosis of Ankle Joint: A Rare Case Report

Maitri M. Pate¹, Dhruvkumar M², Mukundkumar V³, Palak J Patel⁴, Mauravi V Patel⁵, Harsh D Patel⁶

How to cite this article:

Maitri M. Patel, Dhruvkumar M. Patel, Mukundkumar V Patel, et al., Isolated Tuberculosis of Ankle Joint: A Rare Case Report. Indian J Comm Dis 2020;6(1):25-27.

Abstract

Background: Isolated tuberculosis of any joint is a rare entity and often mimics other conditions. Thus, diagnosis and timely treatment is often delayed in these cases. *Clinical case:* We here report a unique case of tuberculosis of synovial tissue of ankle in a 46 year old female. She had suffered progressive left ankle swelling for ten years and left lower limb weight bearing difficulty since last 2 years prior to the presentation. She was not having past or contact history of tuberculosis. She was not having constitutional symptoms or any evidence of pulmonary tuberculosis. The diagnosis was confirmed by MRI study of the joint and typical histopathological findings of trucut biopsy from the thickened synovium. Relevant literature pertaining to tubercular arthritis has also been reviewed at length.

Keywords: Ankle joint; Tuberculosis; Articular TB; Extra pulmonary TB; MRI of joint.

Introduction

Tuberculosis is a very common disease in the developing world and this disease may have diverse presentations. Overall in India, there is a recent trend of increasing incidence of extra-pulmonary tuberculosis, especially bone and joint tuberculosis¹. A large number of tuberculosis cases occurs in the economically productive age group and this leads to significant public health burden for any society.¹

While commoner presentations of the disease like hemoptysis or ascites are comparatively easier to diagnose, bone and joint tuberculosis often presents a diagnostic and therapeutic challenge. Especially tuberculosis of single joint is notorious

for their chronic course and delay in diagnosis.² We here present a case of isolated tuberculosis of the ankle joint from Eastern India.

Case report

A 46 year old female housewife presented with gradually progressive swelling and pain in her left ankle for the last ten years. She was unable to bear weight on that ankle for the last two years. When she presented to this hospital, the left ankle was found to be swollen (Fig. 1), tender with normal local temperature. Attempt to stand on that leg led to shooting pain. Both active and passive range of motion of the left ankle joint was limited to 20°. Joint fluctuation was negative. There was no other arthropathy anywhere else in her body. There was

no skin rash, oral ulcer, dysuria, lymphadenopathy or fever. Vital signs were stable. Neurological examination of lower limbs was completely normal. There was no bleeding manifestation. She did not have a history of diabetes or any rheumatological disorder. Also, there was no history of trauma or any surgery to that joint or in the vicinity. She had never had any contact with a tuberculosis patient.

The patient had been treated for this joint disease for a long time with analgesics. She had also received various indigenous therapies like local heat application and massage. However, she denied taking any steroids or other immunosuppressive drugs, either locally or systemically.

Initial laboratory examinations revealed hemoglobin of 8 gm/dl with total leukocyte count of 9000/ μ L (N: 70; L: 22) and platelet count of 145000/ μ L. ESR was 67 mm in the 1st hour. CRP was 23 mg/L (N< 6). X ray of the feet revealed (Fig. 2) gross osteoporosis of the bones of left feet around ankle. However, the joint margins were maintained. Mantoux test was 6 mm. Serum Anti-nuclear factor and rheumatoid factor were negative. Serum uric acid was 4.8 mg/dl. A chest X-ray did not reveal any abnormality. MRI scan of the left ankle revealed (Fig. 3) massively thickened synovium of the left ankle with intense contrast enhancement. T 1 images showed signal changes in the periarticular bones too. There was minimal collection in the joint space. Thus, joint aspiration was not attempted. Finally, a USG-guided trucut needle biopsy was taken from the synovium which showed (Fig. 4) frank caseous granuloma in the synovial tissue with Langerhans giant cells. This was suggestive of tuberculosis.

The patient was started on 4-drug anti-tubercular therapy (daily regime) with ankle orthosis. At six months' follow up, the pain and swelling has decreased. However, she still has difficulty in bearing weight on that joint.



Fig. 1: The swollen left ankle of the patient



Fig. 2: X ray of left ankle showing peri-articular osteoporosis (compare with right side)



Fig. 3: MRI of left ankle showing uniformly thickened synovium enhanced with contrast




Fig. 4: Synovial biopsy specimen showing caseous granuloma (A) and Langhans Giant cells (B)

Discussion

Arthritis in tuberculosis is caused either by direct infiltration of the joint by mycobacteria or aseptic reactive arthritis.² The latter usually tends to be polyarticular while actual tubercular infection usually involves a single joint. Osteoarticular tuberculosis can be of various types like spondylitis, peripheral arthritis, tenosynovitis, osteomyelitis or Poncet's disease.³ Isolated joints are usually involved by haematogenous spread or inoculation

from local lymph nodes, foci of tubercular osteomyelitis or other contiguous sites. Early diagnosis and full treatment can lead to complete resolution of the joint pathology. However, often tubercular monoarthritis is initially misdiagnosed as traumatic or degenerative arthritis and by the time the proper diagnosis is made, bone and cartilage in and around the joint may be destroyed irreversibly.⁴ Sometimes, tubercular osteomyelitis near a joint may also mimic arthritis.⁵

Diagnosis of tubercular arthritis is done by microbiological examination of synovial effusion or histological study of joint tissue.⁶ Unlike pleural or peritoneal fluid, tests like synovial fluid protein or ADA levels are not deemed useful. In our case, the diagnosis was clinched by finding caseous granuloma in the synovial tissue.

Tubercular arthritis can have features like pannus formation, cartilage destruction, bone erosion, abscess, periarticular tenosynovitis or bursitis and bone marrow edema.³ Many of these features are well delineated on MRI. Common MRI features of tubercular arthritis are hypertrophy and intense early contrast enhancement of synovium on T1, bright tendon sheaths on T2, abscesses, bursa around joints and invading granulation tissue forming cystic structures (hypo intense on T1 and hyperintense on T2) inside bones.³ In our case, synovial hypertrophy with contrast enhancement was seen. Sometimes, some other inflammatory arthritis like rheumatoid arthritis may have similar presentations. In such cases, subtle signs like large bone erosions, uniform synovial thickening and extra-articular cysts, as mentioned above, point in favour of tubercular aetiology.⁷

Isolated monoarticular involvement as first manifestation of tuberculosis is very rarely reported. Sometimes, tuberculosis may complicate a pre-existing joint disease.⁸ In such cases, it is very important to differentiate between a monoarticular presentation of a polyarticular disease and novel tubercular infection. Sometimes, treatment for

pre-existing arthritis with repeated local steroid injections may predispose to local tuberculosis.⁸

Conclusion

Tuberculosis of a joint without constitutional symptoms is a very rare entity. Hence internists should maintain a high degree of suspicion for this disease in unexplained chronic monoarthritis. Timely intervention can prevent subsequent loss of joint function to a great extent.

References

1. Arulprakash N, Narayanan L, Narayanan S, et al. A Young Patient with Stroke and Primary Tuberculosis. *J Neurosci Rural Pract* 2018 Oct-Dec;9(4):613-615.
2. Rajakumar D, Rosenberg AM et al. Mycobacterium tuberculosis monoarthritis in a child. *Pediatr Rheumatol Online J* 2008 Sep 18;6:15.
3. Sawlani V, Chandra T, Mishra RN, et al. MRI Features of Tuberculosis of Peripheral Joints. *Clinical Radiology* 2003 Oct;58:755-62.
4. Hunfeld KP, Rittmeister M, Wichelhaus TA, et al. Two cases of chronic arthritis of the forearm due to Mycobacterium tuberculosis. *Eur J Clin Microbiol Infect Dis* 1998 May;17:344-8.
5. Desikan P, Verma R, Tiwari K, et al. Acute Monoarthritis of the Wrist Joint: Tuberculosis or Not? *J Wrist Surg* 2016 Mar;5(1):77-9.
6. Al-Sayyad MJ, Abumunaser LA. Tuberculous arthritis revisited as a forgotten cause of monoarticular arthritis. *Ann Saudi Med* 2011 Jul-Aug;31(4):398-401.
7. Choi J, Koh SH, Hong S, et al. Rheumatoid Arthritis and Tuberculous Arthritis: Differentiating MRI Features. *American Journal of Roentgenology* 2009 Nov;193:1347-53.
8. Arthanari S, Yusuf S, Nisar M et al. Tuberculosis of the Knee Complicating Seronegative Arthritis. *The journal of Rheumatology* 2008 Jun;35:1227.