

Osteogenesis Imperfecta: A Case Report

Sanjay Sudhakar Joshi¹, Anjali Kurup², Ashishkumar Patil³

Author's Affiliation:

¹Professor and Head, ²Junior Resident, ³Senior Resident, Department of Pediatrics, ACPM Medical College, Dhule, Maharashtra 424002, India.

Corresponding Author:

Anjali Kurup, Junior Resident, Department of Pediatrics, ACPM Medical College, Dhule, Maharashtra 424002, India.

E-mail: anjalikurup55@gmail.com

Received on: 19.02.2020

Accepted on: 16.03.2020

Abstract

Three year old girl child born out of non consanguineous marriage, 4th by birth order presented with severe respiratory distress and was unable to walk since the age of 1.5 years due to multiple recurrent fractures following minor trauma. Her 3 siblings died due to similar complaint. On examination she was malnourished, frontal bossing was present, oral examination dentition normal, severe respiratory distress with b/l crepitations and gallop rhythm heard. X-ray long bones showed multiple fractures with osteopenia. 2d echo-suggestive of dilated cardiomyopathy with cardiac failure. *Results:* Based on clinical examination and radiological features, and on ruling out other differentials: metabolic bone disease, fanconi syndrome, child abuse diagnosis is osteogenesis imperfecta.

Keywords: Osteogenesis imperfect; Recurrent fractures; Dilated cardiomyopathy.

How to cite this article:

Sanjay Sudhakar Joshi, Anjali Kurup, Ashishkumar Patil. Osteogenesis Imperfecta: A Case Report. Indian J Matern Fetal Neonatal Med. 2020;7(1):35-39.

Introduction

Osteogenesis imperfecta (brittle bone disease) is the most common genetic cause of osteoporosis. It is a connective tissue disorder exhibiting a wide range of clinical severity ranging from multiple fracture in utero and perinatal death to benign nearly asymptomatic with mild predisposition to fracture and normal life span. Most of them are autosomal dominant inheritance although there are variants of autosomal recessive and de novo mutation. Mutation in one of the Type I collagen genes is commonly associated with osteogenesis imperfecta.

Case History

Three year old girl child born out of non consanguineous marriage to healthy parents, fourth by birth order presented with severe respiratory distress. She was complaining of fever, cough and

difficulty in breathing since 10 days. She was unable to walk since the age of 18 months due to multiple recurrent fractures following minor trauma. She had tenderness on touching both upper and lower limbs for which no medical help was taken. She was born full-term by spontaneous vaginal delivery without fetal distress or asphyxia, birth weight 2.5 kg. At birth no fractures were present. History of repeated lower respiratory tract infection was given. She was breastfed. With no delayed teeth eruption, and was able to sit at eight months and walk independently at 16 months and has normal cognitive development. Her 3 siblings died due to similar complaint. First girl child died at the age of 9 months, it was a sudden death, she had history of repeated lower respiratory tract infection and fractures. Second male child had history of fractures following minor trauma since the age of 1 year, developmentally he was normal, died at the age of 2 years due to severe pneumonia, was sudden death. Third girl child had similar history died at the age of 3 years.




Fig. 1: Pedigree Chart

Clinical findings

- She was short stature with stunted growth; conscious, oriented to time, place and person.
- Frontal bossing present.
- On oral examination, dentition is normal.
- Barrel shaped rib cage-Pectus carinatum leading to the presenting complain of respiratory distress.
- Harrison groove and pallor seen.
- Pitting edema present on lower limbs.
- Repeated fractures followed by healing resulting in severe limb deformities present.

Anthropometry

- Head circumference - 44 cm below 3rd percentile.
- Height - 72 cm, below 3rd percentile.
- Upper Segment - 39 cm, Lower Segment - 33 cm;
- Chest circumference - 37cm.
- Weight - 5.7 kg, Grade IV PEM according to IAP classification.

Systemic examination

- On inspection trachea is central in position, barrel shaped chest, tachypnea (RR-38/min), subcost and intercostal retraction with nasal flaring seen. Apical impulse seen in 6th IC space lateral to mid clavicular line.
- On palpation hyperdynamic apical impulse and parasternal heave felt.

- On percussion left heart border 2 cm lateral to mid clavicular line.
- On auscultation bilateral crepitations, tachycardia with gallop rhythm heard.
- Liver was 6 cm palpable below the subcostal margin.
- No evidence of hearing impairment, scoliosis and neurological examination was unremarkable.

Laboratory Investigations

- Hb - 6.3 mg/dl; MCV - 65; MCH - 20; MCHC - 32
- suggestive of Nutritional Anemia.
- ESR- 25, CRP- 1.3 mg/dl (up to 6.0 mg/dl)
- Liver function test and kidney function test was normal.
- Serum Calcium - 7.8 mg/dl (9.0-11.0).
- Alkaline Phosphatase - 325IU/L (30-300).
- Serum Phosphorous - 4.1 mg/dl (2.5-5)
- Parathyroid Hormone (PTH) - 33.4 pg/ml (12-95 pg/ml)
- 1, 25-Dihydroxyvitamin D-51.8 ng/ml (sufficient 21-100)
- Thyroid profile, phosphorous-inorganic urine spot, calcium urine spot were within normal range.
- Urine analysis for detecting glycosuria and aminoaciduria (positive dipstick for protein) was normal.



Fig. 2: X-ray right upper limb-osteopenia and displaced fracture of radio ulnar joint with multiple linear fractures.



Fig. 3: X-ray of Right lower limb shows- Multiple non-displaced linear fractures with osteopenia. Sclerosis seen with callus formation in lower one third of diaphysis of right femur suggestive of previous fractures.



Fig. 4: Chest X-ray - AP view:
• Cardiomegaly
• Prominent right superior mediastinum.

On 2D Echo -

- Ejection fraction 35%,
- Left ventricular wall thickening with irregular wall hypokinesia with Increased end-diastolic diameter.
- severe pulmonary hypertension.

Above findings are suggestive of features of Dilated Cardiomyopathy with heart failure.

Discussion

Three year old girl child presented to us in severe respiratory distress with cardiac failure due to underlying dilated cardiomyopathy. On the basis of clinical features and laboratory investigation serum calcium, phosphorus, alkaline phosphatase and parathyroid hormone (PTH); 1,25 dihydroxy vitamin D, S.creatinine and electrolytes all appeared to be within the normal range hence metabolic bone disease-rickets due to Vitamin D deficiency, phosphorus deficiency, tumor induced rickets, calcium deficiency, fanconi syndrome was ruled out. Patient is suffering from osteogenesis imperfecta with dilated cardiomyopathy, which can also be suspected in other siblings as they also had similar complaints and history of sudden death which may be due to cardiac cause.

Osteogenesis imperfecta (brittle bone disease) is the most common genetic cause of osteoporosis. The incidence is 1:20,000 and occurs in all races and ethnicity.¹ It is a connective tissue disorder exhibiting a wide range of clinical severity ranging from multiple fracture in utero and perinatal death to benign nearly asymptomatic with mild predisposition to fracture and normal life span. Most of them are autosomal dominant inheritance although there are variants of autosomal recessive and de novo mutation.^{2,3}

Osteogenesis imperfecta (OI) occurs due to mutation in one of the Type I collagen genes. Since Type I collagen is one of the main structural components of connective tissues; this disease can have various extra skeletal clinical manifestations.⁴ In general, Type I collagen is the most important constituent of different parts of the cardiovascular system, including the heart valves, chordae tendineae, fibrous rings of the heart, the interventricular septum, aorta, and most other arteries.^{5,6} The collagen fibers in the ventricular myocardium contribute to the tensile stiffness and maintain the architecture of the myocytes.⁷

Several studies have reported the association of OI with various cardiovascular abnormalities, such as abnormalities of the aortic root and valve, mitral valve abnormalities, systemic hypertension, and tetralogy of Fallot.⁸⁻¹⁰ Crosssectional studies have reported increased aortic root diameter, increased prevalence of diastolic dysfunction, and valvulopathies in patients with OI, but most of the included patients were asymptomatic despite cardiovascular pathology.¹⁰

In a study carried out by Vetter and colleagues, 58 children aged 1 to 16 years with various forms of OI were evaluated and found the septal hypertrophy and posterior left ventricular wall thickening in 40% and 68% of cases, respectively.⁹ Dilated cardiomyopathy (DCM), formerly termed congestive cardiomyopathy, is characterized by enlargement of the cardiac chambers, contractile dysfunction of the myocardium, and, in later stages, by diminished cardiac output and symptoms of congestive heart failure. Familial dilated cardiomyopathy can occur secondary to an underlying inherited disease process, such as in osteogenesis imperfecta.¹¹⁻¹³

Hence in this case death of all 3 siblings can be attributed to familial dilated cardiomyopathy due to Osteogenesis Imperfecta as all of them had history of recurrent fractures and lower respiratory tract infection and sudden death. Out of 5 siblings 4 were affected with osteogenesis imperfecta with dilated cardiomyopathy.

Conclusion

Patient is suffering from osteogenesis imperfecta with dilated cardiomyopathy, which can also be suspected in other siblings as they also had similar complaints and history of sudden death which may be due to cardiac cause. Hence it is essential for every patient of osteogenesis imperfecta screened for cardiovascular disease.

References

1. Marini JC. Osteogenesis imperfecta. In: Nelson WE, Behrman RE, Kliegman RM, Arvin AM, editors. Nelson Textbook of Pediatrics. 18th ed. Philadelphia: W.B. Saunders Company 2007. pp.2887-90
2. Ward LM, Rauch F, Travers R, et al. Osteogenesis imperfecta Type VII: An autosomal recessive form of brittle bone disease. Bone. 2002;31:12-8. [PubMed] [Google Scholar].

3. Glorieux FH, Rauch F, Plotkin H, et al. Type V osteogenesis imperfecta: A new form of brittle bone disease. *J Bone Miner Res* 2000;15:1650-8. [PubMed] [Google Scholar].
 4. Weis SM, Emery JL, Beker KD, et al. Myocardial mechanics and collagen structure in osteogenesis imperfecta Murine. *Circ Res* 2000;87(8):663-9.
 5. Millington-Sanders C, Meir A, Lawrence L, Stolinski C. Structure of chordae tendineae in the left ventricle of the human heart. *J Anat.* 1998;192 (Pt 4)(Pt 4):573-581. doi:10.1046/j.1469-7580.1998.19240573.x
 6. Vouyouka AG, Pfeiffer BJ, Liem TK, et al. Phillips The role of type I collagen in aortic wall strength with a homotrimeric J. *Vasc. Surg.*, 2001;33:1263-70.
 7. Weis JL, Emery KD, Becker DJ, McBride Jr., et al. McCulloch Myocardial mechanics and collagen structure in the osteogenesis imperfecta murine (oim) *Circ.* 2000;87:663-69. View Record in Scopus Google Scholar
 8. White NJ, Winearls CG, Smith R. Cardiovascular abnormalities in osteogenesis imperfecta. *Am Heart J* 1983;106:1416-20.
 9. Vetter U, Maierhofer B, Muller M, et al. Osteogenesis imperfecta in childhood: cardiac and renal manifestations. *Eur J Pediatr* 1989;149:184-87.
 10. Hortop J, Tsipouras P, Hanley JA, et al. Cardiovascular involvement in osteogenesis imperfecta. *Circulation* 1986;73:54-61.
 11. Ashournia H, Johansen FT, Folkestad L, et al. Bixen Heart disease in patients with osteogenesis imperfecta: A systematic review *Int. J. Cardiol* 2015;196:149-57. Article Download PDFView Record in Scopus Google Scholar
 12. Towbin JA. Molecular genetic aspects of cardiomyopathy. *Biochem Med Metab Biol* 1993;49:285-320.
 13. Emery AEH. *Duchenne Muscular Dystrophy*, 2nd ed. Oxford Monographs on Medical Genetics. Vol. 24. Oxford University Press, Oxford, England, 1993.
 14. *Harrison's Principles of Internal Medicine*, 13th ed. (Eds. Isselbacher KJ, Braunwald E, Wilson JD, Martin JB, Fauci AS, Kasper DL). McGraw-Hill, New York 1994. pp.2206-221.
-

SUBSCRIPTION FORM

I want to renew/subscribe international class journal “**Indian Journal of Maternal-Fetal and Neonatal Medicine**” of Red Flower Publication Pvt. Ltd.

Subscription Rates:

- Institutional: INR10000/USD781

Name and complete address (in capitals): _____

Payment detail:

Online payment link: <http://rfppl.co.in/payment.php?mid=15>

Cheque/DD: Please send the US dollar check from outside India and INR check from India made payable to 'Red Flower Publication Private Limited'. Drawn on Delhi branch.

Wire transfer/NEFT/RTGS:

Complete Bank Account No. 604320110000467

Beneficiary Name: Red Flower Publication Pvt. Ltd.

Bank & Branch Name: Bank of India; Mayur Vihar

MICR Code: 110013045

Branch Code: 6043

IFSC Code: BKID0006043 (used for RTGS and NEFT transactions)

Swift Code: BKIDINBBDOS

Term and condition for supply of journals

1. Advance payment required by Demand Draft payable to **Red Flower Publication Pvt. Ltd.** payable at **Delhi**.
2. Cancellation not allowed except for duplicate payment.
3. Agents allowed 10% discount.
4. Claim must be made within six months from issue date.

Mail all orders to

Subscription and Marketing Manager

Red Flower Publication Pvt. Ltd.

48/41-42, DSIDC, Pocket-II

Mayur Vihar Phase-I

Delhi - 110 091(India)

Phone: 91-11-45796900, 22754205, 22756995, Cell: +91-9821671871

E-mail: sales@rfppl.co.in