

Studies on Bee and Pollination is Changing Environment: A Review

R Padmavathi¹, P Sethuraj²

How to cite this article:

R Padmavathi, P Sethuraj/Studies on Bee and Pollination is Changing Environment: A Review/Indian Journal of Agriculture Business 2023;9(1):21-29.

Abstract

This a review of the early literature demonstrating the adequacy of pollination by local bee populations before intensive and extensive agricultural practices. Although not always the best pollinators, honey bees were used in crops when populations of other bees crashed. Publications on managing and conserving pollen bees are included because interest in sustainable agriculture and biodiversity of bees and host plants is increasing.

Keywords: Apoidea; Pollination; Sustainable; Agriculture; Environment.

INTRODUCTION

Flower visiting habits, bees (Apoidea) are usually the most effective pollinators of the numerous crops and wild plants that are not wind pollinated. The study of this important subject is difficult, due to the complex interactions between the bees, their hosts, and various environmental and economic conditions; also, existing knowledge is widely dispersed in the biological and other literature, Clements and Long, 1923; Grant, (1949)²⁸ Schmid, (1975).¹⁰⁹ This review of interesting and still timely publications was compiled because many of

the early reports are not accessible by computerized literature search, and others are in journals not searched by pollination biologists. Thus, there have been many oversights and omissions. In spite of many published reports to the contrary, most people continue to simplistically credit the common honey bee (*Apis mellifera* L) with nearly all of the insect pollination in nature Barclay and Moffett, (1984)¹⁴ and agricultural crops (Anonymous, (1973)⁴ Cheung, 1973; McGregor, 1976; Levin, 1(983)¹⁰⁴ Robinson et al,1989; Southwick and Southwick, (1992).¹¹⁷

At issue is the logical valuation of publicly shared assets (bees), most species and populations belonging to nobody, yet benefiting all of us through pollination (in economics, a condition referred to as an 'externality'). Honey bees and certain stingless honey bees were brought into domestication or management long ago, but for the purpose of their honey and wax production rather than for any superior pollinating abilities. Pollination of crops was not a consideration at that time. Because they were owned by beekeepers and thus already had value in the economic system (in economic terms, and could be managed, honey bees

Author's Affiliation: ¹Assistant Registrar, ²Assistant Professor, Department of Physical Education, Alagappa University, Karaikudi 630003, Tamil Nadu, India.

Coresponding Author: R Padmavathi, Assistant Registrar, Department of Physical Education, Alagappa University, Karaikudi 630003, Tamil Nadu, India.

E-mail: drponnuseturaj@yahoo.co.in

Received on: 14.02.2023

Accepted on: 25.03.2023

began to be used as pollinators when populations of other bees (pollen bees) and crop yields declined, due to new adverse agronomic and environmental impacts. The challenge of bee and pollination economics even launched distinguished career in economics (Voorhies et al, 1933).¹³⁴ The honey bee is a good general pollinator, but not for all crops, under all conditions, just as the prolific freshwater carp is a good fish, but others may be more suited to different environments such as the sea, tastier, or less bony. The pollination need is similar to that in fisheries and aquaculture, where, as 'free' wild fish populations decline, fish prices rise, and ever more species are brought into domestication (aquaculture). Since the Roman empire, the common carp has been the world's major cultured food fish, but due to recently reduced populations of wild fish there are now many more.

The rational assessment and valuation of things that have been historically considered 'free', such as clean air, adequate water, fish, and pollinators, has become a major concern in ecology, as scarcities of such things increase (Meyer and Turner, (1992)⁷⁸ Vitousek, (1994).¹³² It seems to be human nature to unwittingly use up or destroy our resources, unless they can be specifically identified as worth saving, for economic reasons, or as ultimately beneficial to human life.

Thus, it is time to protect our native beneficial Apoidea (**Honeybee**) through habitat conservation and sustainable agriculture, and to augment them with selected, managed pollinators that are best adapted to most efficiently pollinate the flowers of our crops.

DELAYED RECOGNITION

The bias in favor of maladapted but manageable honey bees is clearly illustrated by alfalfa (lucerne) pollination. Henslow (1867)⁴⁸ first reported the tripping mechanism of the flower, and he noted that honey bees took only nectar, without tripping (pollinating). This was confirmed repeatedly (Muller, (1873)⁸⁴ Brand and Westgate, 1909;¹³⁶ Westgate, 1911; Piper et al, (1914)⁴² Hadfield and Calder, 1936; Tysdal, 1940; Knowles, 1943; Linsley, (1946)⁶⁹ Akerberg and Lesins, (1949)⁶⁷ Stephen, (1955)¹¹⁹ Semmens et al, (1993)¹¹⁴ many more recent authors). Other species of bees tripped the flowers and were soon found by astute observers to be efficient pollinators (Brand and West-gate, (1909)²⁴ Piper et al, 1914; Archer, (1917)⁵ Sladen, (1918)¹¹⁶ Helmbold, 1929; Hadfield and Calder, (1936)¹⁰⁶

Salt, 1940; Tysdal, (1940)¹²⁸ Knowles, (1943)⁶³ Stapel, 1943; Linsley, (1946)⁶⁹ Akerberg and Lesins, (1949).²

Recommendations for conservation of habitats where bees could nest (Vansell, (1951)³⁵ Franklin, (1952)¹¹⁹ Stephen, 1955), and the rearing or management of selected bee species (Knowles, (1943)⁶³ Peck and Bolton, (1946)⁹⁴ Bohart, 1972)²⁰ began to be made as populations of native bees crashed, due to the new use of insecticides and the continuing destruction of habitat. In 1943, a Canadian grower, WD Clarke, first raised *Megachile* for alfalfa pollination (Peck and Bolton, 1946)⁹⁴ Nevertheless, because honey bees were manageable and could distribute pollen from heat stressed, open flowers, they still were promoted by scientists and others for alfalfa pollination everywhere (Akerberg 2 and Lesins, (1949)⁶⁸ Vansell, (1951)³⁵ Franklin, (1952)¹³⁰ Anon, (1973)⁴ Levin, (1983)⁶⁸ Olmstead and Wooten, 1987; Robinson et al, 1989).¹⁰⁴ Recently, economic data confirming the superiority of managed *Megachile rotundata* F and *Nomia melanderi* Ckll were published (Olmstead and Wooten, 1987; Peterson et al, 1992; Wichelns et al, 1992).¹⁴⁰

A similar situation exists in apples and related orchard crops. Before the introduction of insecticides, early observers reported adequate pollination by native bees alone, and most of them recommended the conservation of uncultivated lands in or near orchards to provide habitat for their nesting (Sax, (1922)¹⁰⁷ Hutson, 1926; Wilson, (1926)¹⁴⁵ 1933; Atwood, (1933)⁷ Voorhies et al, 1933; Brittain and Newton, (1934)²⁵ Howlett, (1934)⁵² Musychenko, (1937)⁸³ Orsono-Mesa, 1947). Musychenko (1937)⁸³ first recommended the management of pollen bees to augment their populations, and a Japanese apple grower, E Matsuyama, seems to have been the first to do so, using *Osmia cornifrons* (Rad) in the 1930s (Maeta and Kitamura, 1981). At present, this superior bee is managed to pollinate a third of Japan's apples (Sekita and Yamada, 1993)¹¹³ and its use has spread to other countries (Parker et al, 1987).³⁹ Although polylectic honey bees are used in orchards, they obtain little nectar, and, having a long flight range, they tend to stray, to forage elsewhere (Voorhies et al, (1933)¹³⁴ Cheung, (1973)²⁷ Growers thus pay for hive rentals that may benefit their neighbors or even undesirable weeds (Voorhies et al, (1933)¹³⁴ Loring, 1981; Goltz, (1987)⁷¹ Mac-Fawn, (1993)³ Because most managed pollen bees are gentle, easy to keep, prefer the crop host, have short flight ranges, stay in the orchard, work rapidly, and collect and distribute pollen efficiently, interest in keeping them is increasing. *O lignaria*

propinqua Cr, O cornuta (Latr), and other *Osmia* spp are now being managed (Parker et al, (1987)⁸⁹ Torchio, 1990, (1991)¹⁸⁶ Bosch, (1994)²¹ Bosch and Blas, 1994; Marquez et al, 1994)⁷⁵

Blueberries and cranberries (*Vaccinium* spp) are other crops that are better pollinated by pollen bees than honey bees, as discovered by Shaw et al (1939)¹¹⁵ and repeatedly confirmed (Wood, (1961),¹⁴⁸ (1979)¹⁴⁹ Boulanger et al, 1967; Bigras-Huot and Jobin, 1972; Kevan et al, (1983)⁸⁹ Parker et al, (1987)⁸⁹ Harder and Thomson, 1989). The management of several new pollinators is recently developing (Parker et al, (1987; Batra, (1994)¹⁵ Stubbs et al, 1994). New techniques for mass rearing bumble bees (Ptacek, (1985)¹²⁹ Roseler, 1985; van Heemert et al, 1990; Vogel and Westerkamp, 1991) permit their field use in blueberries and cranberries, and their widespread use in glasshouses on other crops, such as the hothouse tomato. For best yields, these crops require buzz-pollination, which honey bees have long been known to be unable to perform (Fink, 1896; Herbst, (1918)³⁴ Orsono-Mesa, 1947; Rick, 1950)¹⁰³ Unfortunately, the findings of these authors were ignored for decades, and instead of efforts to manage the best pollinators, honey bees were used.

Different pollinators have different pollinating abilities, which vary with the bee species (its size, hairiness, quickness, fidelity, longevity, learning ability, flight range, cold tolerance, season, flower handling ability, etc). The appropriate match between each bee species and its host is critical, because a poorly adapted pollinator may even function as a parasite by removing nectar and pollen, without achieving pollination. Some early reports were relatively neglected (Allard, 1911; Green, (1955)⁴¹ Fronk and Slater, 1956; Medler, (1956),⁷⁷ but as knowledge of ecosystem dynamics and change increased, more attention was paid to this subject, especially the relationships between native bees, honey bees and plants (Primack and Silander Jr, (1975)¹⁶⁸ Schemske and Horvitz, 1984; Herrera, 1987, (1989)⁵⁰ Harder and Thomson, (1989)⁴⁶ Thomson and Thomson, 1989, (1992)¹⁵⁰ Young and Stanton, 1990; Westerkamp, 1991a,b; Willmer, (1991)^{144, 146} Wilson and Thomson, (1991)²⁶ Chagnon et al 1993; Kunin, (1993)⁶⁶ Cresswell, 1994; Marquez et al, 1994a,b)⁷⁵

The Conservation of Native Bees

Because unmanaged native bees are so important as pollinators of crops and wild plants, and interest in maintaining biodiversity is growing (Prescott-Allen and Prescott Allen, (1986)³ Parker et al, 1987; Torchio, (1991)¹²⁵ general awareness of the diversity

of bee species and public knowledge of their value to ecosystems is increasing (Morse, (1960) 81 Malyshev, 1963; Michener, 1979; Schreck and Schedl, (1979)¹¹⁰ Westrich, 1983b, 1990; MacKenzie and Winston, (1984)¹¹² Scott Dupree and Winston, 1987; Zockler, 1988; Banaszak, (1989a)¹² O'Toole, (1993)¹ Adams and Senft, 1994). The impacts of various human activities on populations of bees and on their habitats have been studied, especially in Europe, where long and detailed records have been kept (Westrich, 1990).¹³⁹ Impacts in Europe include forestry, cultivation patterns, pollution, drainage, irrigation, construction, parks as refuges, urbanization, herbicides and insecticides, road traffic, weed invasions, and host plant destruction (Emeis, (1964)³³ Pawlikowski, (1967)⁹¹ Benedek, (1970)¹⁷, 1972; Haeseler, (1972)⁴³, 1978, 1982; Peters, (1972)⁵⁴ Jacob-Remacle, 1976, 1984, 1992; Williams, (1982)¹⁴¹, 1986, 1989; Banaszak, (1983)¹³, 1985, 1987, (1989b)¹² Prescott-Allen, 1986; Banaszak and Manole, (1987)⁶⁴ Kosior, 1987; Tanacs, 1987; Archer, (1989)⁵ Kratochwil and Klatt, 1989; Pawlikowski, 1989a, b; Torres et al, (1989)^{127,37} Fussell and Corbet, 1991; Schwenninger, 1992; Westrich, 1983a; Osborne et al, (1991)⁶⁷ Vogel and Westerkamp, 1991; Lagerlof et al, 1992; Donath, 1994).³⁰

Recent studies have documented similar impacts and population changes in Japan (Munakata, (1984)⁸² and in North and Central America (Plowright et al, (1978)⁹⁸ Miliczky and Osgood, 1979; Wood, (1979)¹⁴⁸ Ginevan et al, (1980)³⁸ Johansen et al, (1983)⁵⁸ Hansen and Osgood, 1984; Thomson et al, 1985; Kevan et al, (1990b)⁶⁰ Vinson and Frankie, 1990; Johansen and Mayer, (1990)⁶² Kevan et al 1990a).⁶¹

Exotic Pollinators

The accidental or deliberate importation of pollinators into new geographic regions has lately become a concern to some environmentalists. The widespread European honey bee and bumble bees imported into some regions (Dunning, 1886)³² may be displacing native bees or other pollinators of the native flora, or they may provide unsatisfactory pollination of these plants (Pyke and Balzer, 1983; Vogel and Westerkamp, 1991; Westerkamp, 1991; Wilson and Thomson, 1991; Thomson and Thomson, 1992; Kato, 1993; Paton, 1993). *Bombus terrestris* L is now widely distributed for pollination of crops and is another potential competitor (Donovan, 1990; Semmens et al, 1993; Kato, 1993).⁵⁹ Although several species of solitary bees have been distributed to new regions for pollinating crops,

they are less likely to displace native bees than is the ubiquitous European honey bee, due to their greater host specificity, climatic limitations, shorter foraging ranges, specific conditions for nesting, and brief adult life (Donovan, 1990).³¹ The principles that apply to the importation and establishment of exotic pollinators resemble those of classical biological control (Batra, 1982),¹⁶ in which beneficial organisms are sought near the center of origin of the problem causing organism in a similar climate, observed and tested for lack of harm to other beneficial or rare organisms, imported into quarantine where their parasites can be eliminated, tested with hosts again, in confinement and then released for a specific purpose into their new environments. Considerable effort has gone into identifying the most efficient pollinators of several Eurasian crops that may be worthy of importation into North America, South America, Australia and New Zealand where native pollinators are not well adapted to pollinate these crops (reviewed in Parker et al, 1987).⁸⁹ Conversely, American pollinators of sunflowers, cotton, passion fruit, squashes and gourds, avocado, tomato, chili, blueberries and cranberries have been studied in areas where these crops are believed to have originated, the goal being to introduce their pollinating bees into areas where the crops are now grown without their most efficient pollinators (Parker et al, 1987).⁸⁹

CONCLUSION

Large human populations can currently only be maintained by agriculture, which must alter native environments. Humans, like all organisms, may be expected to increase in population (absent natural enemies), until all available resources are fully exploited, either directly (as in agriculture), or indirectly, for example, when biota and other natural resources have been assessed and then reserved for possible future value (in economic terms, internalized). The impact of humans is large, complex, and irreversible (Vitousek, 1994).¹³² One way to provide for more habitat in which to conserve native bees and other biodiversity, as well as nonrenewable resources, would be to improve the yields on existing crop lands. This can be accomplished, in part, by introducing the most efficient pollinators for these crops.

REFERENCES

1. Adams S, Senft D (1994) The busiest of bees. *Agric Res* 42, 8-12
2. Akerberg E, Lesins K (1949) Insects pollinating alfalfa in central Sweden. *Ann R Agric Coll Sweden* 16, 630-643
3. Allard HA (1911) Some experimental observations concerning the behavior of various bees in their visits to cotton blossoms. *Am Nat* 45, 607-685
4. Anonymous (1973) Winged nations. *Agric Res* 21, 2
5. Archer LC (1917) The production of alfalfa seed in southern Idaho. *Univ Idaho Agric Exp Sta Bull* 101, 1
6. Archer ME (1989) The wasps and bees (Hymenoptera: Aculeata) of Allertorpe Common before and after coniferization. *Naturalist (Leeds)* 114, 129-136
7. Atwood CE (1933) Studies on the Apoidea of western Nova Scotia with special reference to visitors to apple bloom. *Can J Res* 9, 443-457
8. Banaszak J (1983) Ecology of bees (Apoidea) of agricultural landscape. *Polish Ecol Stud* 9, 421-505
9. Banaszak J (1985) Rural parks as refuges of the fauna of insects which pollinate plants. *Chronmy Przyr Ojcz* 41, 5-9
10. Banaszak J (1987) Fauna of wild bees of Wielkopolska lowland during the last 50 years. *Badania Fizjogr Polska Zachodnia Ser C Zool* 67-77
11. Banaszak J (1989a) Investigations on natural resources of pollinators. In: *Unconventional Methods in Lucerne Breeding Proc, Medicago sativa Working Group Meeting, Radzikow, Poland*
12. Banaszak J (1989b) Protection of pollinating insects in Poland. *Kosmos* 38, 363-374
13. Banaszak J, Manole T (1987) Diversity and density of pollinating insects (Apoidea) in the agricultural landscape of Rumania. *Polskie Pismo Entomol* 57, 747-766
14. Barclay JS, Moffett JO (1984) The pollination value of honey bees to wildlife. *Am Bee J* 124, 497-498, 551
15. Batra SWT (1982) Biological control in agroecosystems. *Science* 25, 134-139
16. Batra SWT (1994) *Anthophora pilipes villosula* Sm (Hymenoptera: Anthophoridae), a manageable Japanese bee that visits blueberries

- and apples during cool, rainy, spring weather. Proc Entomol Soc Wash 96, 98-119
17. Benedek P (1970) Effects of plot size, stand density and state of flowering on wild bees pollinating lucerne. Acta Agron Acad Sci Hungaricae 19, 283-292
 18. Benedek P (1972) Possible indirect effect of weed control on population changes of wild bees pollinating lucerne. Acta Phytopathol Acad Sci Hungaricae 7, 267-278
 19. Bigras-Huot F, Jobin LJ (1972) La pollinisation du bleuets au Lac-St-Jean. I. Role de l'abeille domestique *Apis mellifera* L. Ann Soc Entomol Quebec 16, 138-169
 20. Bohart GE (1972) Management of wild bees for the pollination of crops. Annu Rev Entomol 17, 287-312
 21. Bosch J (1994) Improvement of field management of *Osmia cornuta* (Latreille) (Hymenoptera, Megachilidae) to pollinate almond. Apidologie 25, 71-83
 22. Bosch J, Blas M (1994) Foraging behaviour and pollinating efficiency of *Osmia cornuta* (Latr) and *Apis mellifera* L on almond (Hymenoptera, Megachilidae and Apidae). Appl Entomol Zoo 28-29, 1-9
 23. Boulanger LW, Wood GW, Osgood EA, Dirks CO (1967) Native bees associated with the lowbush blueberry in Maine and eastern Canada. Maine Agric Exp Sta Canada Agric Res Sta Tech Bull 26
 24. Brand CJ, Westgate JM (1909) Alfalfa in cultivated rows for seed production in semiarid regions. US Dept Agric Bur Industry Circular No 24, 1-23
 25. Brittain WH, Newton DE (1934) Further observations on the pollen constancy of bees. Can J Res 10, 255- 263
 26. Chagnon M, Gingras J, de Oliveira D (1993) Complementary aspects of strawberry pollination by honey and indigenous bees (Hymenoptera). J Econ Entomol 86, 416-420
 27. Cheung SNS (1973) The fable of the bees: an economic investigation. J Law Econ 16, 11-33
 28. Clements FE, Long FL (1923) Experimental pollination, an outline of the ecology of flowers and insects, Carnegie Inst, Washington, DC, USA
 29. Cresswell JE (1994) A method for quantifying the gene flow that results from a single bumblebee visit using transgenic oilseed rape, *Brassica napus* L cv Westar. Transgenic Res 3, 1343-1377
 30. Donath H (1994) Gefährdung und Schutz unserer Hummeln. Naturschutz-Arbeit Berlin Brandenburg 21, 1-5
 31. Donovan BJ (1990) Selection and importation of new pollinators to New Zealand. NZ Entomol 13, 26-32
 32. Dunning JW (1886) The importation of humble bees into New Zealand. Trans R Entomol Soc London 6, 32-34
 33. Emeis W (1964) Beobachtungen über den Rückgang häufiger Bienenarten in Schleswig-Holstein. Faunist Mitt Norddeutschland 2, 152-154
 34. Fink B (1896) Pollination and reproduction of *Lycopersicon esculentum*. Minn Bot Stud Bull 9, 636-643
 35. Franklin WW (1952) Wild bees as supplementary pollinators. Am Bee J 92, 290-291
 36. Fronk WD, Slater JA (1956) Insect fauna of cucurbit flowers. J Kans Entomol Soc 29, 141-146
 37. Fussell M, Corbet SA (1991) Bumblebee habitat requirements: a public survey. Acta Hort 288, 159-163
 38. Ginevan ME, Lane DD, Greenberg L (1980) Ambient air concentration of sulfur dioxide affects flight activity in bees. Proc Natl Acad Sci USA 77, 5631-5633
 39. Goltz L (1987) Migratory beekeeping. Am Bee J 127, 29-46
 40. Grant V (1949) Arthur Dobbs (1750) and the discovery of the pollination of flowers by insects. Torrey 76, 217-219
 41. Green JM (1955) Bumblebees breed cotton too! What's New in Crops and Soils 7, 1-2
 42. Hadfield JW, Calder RA (1936) Lucerne (*Medicago sativa*). Investigations relative to pollination and seed production in New Zealand. NZJ Sci Tech 17, 577-594
 43. Haeseler V (1978) Zum Auftreten aculeater Hymenopteren in gestorten Hochmoorresten des Fintlandsmoores. Drosera 78, 57-76
 44. Haeseler V (1982) Ameisen, Wespen und Bienen als Bewohner gepflasterter Bürgersteige, Parkplätze und Strassen (Hymenoptera: Aculeata). Drosera 82, 17-32
 45. Haeseler V (1972) Man-made habitats (deforested area, gravel pit, city gardens and parks) as refuges for insects, exemplified by the Hym Aculeata. Zool Jb Syst 99, 133-212 (in German)
 46. Hansen RW, Osgood EA (1984) Effects of a split application of Sevin-4-oil on pollinators and fruit set in a spruce-fir forest. Can Entomol 116, 457-464
 47. Harder LD, Thomson JD (1989) Evolutionary options for maximizing pollen dispersal of animal-pollinated plants. Am Nat 133, 323-344
 48. Helmbold F (1929) Untersuchungen über

- die Befruchtungsverhältnisse, über die Bedingungen und über die Vererbung der Samenerzeugung bei Luzerne (*Medicago sativa* und Bastardluzerne). *Z Pflanzen zuchtung* 14, 113-173
49. Henslow G (1867) Note on the structure of *Medicago sativa* as apparently affording facilities for the intercrossing of distinct flowers. *J Linn Soc Bot* 9, 327-329
 50. Herbst P (1918) Über Lautau ßerungen einiger chilenischer Blumenwespen (Apidae). *Dsch Entomol Z* 93-96
 51. Herrera CM (1987) Components of pollinator 'quality': comparative analysis of a diverse insect assemblage. *Oikos* 50, 79-90
 52. Herrera CM (1989) Pollinator abundance, morphology, and flower visitation rate: analysis of the 'quality' component in a plant-pollinator system. *Oecologia* 80, 241-248
 53. Howlett FS (1934) Pollination of the apple in Ohio. *Bull Ohio Agric Exp Sta* 167, 65-70
 54. Hutson R (1926) Relation of the honeybee to fruit pollination in New Jersey. *NJ Agr Exp Sta Bull* 434, 2-32
 55. Jacob-Remacle A (1976) Une opération nichoirs artificiels pour hyménoptères dans 3 jardins de Liège. *Bull Ann Soc R Belge Entomol* 112, 219-242
 56. Jacob-Remacle A (1984) Étude écologique du peuplement d'hyménoptères aculéates survivant dans la zone la plus urbanisée de la ville de Liège. *Bull Ann Soc R Belge Entomol* 120, 241-262
 57. Jacob-Remacle A (1992) Les abeilles solitaires: des insectes pollinisateurs peu connus. *Insectes* 84, 20-22
 58. Johansen CA, Mayer DF, Eves JD, Kious CW (1983) Pesticides and bees. *Environ Entomol* 12, 1513-1518
 59. Johansen CA, Mayer DF (1990) Pollinator Protection: A Bee and Pesticide Handbook, Wicwas Press, Cheshire, CT, USA
 60. Kato M (1993) Impacts of the introduction of *Bombus terrestris* colonies upon pollination mutualism in Japan. *Honeybee Sci* 14, 110-114 (in Japanese)
 61. Kevan PG, Gadawski RM, Kevan SD, Gadawski SE (1983) Pollination of cranberries, *Vaccinium macro-carpon*, on cultivated marshes in Ontario. *Proc Entomol Soc Ontario* 114, 45-53
 62. Kevan PG, Clark EA, Thomas VG (1990a) Pollination: a crucial ecological and mutualistic link in agroforestry and sustainable agriculture. *Proc Entomol Soc Ontario* 121, 43-48
 63. Kevan PG, Clark EA, Thomas VG (1990b) Insect pollinators and sustainable agriculture. *Am J Alternative Agric* 5, 13-22
 64. Knowles RP (1943) The role of insects, weather conditions and plant character in seed setting of alfalfa. *Sci Agric* 24, 29-50
 65. Kosior A (1987) Impact of economic activity upon bumblebee *Bombus Latr* population in the west Bieszczady Mts. *Ochrona Przyrody* 45, 239-262
 66. Kratochwil A, Klatt M (1989) Bee species of ruderal sites in the city of Freiburg: Br (FRG) Submediterranean elements in mosaic habitats with high persistence. *Zool Jb Syst* 116, 379-389
 67. Kunin WE (1993) Sex and the single mustard: population density and pollinator behavior effects on seed-set. *Ecology* 74, 2145-2160
 68. Lagerlof J, Stark J, Svensson B (1992) Margins of agri-cultural fields as habitats for pollinating insects. *Agric Ecosyst Environ* 40, 117-124
 69. Levin MD (1983) Value of bee pollination to US agriculture. *Bull Entomol Soc Am* 29, 50-51
 70. Linsley EG (1946) Insect pollinators of alfalfa in California. *J Econ Entomol* 39, 18-29
 71. Loring M (1981) Bees and the Law. Dadant and Sons, Hamilton, IL, USA
 72. MacFawn D (1993) Pollination pricing. *Am Bee J* 133 845-847
 73. MacKenzie KE, Winston ML (1984) Diversity and abundance of native bee pollinators on berry crops and natural vegetation in the lower Fraser Valley, British Columbia. *Can Ent* 116, 965-974
 74. Maeta Y, Kitamura T (1981) Pollinating efficiency by *Osmia cornifrons* (Radoszkowski) in relation to required number of nesting bees for economic fruit production. *Honeybee Sci* 2, 65-72
 75. Malyshev SI (1963) Wild Pollinizers in the Service of Man, *Izd-vo Nauka, Moskva, Russia*
 76. Marquez J, Bosch J, Vicens N (1994) Pollens collected by wild and managed populations of the potential orchard pollinator, *Osmia cornuta* (Latr). *J Appl Entomol* 117, 353-359
 77. McGregor SE (1976) Insect pollination of cultivated crop Plants. Agricultural Research Service, US Dept Agric
 78. Medler JT (1956) Principles and methods for the utilization of bumblebees in cross-pollination of crops. *Proc 10th Int Congr Entomol* 4, 973-981
 79. Meyer WB, Turner BL (1992) Human population growth and global land-use/cover change. *Ann Rev Ecol Syst* 23, 39-61
 80. Michener CD (1979) Biogeography of the bees. *Ann Missouri Bot Gdn* 66, 277-347
 81. Miliczky ER, Osgood EA (1979) The effects of spraying with Sevin-4-oil on insect pollinators

- in a spruce-fir forest. Maine Agric Exp Sta Tech Bull 90, 1-21
82. Morse RA (1960) The abundance of wild bees (Apoidea) in the northeastern United States. *J Econ Entomol* 53, 679-680
 83. Munakata M (1984) Change of wild bee fauna at Narukawa between 1963 and 1973. *J Hokkaido Univ Educ Sect 2B* 34, 19-39
 84. Musychenko G (1937) On the insects pollinating fruit and berry plants and their ecology. *Kief Acad Sci Inst Zool Biol* 14, 226-229
 85. Muller H (1873) Die Befruchtung der Blumen durch Insekten und die gegenseitigen Anpassungen beider, Leipzig, Germany
 86. O'Toole C (1993) Diversity of native bees and agroecosystems. In: *Hymenoptera and Biodiversity* (P LaSalle, ID Gauld, eds), CAB International, Wallingford, UK
 87. Olmstead AL, Wooten DB (1987) Bee pollination and productivity growth: the case of alfalfa. *Am J Agric Econ* 56-63
 88. Orsono-Mesa H (1947) Observaciones antecologicas sobre recoleccion de pollen por vibracion. *Caldasia* 4, 465-467
 89. Osborne J, Williams I, Corbet S (1991) Bees, pollination and habitat change in the European community. *Bee World* 72, 99-116
 90. Parker FD, Batra SWT, Tepedino VJ (1987) New pollinators for our crops. *Agric Zool Rev* 2, 279-304
 91. Paton DC (1993) Honeybees in the Australian environment: does *Apis mellifera* disrupt or benefit the native biota? *Bioscience* 43, 95-103
 92. Pawlikowski T (1967) Influence of farming system on wild bees (Hymenoptera, Apoidea) in agricultural landscape. *Acta Univ Nicolai Copernici Biologia* 35, 153-167
 93. Pawlikowski T (1989a) The structure of wild bee communities from farming areas of different field sizes. *Acta Univ Nicolai Copernici Biologia* 33, 31-46
 94. Pawlikowski T (1989b) The structure of wild bee communities from habitats of field-pine forest ecotone. *Acta Univ Nicolai Copernici Biologia* 33, 101-109
 95. Peck O, Bolton JL (1946) Alfalfa seed production in northern Saskatchewan as affected by bees, with a report on means of increasing the populations of native bees. *Sci Agric* 26, 388-418
 96. Peters G (1972) Ursachen fur den Ruckgang der selten heimischen Hummelarten (Hym, *Bombus et Psithyrus*). *Entomol Ber* 1972, 85-90
 97. Peterson SS, Baird CR, Bitner RM (1992) Current status of the alfalfa leafcutting bee, *Megachile rotundata* as a pollinator of alfalfa seed. *Beescience* 2, 135-142
 98. Piper CV, Evans MW, McKee R, Morse WJ (1914) Alfalfa seed production; pollination studies. *Bull US Dept Agric* No 75, 1-32
 99. Plowright RC, Pendrel BA, McLaren IA (1978) The impact of aerial fenitrothion spraying upon the population biology of bumble bees in southwestern New Brunswick. *Can Entomol* 110, 1145-1156
 100. Prescott-Allen C, Prescott-Allen R (1986) *The First Resource*, Yale University Press, Hartford, CT, USA
 101. Primack RB, Silander Jr JA (1975) Measuring the relative importance of different pollinators to plants. *Nature (Lond)* 255, 143-144
 102. Ptacek V (1985) Testing three methods of rearing bumble bees. *Sbornik Vedeckych Praci Vyzkumny A Slechti telsky Ustav Picinarnsky v Troubsko u Brna* 9, 59-67
 103. Pyke GH, Balzer L (1983) The effects of the introduced honeybee (*Apis mellifera*) on Australian native bees. *Hawkesbury Agric Coll NSW* 18-22
 104. Rick CM (1950) Pollination relations of *Lycopersicon esculentum* in native and foreign regions. *Evolution* 4, 110-122
 105. Robinson WS, Nowogrodzki R, Morse RA (1989) The value of honey bees as pollinators of US crops. *Am Bee J* 129, 411-486
 106. Roseler PF (1985) A technique for year-round rearing of *Bombus terrestris* (Apidae, Bombini) colonies in captivity. *Apidologie* 16, 165-170
 107. Salt RW (1940) Utilize wild bees in alfalfa seed production. *Dom Dept Agric, Lethbridge Entomol Lab Leaflet No 10 FCII No 266*, 1
 108. Sax K (1922) Sterility relationships in Maine apple varieties. *Maine Agric Exp Sta Bull* 307, 61-76
 109. Schemske DW, Horvitz CC (1984) Variation among floral visitors in pollination ability: a precondition for mutualism specialization. *Science* 225, 519-521
 110. Schmid R (1975) Two hundred years of pollination biology: an overview. *The Biologist* 57, 26-35
 111. Schreck E, Schedl W (1979) Die Bedeutung des Wildbienen - Anteils bei der Bestäubung von Apfelblüten an einem Beispiel in Nordtirol (Osterreich). *Ber Nat Med Ver Innsbruck* 66, 95-107
 112. Schwenninger HR (1992) Investigations on the influence of intensity of cultivation in rural areas on the occurrence of insects with special reference to the wild bees (Hymenoptera: Apoidea). *Zool Jb Syst* 119, 543-561
 113. Scott-Dupree CD, Winston ML (1987) Wild bee pollinator diversity and abundance in orchard

- and uncultivated habitats in the Okanagan Valley, British Columbia. *Can Entomol* 119, 735-745
114. Sekita N, Yamada M (1993) Use of *Osmia cornifrons* for pollination of apples in Aomori prefecture, Japan. *Japan Agric Res Quart* 26, 264-270
 115. Semmens TD, Turner E, Buttermore R (1993) *Bombus terrestris* (L) (Hymenoptera: Apidae) now established in Tasmania. *J Aust Entomol Soc* 32, 346
 116. Shaw FR, Bailey JS, Bourne AJ (1939) The comparative value of honeybees in the pollination of cultivated blueberries. *J Econ Entomol* 32, 872-874
 117. Sladen FWL (1918) Pollination of alfalfa by bees of the genus *Megachile*. *Agr Gaz Canada* 5, 125-126
 118. Southwick EE, Southwick LJ (1992) Estimating the economic value of honey bees (Hymenoptera: Apidae) as agricultural pollinators in the United States. *J Econ Entomol* 85, 621-633
 119. Stapel C (1943) Über die Befruchtung der Luzerne durch Insekten in Danemark. *Entomol Medd* 23, 224-239
 120. Stephen WP (1955) Alfalfa pollination in Manitoba. *J Econ Entomol* 48, 543-548
 121. Stubbs CS, Drummond FA, Osgood EA (1994) *Osmia ribifloris biedermanni* and *Megachile rotundata* (Hymenoptera: Megachilidae) introduced into the lowbush blueberry agroecosystem in Maine. *J Kans Entomol Soc* 67, 173-185
 122. Tanacs L (1987) The trend of the composition of the wild-bee population on anthropogenic effects in the biotopes of the Kiskore storage-tank region at the Tisza-Valley. *Tiscia (Szeged)* 22, 109-119
 123. Thomson JD, Plowright RS, Thaler GR (1985) Maticil insecticide spraying, pollinator mortality, and plant fecundity in New Brunswick forests. *Can J Bot* 63, 2056-2061
 124. Thomson JD, Thomson BA (1989) Dispersal of *Erythronium grandiflorum* pollen by bumblebees: implications for gene flow and reproductive success. *Evolution* 43, 657-661
 125. Thomson JD, Thomson BA (1992) Pollen presentation and viability schedules in animal-pollinated plants: consequences for reproductive success. In: *Ecology and Evolution of Plant Reproduction: New Approaches* (R Wyatt, ed), Chapman and Hall, New York, NY, USA, 1-24
 126. Torchio PF (1990) Diversification of pollination strategies for US crops. *Environ Entomol* 19, 1649-1656
 127. Torchio PF (1991) Use of *Osmia lignaria propinqua* (Hymenoptera: Megachilidae) as a mobile pollinator of orchard crops. *Environ Entomol* 20, 590-596
 128. Torres F, Gayubo SF, Asensio E (1989) Efecto de la presión urbana sobre abejas y avispas en Salamanca. V. Superfamilia Apoidea. *Commun Inst Nac Invest Agrar Ser Recur Nat* 52, 1-49
 129. Tysdal HM (1940) Is tripping necessary for seed setting in alfalfa? *J Am Soc Agron* 32, 570-585
 130. Van Heemert C, de Ruijter A, van den Eijnde J, van der Steen J (1990) Year-round production of bumble bee colonies for crop pollination. *Bee World* 71, 54-56
 131. Vansell GH (1951) Use of honey bees in alfalfa seed production. *US Dept Agric Circ* 876, 1-11
 132. Vinson SB, Frankie GW (1990) Territorial and mating behavior of *Xylocopa fimbriata* F and *Xylocopa gualanensis* Cockerell from Costa Rica. *J Insect Behav* 13-32
 133. Vitousek PM (1994) Beyond global warming: ecology and global change. *Ecology* 75, 1861-1876
 134. Vogel S, Westerkamp C (1991) Pollination: an integrating factor of biocenoses. In: *Species Conservation: A Population-Biological Approach* (A Seitz, V Loeschoke, eds), Birkhauser Verlag, Basel, Switzerland
 135. Voorhies EC, Todd FE, Galbraith JK (1933) Economic aspects of the bee industry. *Univ Calif Agric Ext Bull* 555, 1-110
 136. Westerkamp C (1991) Honeybees are poor pollinators why? *Pl Syst Evol* 177, 71-75
 137. Westgate VL (1911) The alfalfa caterpillar. *US Dept Agric Bur Entomol Circular* No 133, 1
 138. Westrich P (1983a) Die Bienenfauna des Leudelsbach tals bei Markgroningen und ihre Veränderungen im Verlauf von 50 Jahren (Hymenoptera: Apoidea). *Jb Ges Naturk Württemberg* 138, 271-285
 139. Westrich P (1983b) Die Bienen Baden-Württembergs 1. Megachilidae (Hymenoptera: Apoidea). *Stuttgarter Beitr Naturk Ser A (Biologie)* 1-50
 140. Westrich P (1989-1990) Die Wildbienen Baden-Württembergs. Verlag Eugen Ulmer, Stuttgart, 2 vol
 141. Wichelns D, Weaver TF, Brooks PM (1992) Estimating the impact of alkali bees on the yield and acreage of alfalfa seed. *J Prod Agric* 5, 512-518
 142. Williams PH (1982) The distribution and decline of British bumble bees (*Bombus* Latr.). *J Apic Res* 21, 236-245
 143. Williams PH (1986) Environmental change and the distributions of British bumblebees

- (*Bombus Latr.*). *Bee World* 67, 50-61
144. Williams PH (1989) Bumblebees and Their Decline in Britain. Centr Assoc Beekeepers, Essex, UK
145. Willmer PG (1991) Constraints on foraging by solitary bees. In: *The Behavior and Physiology of Bees* (LJ Goodman, RC Fisher, eds), CAB International, Wallingford, UK, 121-148
146. Wilson GF (1926) Pollination of hardy fruits: insect visitors to fruit blossoms. *Ann Appl Biol* 16, 602-628
147. Wilson GF (1933) Pollination in orchards. *JR Hort Soc* 58, 125-138
148. Wilson P, Thomson JD (1991) Heterogeneity among floral visitors leads to discordance between removal and deposition of pollen. *Ecology* 72, 1503-1507
149. Wood GW (1961) The influence of honeybee pollination on fruit set of the lowbush blueberry. *Can J Plant Sci* 41, 332-335
150. Wood GW (1979) Recuperation of native bee populations in blueberry fields exposed to drift of Fenitrothion from forest spray operations in New Brunswick. *J Econ Entomol* 72, 36-39
151. Young HJ, Stanton ML (1990) Influences of floral variation on pollen removal and seed production in wild radish. *Ecology* 71, 536-547
152. Zockler C (1988) Feuchtwiesenflora und blütenbe-suchende Insekten. Ihre Bedeutung für die Grünlandextensivierung-Faun. *Okol Mitt* 6, 5-19.

