

Comparative Study of the Effect of Dexmedetomidine v/s Fentanyl on Intraoperative Hemodynamic Response in Robot Assisted Lower Abdominal Onco-Surgeries in Steep Trendlenburg Position

Rachana ND¹, Mamatha HS², Sumitha CS³, Kavitha Lakshman⁴, Namrata Ranganath⁵

¹⁻⁴Assistant Professor, ⁵Professor and Head, Department of Anaesthesiology and Pain Relief, Kidwai Cancer Institute, Dr. M.H Marigowda Road, Near Dairy Circle, Bangalore, Karnataka 560029, India.

Abstract

Introduction: Carbon dioxide insufflation for pneumoperitoneum causes increase plasma levels of catecholamines and vasopressin which leads to hemodynamic disturbances. These effects are exaggerated by the trendelenburg position and long duration of surgery in robot assisted surgeries. This study was designed to evaluate the effect of Dexmedetomidine and fentanyl on hemodynamics in robot assisted surgeries in trendelenburg position. **Methods:** After obtaining Ethical committee clearance and patients consent, randomised comparative study on 40 ASA I and II patients, who were randomly allocated to receive either dexmedetomidine or fentanyl intravenous infusion was done. Patients with ASA III & IV, on β blockers, HR < 55 bpm were excluded. Intraoperative Hemodynamics, recovery and emergence was assessed. Students 't' test was used. **Results:** The study included 40 patients undergoing robotic lower abdominal surgeries requiring steep trendelenburg positions. Dexmedetomidine group of patients showed better intra operative hemodynamics at various time intervals as compared to Fentanyl group of patients.

Keywords: Robot; Trendlenburg position; Dexmedetomidine.

How to cite this article:

Rachana ND, Mamatha HS, Sumitha CS *et al.* Comparative Study of the Effect of Dexmedetomidine v/s Fentanyl on Intraoperative Hemodynamic Response in Robot Assisted Lower Abdominal Onco-Surgeries in Steep Trendlenburg Position. Indian J Anesth Analg. 2019;6(4):1191-1196.

Introduction

Robotic assisted surgeries are becoming popular due to its several benefits like early recovery, less tissue damage, minimal incision and shorter hospital stay. Anaesthetic management has become complicated in these surgeries due to cardiopulmonary changes occurring during creation of pneumoperitoneum with CO₂ with patients in trendlenburg position and maintenance of immobility of patients

throughout the surgery. Pneumoperitoneum is associated with sympathetically mediated adverse haemodynamic effect like elevation of arterial pressure, heart rate, decrease in cardiac output due to pneumoperitoneum and increase of systemic and pulmonary vascular resistances. Peritoneal insufflations also results in ventilatory and respiratory changes and can contribute to stress response. These effects are further exaggerated by the trendlenburg position.

Dexmedetomidine, an alpha 2 selective

Corresponding Author: Mamatha HS, Assistant Professor, Department of Anaesthesiology and Pain Relief, Kidwai Cancer Institute, Dr. M.H Marigowda Road, Near Dairy Circle, Bangalore, Karnataka 560029, India.

E-mail: drmamathah@gmail.com

Received on 15.04.2019, **Accepted on** 14.05.2019



This work is licensed under a Creative Commons Attribution-NonCommercial-ShareAlike 4.0.

adrenoceptor agonists is known to maintain stable hemodynamics due to its effect on central sympatholytic system. Fentanyl being an opioid analgesic acts via opioid receptors to blunt hemodynamic responses. Opioids are known to cause acute hyperalgesia [1] in the immediate postoperative period and also ileum and urinary retention with prolonged sedation. Recent studies have shown that Dexmedetomidine has an opioid equivalent analgesic actions in laparoscopic surgeries [2,3].

The study was aimed to compare the effect of Dexmedetomidine Vs Fentanyl on hemodynamic response to tracheal intubation, following pneumoperitoneum and intraoperative period in patients undergoing robotic oncology surgeries in trendelenberg position.

Methods

After obtaining approval from hospital ethical committee and informed consent from all the patients, a prospective randomised controlled study was conducted on forty patients aged 18-60 years, with American Society Anaesthesiologists (ASA) status I and II who were undergoing elective robotic oncology surgeries requiring trendelenberg position.

Exclusion criteria were patient with acute and chronic renal failure, compromised cardiovascular function, severe deranged liver function, patients with ASA Grade III and IV, emergency cases, patients on β blocker and patients with HR <55/min.

The study drugs were then prepared as follows: 2 ml (200 μ g) of study drug Dexmedetomidine was diluted in 48ml of normal saline to make 50 ml (concentration 4 μ g/ml). 4 ml (200 μ g) of study drug fentanyl was diluted in 46 ml normal saline to make 50 ml (conc. 4 μ g/ml).

Intra Venous access was secured with a 18 G cannula and infusion of Ringer's lactate was started.

The prepared drug Dexmedetomidine or fentanyl was given as follows:

Group-D (N=20): IV Dexmedetomidine 1 μ g/kg/hr was given as loading dose over 10 minutes prior to induction and then continued intraoperatively with Dexmedetomidine of 0.5 μ g/kg/hr IV infusion.

Group-F (N=20): IV fentanyl 1 μ g/kg/hr was given as a loading dose over 10 minutes prior to induction and then continued intraoperatively with fentanyl of 0.5 μ g/kg/hr IV infusion.

Hemodynamic parameters as ECG, HR, RR, SBP,

DBP, MAP and SpO₂ were recorded at baseline.

The patient was pre-oxygenated with 100% oxygen for three minutes. Following this, Inj. Glycopyrrolate 0.2 mg IV (as and when required when HR <50/min), Inj. Ondansetron 4 mg I.V and IV Midazolam 1mg were given before induction.

Anaesthesia induced with IV Propofol 1 mg/kg and inhalation agent with vecuronium as a muscle relaxant. Intraoperatively anaesthesia maintained with Inhalational agents, Muscle relaxant infusion and Study drug infusion with Positive Pressure Ventilation.

Intraoperatively parameters like ECG, HR, SBP, DBP, MAP, SpO₂, and EtCO₂ were continuously recorded.

The above parameters were then recorded at predetermined time intervals as follows: prior to infusion of study drug, 10 minutes after the study drug, after Inj. of induction drug, after intubation, five minutes after intubation, after pneumoperitoneum, 15 minutes after pneumoperitoneum, 30 minutes after pneumoperitoneum, 45 minutes after pneumoperitoneum, 60 minutes after pneumoperitoneum, every 30 mins thereafter till the release of pneumoperitoneum and five minutes after release of pneumoperitoneum.

Adverse effects like bradycardia, tachycardia, hypotension, hypertension, nausea, vomiting, respiratory depression, if any, noted during operative procedure, were treated as follows:

Bradycardia - (HR <50/min): Inj. Glycopyrrolate 0.2 mg I.V

Tachycardia - (HR >30% above baseline value): Inj. Propofol 20 mg I.V in titrated dose

Hypotension - (SBP <60 mmHg) Inj. Ephedrine 6 mg I.V in titrated dose.

Hypertension - (SBP > 140 mmHg): Inj. Propofol 20 mg I.V in titrated dose and 0.25% bupivacaine 5ml of epidural top up and IV Fentanyl 25 microgram.

Infusion of drug (Dexmedetomidine or fentanyl) was stopped and isoflurane was discontinued 10 minutes before reversal. Residual paralysis was reversed with Inj. Neostigmine 0.05 mg/kg IV, and Inj. Glycopyrrolate 8 μ g/kg IV. Patient was extubated once extubation criterias were fulfilled.

Parameters were again recorded at five minutes after extubation.

Results

Demographic profile including age, sex, weight,

ASA physical status, duration of anaesthesia and type of operation were comparable in both groups and not of much statistically significant.

The preoperative baseline haemodynamic parameters like mean HR, SBP, DBP, SpO₂ in Group D and in Group F were not significant between two groups.

At 10 mins after Loading Dose there was 18% fall in HR, 12.5% fall in SBP, 10.5% fall in DBP and 9.5% fall in MAP in Group D as compared to 2% fall in HR, 11.3% fall in SBP, 9.6% in DBP and 10.5% fall in MAP fall in Group F patients.

After intubation there was 21% fall in HR, 13.3% fall in SBP, 9.4% fall in DBP and 9.5% fall in MAP in Group D patients as compared to 2% fall in HR, 7.6% fall in SBP, 3.6% fall in DBP, 6.7% fall in MAP in Group F patients.

After insufflation there was 29.8% fall in HR, 20% fall in SBP, 17.5% fall in DBP and 17% fall in MAP in Group D patients as compared to 12.3% fall in HR, 7.6% fall in SBP, 3.5% increase in DBP and 2% fall in MAP in Group F patients.

Throughout intraoperative period of pneumoperitoneum Group D showed significant fall in HR, SBP, DBP, MAP from baseline and more stable values at all points of time intervals where as it remains similar to baseline values or at sometimes above baseline values in Group F patients (Graph 1,2,3).

At extubation there was 24.5% fall in HR, 13.98% fall in SBP, 10.5% fall in DBP and 10.5% fall in MAP in Group D patients as compared to 5.6% fall in HR, 4.6% fall in SBP, 0% variation in DBP and 2% increase in MAP in Group F patients.

Intraoperative hypertension was found in 3 patients (15%) in Group D and 6 patients (30%) in Group F patients and 3 (50%) patients in Group F needed treatment of Hypertension.

Intraoperative Bradycardia was found in 4 (20%) patients in Group D and 2 (50%) patients required intervention.

Intraoperative tachycardia was found in 5 (25%) patients in Group F. 8 patients (40%) of patients were sedated at the end of surgery but arousable to commands with no respiratory depression in group D patients and were shifted to ICU with ET tube in situ.


In fentanyl group of patients, 4 (20%) patients were sedated and responding to painful stimuli with no respiratory depression and were shifted to ICU with ET tube in situ. (Table 1).

Table 1:


	Dexmedetomidine	Fentanyl
Intraoperative Bradycardia	4 (20%)	0
Treatment	2 (50%)	0
Intraoperative Tachycardia	0	5 (25%)
Treatment	0	0
Intraoperative Hypertension	3 (15%)	6 (30%)
Treatment	1 (5%)	4 (20%)
Sedation	8 (40%)	6 (30%)
Respiratory Depression	0	0
ECG Abnormality	0	0

Extubation response was smooth in Dexmedetomidine group of patients as compared to Fentanyl group of patients.


None of our patients in both groups have any respiratory depression. None developed any ECG abnormality.



Graph 1:



Graph 2:



Graph 3:

Discussion

Pneumoperitoneum is known to increase systemic vascular resistance, mean arterial pressures, cardiac filling pressures and cardiac index. CO₂ insufflation results in hypercarbia due to peritoneal absorption of CO₂ which results in increase in HR, BP and increased risk of arrhythmias due to sympathetic nervous system stimulation and also decreases myocardial contractility [4,5].

The hemodynamic changes are further exaggerated with the increasing degree of steep trendelenberg position [6].

The opioid free and opioid sparing analgesic techniques are known to reduce the side effects associated with opioids and hence helps in faster recovery, early ambulation, early oral intake [7]. Side effects like PONV, postoperative ileus seen in these kind of surgeries along with postoperative

hyperalgesia are known to be aggravated with the use of opioids [1].

Alpha 2 agonists are shown to be effective as sole analgesic agents intraoperatively with better and stable hemodynamic control with sympatholytic, antinociceptive and sedative properties [8, 9]. A variety of pharmacological agents namely opioids, beta blockers, calcium channel blockers, combined alpha and beta blockers, lignocaine, and alpha-2 receptor agonists have been used to maintain stable hemodynamics in the perioperative period. Fentanyl being widely used intraoperatively for control of hemodynamics, is associated with side effects such as respiratory depression and increased incidence of postoperative nausea and vomiting (PONV). No studies have been done to study the effects of Dexmedetomidine and fentanyl on the hemodynamic profile in robotic surgeries requiring steep trendelenberg positions. Hence, this study was done to compare the efficacy of the two study drugs

on the hemodynamic changes intraoperatively in robotic surgeries.

Our study demonstrates that use of Dexmedetomidine for intraoperative infusion helps attenuates stress responses to different noxious stimuli during surgery and helps maintain haemodynamic stability perioperatively.

Our observation shows significantly less increase in HR in Dexmedetomidine group as in comparison with fentanyl group after intubation, after pneumoperitoneum, intraoperative period and after extubation which is in accordance with that of other studies. This is because Dexmedetomidine effectively blunts sympatho adrenal response to intubation and has good sympatholytic activity. Suparto *et al.*, [13] concluded that both Dexmedetomidine at 1 µg/Kg and fentanyl at 1 µg/Kg given intravenously as single bolus dose prior to anaesthesia induction produced lowering of blood pressures and cardiac rates, with significantly lower mean heart rates with Dexmedetomidine i.e., 21% decrease in Dexmedetomidine group vs 2% decrease in fentanyl group.

In our study, there is less fluctuation in SBP, DBP and MAP in Dexmedetomidine group in comparison to fentanyl after intubation, after pneumoperitoneum and after extubation. This implies that Dexmedetomidine had attenuated stress response perioperatively. Similar findings were observed by Patel CR *et al.* [15] and Jayshree P Vaswani *et al.* [10] and also Feld JM *et al.* [11] in which HR and BP decreased.

Intraoperatively throughout hemodynamics was more stable in Dexmedetomidine (0.5 µg/kg infusion) group of patients when compared to Fentanyl (0.5 µg/kg infusion) group of patients which is consistent with that of Sharif SM *et al.*, [16] who observed that both Dexmedetomidine (1 µg/kg) and fentanyl (2 µg/kg), when used as premedicant before induction attenuated the haemodynamic response to pneumoperitoneum during laparoscopic surgeries.

Hall JE *et al.*, [17] in their study concluded that small doses of Dexmedetomidine led to significant sedation which could be reversed by the help of verbal or physical stimuli and it resolved completely after two hour of termination of infusion. In our study, no stastically significant sedation was seen in any patient though eight patients had sedation score ≤ 3 in Group D.

Group D patients had less extubation response as compared to Group F patients which is similar to Goyal S *et al.* [18] study.

Limitation

We have done a small study of 40 patients and could not analyse the post op analgesic requirement and total amount of the study drug consumed, cost effectiveness of the study drugs and also requirement of inhalational agents in the two study groups.

Conclusion

To conclude Dexmedetomidine causes greater attenuation of hemodynamic response to tracheal intubation, following pneumoperitoneum and perioperatively with stable hemodynamics as compared to Fentanyl in surgeries requiring pneumoperitoneum in steep trendlenberg position. Dexmedetomidine patients had better sedation score and easy arousability.

Hence, intravenous premedication with Dexmedetomidine in dose of 1µg/kg as loading dose over 10 minutes prior to induction in robotic assisted surgeries requiring steep trendlenberg position followed by 0.5 µg/kg infusion till surgery is over, may be recommended for better haemodynamic stability during perioperative period.

References

1. Angst MS, Clark JD. Opioid-induced hyperalgesia: A qualitative systematic review. *Anesthesiology* 2006;104:570-87.
2. Tufanogullari B, White PF, Peixoto MP, Kianpour D, Lacour T, Griffin J, *et al.* Dexmedetomidine infusion during laparoscopic bariatric surgery: The effect on recovery outcome variables. *AnesthAnalg.* 2008;106:1741-8.
3. Jung HS, Joo JD, Jeon YS, Lee JA, Kim DW, In JH, *et al.* Comparison of an intraoperative infusion of dexmedetomidine or remifentanyl on perioperative haemodynamics, hypnosis and sedation, and postoperative pain control. *J Int Med Res* 2011;39:1890-9.
4. Hirvonen EA, Nuutinen LS, Kauko M. Haemodynamic changes due to Trendelenburg positioning and pneumoperitoneum during laparoscopic hysterectomy. *Acta Anaesthesiol Scand.* 1995;39:949.
5. Cunningham AJ, Turner J, Rosenbaum S, Rafferty T. Transoesophageal echocardiographic assessment of haemodynamic function during laparoscopic cholecystectomy. *Br J Anaesth.* 1993;70:621.
6. Kadono Y, Yaegashi H, Machioka K, Ueno S, Miwa S, Maeda Y, Miyagi T, Mizokami A, Fujii Y, Tsubokawa T, Yamamoto K, Namiki M. Cardiovascular and

- respiratory effects of the degree of head-down angle during robot-assisted laparoscopic radical prostatectomy. *Int J Med Robot.* 2013 Mar;9(1):17-22. doi: 10.1002/rcs.1482. Epub 2013 Jan 25.
7. Cepeda MS, Farrar JT, Baumgarten M, Boston R, Carr DB, Strom BL. Side effects of opioids during short-term administration: Effect of age, gender, and race. *ClinPharmacolTher.* 2003;74:102-12.
 8. Feld JM, Laurito CE, Beckerman M, Vincent J, Hoffman WE. Non-opioid analgesia improves pain relief and decreases sedation after gastric bypass surgery. *Can J Anaesth.* 2003;50:336-41.
 9. Bhana N, Goa KL, McClellan KJ. Dexmedetomidine. *Drugs.* 2000;59:263-8.
 10. Jayshree P Vaswani *et al.* Jayshree P Vaswani *et al.*, Dexmedetomidine vs Fentanyl: Effect on the Haemodynamic Response in Laparoscopic Surgeries *Journal of Clinical and Diagnostic Research.* 2017 Sep;11(9):UC04-UC08.
 11. Feld JM, Feld JM, Hoffman WE, Stechert MM, Hoffman IW, Ananda RC. Fentanyl or dexmedetomidine combined with desflurane for bariatric surgery. *J Clin Anesth.* 2006;18:24-8.
 12. Lovina Neil *et al.* Effect of Dexmedetomidine Versus Fentanyl on Haemodynamic Response to Patients Undergoing Elective Laparoscopic Surgery: A Double Blinded Randomized Controlled Study *Journal of Clinical and Diagnostic Research.* 2017 Apr;11(4):UC01-UC04].
 13. Suparto, Flores OC, Layusa CAA. A Randomized controlled trial on the effectiveness of dexmedetomidine versus fentanyl in attenuating the sympathetic response to direct laryngoscopy and endotracheal intubation.
 14. Maj Kedoki *Indon.* 2010; 60:126-32. [Who concluded that both Dexmedetomidine at 1 µg/Kg and fentanyl at 1 µg/Kg given intravenously as single bolus dose prior to anaesthesia induction produced lowering of blood pressures and cardiac rates, with significantly lower mean heart rates with Dexmedetomidine.]
 15. Patel CR *et al.* Effect of intravenous infusion of Dexmedetomidine on perioperative haemodynamic changes and postoperative recovery: a study with entropy analysis. *Indian J. Anaesth.* 2012;56:542-46.
 16. Shareef SM *et al.* A comparative study of the haemodynamic changes with Dexmedetomidine Vs Fentanyl during laparoscopic cholecystectomies under general anaesthesia. *J of Dental and Medical Sciences.* 2016;15:01-09.
 17. Hall JE, Uhrich TD, Barney JA, Arain SR, Ebert TJ. Sedative, amnestic, and analgesic properties of small-dose dexmedetomidine infusions. *Anaesth Analg.* 2000;90:699-705.
 18. Goyal S, Gupta KK, Mahajan V. A comparative evaluation of intravenous dexmedetomidine and fentanyl in breast cancer surgery: A prospective, randomized, and controlled trial. *Anesth Essays Res.* 2017;11:611-6.
-
-