

Right Middle Cerebral Artery Infarct after Minor Head Trauma in an Infant

Shubham Tripathi¹, Arnab Nag², Indranil Das³

How to cite this article:

Shubham Tripathi, Arnab Nag, Indranil Das/Right Middle Cerebral Artery Infarct After Minor Head Trauma in an Infant/
Indian J Emerg Med 2023;9(3):95- 104.

Abstract

Ischemic stroke (IS) in the pediatric population is extremely rare. In this age group, the occurrence of IS often concurs with underlying congenital heart disease, haematological, metabolic or immunological conditions. In contrast, the association between IS and minor head injury in children has been sparse. The authors report a case of a healthy 10 yrs 8 months old male who was found to have a right middle cerebral artery territory infarct. An extensive medical workup was performed, and it was negative for any previously undiagnosed comorbidities. Given the paucity of such cases, the condition and its management are discussed in corroboration with current literature.

Keywords: Stroke in Children; Ischemic CVA; Hemiparesis; Ischemic Stroke.

INTRODUCTION

The incidence of ischemic stroke (IS) in pediatrics is rare. Conversely, in the aging population, strokes are common with well established risk factors associated with IS include nutrition, hypertension, coagulopathy disorders, carotid stenosis, and patent foramen ovale.¹ However, in young adults, the list of potential stroke causes is extensive. According to the *Toast*

(Trial of Org 10172 in Acute Stroke Treatment) criteria, both strokes of unde terminated and of other determined etiology are the most common types among them.² Broadly speaking, causative factors in children can be similar to young adults where by the diagnosis is often linked to a background of congenital heart disease, haematological and, or immunological conditions. Interestingly, there have been reports of IS associated with head injury in patients less than 12 months of age.³ The authors describe the case of a 10 year 8 months - old child who developed progressive unilateral hemiparesis secondary to a left middle cerebral artery (MCA) territory infarct, etiology unknown. Given the in frequency of such cases, the condition is discussed in corroboration with current literature.

Author's Affiliation: ¹Final Year, ²Senior Consultant, ³Consultant and Head, Department of Emergency Medicine, Medica Super Speciality Hospital, Kolkata 700099, India.

Corresponding Author: Arnab Nag, Senior Consultant, Department of Emergency Medicine, Medica Super speciality Hospital, Kolkata 700099, India.

E-mail: drarnabnag@gmail.com

Received on: 25-03-2023

Accepted on: 23-04-2023

CASE REPORT

A previously well 10 year 8 months male of non-consanguineous parents presented to the



Emergency Department. According to the history, there was sudden on set of headache more on right side of head since 2 days, associated with vomiting, left upper limb of weakness, gradually involving left lower limb. Complaint of slurring of speech with deviation of mouth on right side with dribbling of saliva. No loss of consciousness was observed. Physical examination demonstrated that he had a full Glasgow Coma Scale with bilaterally equal and reactive pupils, child was able to understand what we talk as well able to talk but in slurred speech. There was normal extraocular movement. No scalp hematoma, significant skin swelling or bruise was noted. However, his left upper limb demonstrated motor power 2 out of 5 and lower limb 3 out of 5. Muscle tone in right limbs was normal but was reduced in left upper and lower limb. He was admitted for close neuromonitoring. On the following day, he was found to have progressed left lower limb weakness (power 2 out of 5) associated with hypertonia and hyper reflexia. In addition, there was no clinical improvement of his previously documented left upper limb weakness. No neurological deficit observed on his right side. Child had a facial deviation on right side. The remainder of his cranial nerves was intact.

Cl-	104
Lactate	8.6
HCO ₃ -	20.7

A CT Scan (Fig. 2, 3) was done which showed a wedge shaped hypodensity involving right fronto-temporo-parietal region suggestive of acute infarct in right MCA artery, there was no hemorrhage seen.




Fig. 2



Fig. 3

Identifications	
Patient ID	869382
Department (Pat.)	PICU/4
Patient Last Name	[REDACTED]
Patient First Name	[REDACTED]
Sample type	Arterial
T	37.0 °C
Blood Gas Values	
pH	7.309 [7.350 - 7.450]
pCO ₂	43.5 mmHg [35.0 - 45.0]
pO ₂	110 mmHg [80.0 - 100]
Oximetry Values	
ctHb	12.8 g/dL [- -]
sO ₂	98.7 % [- -]
FO ₂ Hb _e	97.9 % [- -]
Rt-Hb _e	1.3 % [- -]
Electrolyte Values	
cK ⁺	3.6 mmol/L [3.5 - 4.5]
cNa ⁺	132 mmol/L [135 - 145]
cCa ²⁺	1.18 mmol/L [1.12 - 1.32]
cCl ⁻	104 mmol/L [98 - 107]
Metabolite Values	
cLac	8.6 mmol/L [0.4 - 2.2]
Temperature Corrected Values	
pH(T)	7.309
pCO ₂ (T)	43.5 mmHg
pO ₂ (T)	110 mmHg
Oxygen Status	
ctO ₂ e	17.7 Vol%
p50 _e	29.23 mmHg
Acid Base Status	
cBase(Ecf)c	-4.1 mmol/L
cHCO ₃ (P st)c	20.7 mmol/L

An ABG (Fig. 1) was done upon arrival to ED.

pH	7.309
PCO ₂	43.5
PO ₂	110
K ⁺	3.6
Na ⁺	132

An Magnetic Resonance Imaging (MRI) brain (Fig. 4, 5) reported restricted diffusion in the right fronto-temporo-parietal regions, in keeping with a right MCA territory infarct. Diffuse thinning of right MCA and intracranial segments of right ICA suggested Vasculitis/Moya disease. No midline shift, hydrocephalus or effacement of the basal




Fig. 4

cisterns was seen.




Fig. 5

MR angiography demonstrated no flow-limiting stenosis of the anterior or posterior cerebral circulation. No arteriovenous malformation was seen. On the next day after admission, the patient had an unfortunate event of convulsions followed by cardiac arrest and was revived with the need of dopamine and dobutamine infusion post ROSC, the patient was intubated and put on a ventilator. The patient had a cardiac arrest at night the day after admission and ROSC was not achieved.

As part of the stroke workup, a comprehensive list of relevant cardiac, hematological, immunological and metabolic investigations was performed to exclude the possibility of underlying medical conditions for the cause of his IS.

Investigation	Results	Reference Values
Hb	12.9 gm/dL	12.5 - 16.1 gm/dL
Platelet	3.25 lakhs / μ L	1.5 - 4.5 / μ L
Neutrophils	86%	40 - 80%
Lymphocytes	12%	20 - 40%
PCV	38.7%	36 - 37%
MCV	89.0 fL	78 - 95 fL
MCH	29.8 pg	26 - 32 pg
MCHC	33.3 gm/dL	32 - 36 gm/dL
RDW	16.0%	11 - 17%
RBC Count	4.34 X 10 ⁶ / μ L	4.2 - 5.6 / μ L
WBC Count	11460 cells / μ L	4000 - 10500 cells / μ L
Serum Urea	11.0 mg/dL	10 - 50 mg/dL
Serum Creatinine	0.43 mg/dL	0.7 - 1.5 mg/dL
Prothrombin Time	15.8 sec	11 - 15 sec
INR	1.22	< = 1
APTT	30.2 sec	30 - 40 sec

His collective results were also reviewed by a hematologist who was of the opinion that the latter values were likely reactive to the intracranial event and hence, deemed equivocal at this stage. A detailed family history was also taken to exclude any hereditary causes for stroke, and this was negative. The patient was started on aspirin for his IS from the day of admission.

DISCUSSION

In comparison to adults, children with strokes present different layers often have unique risk factors that are less common. In Addition, presumptive risk factors for paediatric stroke often differ in children compared with adults.^{7,8} Despite an increased incidence of paediatric stroke, there is often a delay in diagnosis, and patients may remain under and, or misdiagnosed.^{5,7} A key

contributing factor is the limited expressive and interpretive skills of symptoms presented by young children.⁹ Further more, patients may present with subtle symptoms that mimic other diseases, leading to a low level suspicion by the attending clinician.⁵ With the benefit of recent in sights, we are now aware that studies show that the risk of stroke is higher for 2 weeks after trauma. On set is frequently delayed, providing an opportunity for stroke prevention during this period.¹⁰ However, various studies report that the majority of children admitted for minor head injury generally have a good outcome.^{11,16} Presumably, this may cause potential patients to have their diagnosis of IS to be delayed. Therefore, the current challenge is to identify specific children who are at higher risk of developing IS in cohort of children diagnosed with minor head injury.

Pediatric ischemic stroke may be caused

by several factors inducing thrombo-embolic occlusions of cerebral blood vessels and the activation of a complex cascade of events resulting

in a permanent brain damage.^{7,8,11-15} The main risk factors in children older than 28 days and their respective frequencies are summarized in


Attachment A*: Risk Factors For Childhood Arterial Stroke (28 days - 18 years)

Arteriopathies (intra and extracranial) 21-53%	Moya Moya Sickle cell arteriopathy Post-varicella arteriopathy Arterial dissection Primary CNS vasculitis Para/post infectious vasculitis Fibro muscular dysplasia
Cardiac 24-31%	Congenital/acquired heart diseases Patent foramen ovale(PFO) Arrhythmia Previous cardiac surgery/catheterization Endocarditis Other
Hypercoagulable State 28-13%	Protein C deficiency Protein S deficiency Factor V Leiden mutation (G1691A) Prothrombin mutation (PTG20210A) MTHFR (C677T polymorphism) Antithrombin III deficiency Increased factor VIII Hyperlipo proteinemia (alpha) Lupus anticoagulant
Acute Systemic Conditions 22-9%	Infectious diseases Sepsis Shock Acidosis/ Anoxia Other
Haemato Oncological 9-19%	Sickle cell disease Haemolytic/iron deficiency anemia Haematological malignancy Solid extracranial tumors Other
Chronic head and neck disorders 9-10%	Brain tumors Aneurysm/arterio-venous malformations Other cranial/neck tumors Migraine Other
Genetic/metabolic 1%	Mitochondrio pathy (e.g.MELAS, POLG1 mutation) Fabry disease PHACE syndrome ACTA2/COL4A1 mutation Trisomy 21 Deficiencyof Adenosine Deaminase 2 Connective tissue disorders (e.g. Ehlers Danlos-/ Marfan/Loeys Dietz syndrome)
Other 10%	Previous brain surgery Trauma

Non atherosclerotic arteriopathies, cardiac disorders, and prothrombotic states account for most of the cases, with a variable distribution of

their frequency in different geographical areas or age range.

Data collected from the International Pediatric



Stroke Study, a worldwide prospective study on 676 pediatric patients between 0 and 18 years, evidenced a higher prevalence of acute systemic conditions (including dehydration, sepsis, fever, acidosis, shock, anoxia/asphyxia, viral gastroenteritis) in Asia and South America and a lower prevalence of arteriopathies in Asia and chronic systemic conditions (haematological, oncological, and genetic disorders) in Europe and Australia.¹⁵ Non atherosclerotic arteriopathies were the predominant aetiology in all the age ranges, with the highest prevalence in children between 5 and 9 years old, while the highest prevalence of cardiac disorders and acute conditions were reported in patients under the age of 5.^{14,16} The term “non-atherosclerotic arteriopathies” included a group of heterogeneous disorders resulting in lesions or structural abnormalities involving the cerebral blood vessels’ wall as a consequence of infectious,

parainfectious or inflammatory mechanisms but also genetic predisposition or vascular malformation (e.g., focal cerebral arteriopathy, PHACE, sickle cell disease, post-varicella arteriopathy, fibromuscular dysplasia).¹⁷ The VIPS (Vascular Effect of Infection In Pediatric Stroke Study) study identified viral infections in the prior week, recent vaccination, black ethnicity, and rural residence as risk factors for a higher occurrence of arterial ischemic stroke in children.^{18,19} Serological evidence of recent, and mostly asymptomatic, herpes virus infections were detected in 45% of the enrolled patients with a predominance of HSV1 and HSV2 over VZV (respectively 24.5% versus 11.3% of the cases).²⁰ A large prospective cohort study by De Veber *et al.* that recruited 894 children with stroke in Germany, Canada, and UK, aiming to determine the association between prothrombotic conditions and risk of recurrent episodes of stroke, evidenced


a recurrence rate of 17.9% between 1 day and 136 months after the first stroke.²¹ The following conditions were identified as independent risk factors for recurrence: antithrombin deficiency (hazard ratio 3.9; 95% confidence interval 1.4 – 10.9), increased Lipoprotein (a) (hazard ratio 2.3; 95% confidence interval 1.3–4.1) and more than one prothrombotic marker (hazard ratio 1.9; 95% confidence interval 1.1–3.2).²¹

Cardiac conditions associated with stroke encompass a wide spectrum of either acquired or congenital diseases. A study by Rodan *et al.*, that recruited 135 patients with congenital heart

diseases and a diagnosis of acute ischemic stroke from Canadian Pediatric Ischemic Stroke Registry, reported a 10 years recurrence rate of 27%.²² The most common associated risk factors included mechanical heart valve (hazard ratio = 8.8), systemic infection (hazard ratio = 5.7), and prothrombotic state (hazard ratio = 2.9).²² Asaki *et al.* identified two additional risk factors in a retrospective case control study on 52 cardiopathic children who developed an arterial ischemic stroke after invasive procedures: length of ICU hospitalization and post-procedural infections.²³

The most updated pediatric stroke guidelines

For neuro imaging the criteria is described in Attachment B.



state that it is feasible to apply r TPA in children from 2 years of age, with persistent disabling neurological deficits (e.g., Pediatric NIH Stroke Scale score ≥ 6 at the time of intervention) and radio graphically confirmed cerebral large artery occlusion, within 4.5 h of known onset of symptoms.^{2,35,36} The suggested drug dosage is the same used for adults, although according to

known agerelated differences in the fibrinolytic system with children having higher levels of tissue plasminogen inhibitor than adults, it would be reasonable to consider that children might benefit of higher dosages of r TPA.^{2,13} The lacking initiation of an anti-thrombotic treatment for stroke prevention is associated with a 1.5–2.5 fold increased risk of recurrences after a first episode.² Anti-thrombotic

therapies are contraindicated in the acute setting when haemorrhagic stroke has not been excluded and in other cases including Moya Moya disease, surgery within the previous 24th, methotrexate

toxicity, thrombocytopenia with platelet count less than 50,000/mm³, or history of heparin induced thrombocytopenia.²

ATTACHMENT B : CONTRAINDICATIONS TO THROMBOLYSIS**

- Unknown time of symptoms onset
 - Pregnancy (adolescents)
 - Clinical presentation suggestive of subarachnoid haemorrhage (SAH), even if brain imaging is negative for blood
 - Patient who would decline blood transfusion if indicated
 - History of prior intracranial haemorrhage
 - Known cerebral arterial venous malformation, aneurysm, or neoplasm
 - Persistent systolic blood pressure more than 15% above the 95th percentile for age while sitting or supine
 - Glucose less than 2.78mmol/L or more than 22.22mmol/L
 - Bleeding diathesis including platelets less than 100 000, prothrombin time (PT) more than 15s (international normalised ratio (INR) more than 1.4), or elevated activated partial thromboplastin time (aPTT) more than upper limits of the normal range
 - Clinical presentation consistent with acute myocardial infarction (MI) or post-MI pericarditis that requires evaluation by cardiology before treatment
 - Prior stroke, major head trauma, or intracranial surgery within the past three months
 - Major surgery within 10 days
 - Parenchymal biopsy within 10 days (relative contraindication)
 - Gastrointestinal or urinary bleeding within 21 days (relative contraindication)
 - Arterial puncture at non-compressible site or LP within seven days (relative contraindication). Patients who have had a cardiac catheterization via a compressible artery are not excluded
 - Patient with malignancy or within one month of completion of treatment for cancer
 - Patients with an underlying significant bleeding disorder. Patients with a mild platelet dysfunction, mild von Willebrand disease, or other mild bleeding disorders are not excluded
- Stroke related exclusion criteria:**
- Mild deficit (Paediatric National Institute of Health Stroke Scale (PedNIHSS) less than 4) at start of tPA infusion or at time of sedation for neuroimaging, if applicable
 - Severe deficit suggesting large territory stroke, with pre-tPA PedNIHSS more than 24, regardless of the infarct volume seen on neuroimaging
 - Stroke suspected to be due to subacute bacterial endocarditis, Moya Moya, sickle cell disease, meningitis, bone marrow, air, or fat embolism
 - Previously diagnosed primary angitis of the central nervous system (PACNS) or secondary central nervous system (CNS) vasculitis. Focal cerebral arteriopathy of childhood is not a contraindication
- Neuroimaging related exclusion criteria:**
- Intracranial haemorrhage on pre-treatment head CT and MRI
 - Intracranial dissection (defined as at or distal to the ophthalmic artery)
 - Large infarct volume, defined by the finding of acute infarct on MRI involving one-third or more of the complete middle cerebral artery (MCA) territory involvement
- Drug-related exclusion criteria:**
- Known allergy to recombinant tissue plasminogen activator
 - Patient who received heparin within four hours must have activated partial thromboplastin time (aPTT) in normal range
 - Low molecular-weight heparin (LMWH) within past 24 hours (aPTT and INR will not reflect LMWH effect)

Specific prevention therapies may be considered for selected aetiologies (e.g., transfusions or hydroxyurea for sickle cell disease, L-arginine for MELAS, agalsidase or migalast for Fabry disease, pyridoxine in combination with folic acid and vitamin B12, methionine restricted, cystine supplemented diet and betaine, for homocystinuria) even if adequate supporting evidences are often not available because of their rare occurrence.^{2,42} The usefulness of other specific strategies (e.g Anticoagulation for pediatric arterial dissection, patent foramen ovale closure, steroids for focal cerebral arteriopathies, or surgical strategies for moyamoya) remain controversial.²

The most recent protocols have all been

similarly developed from multidisciplinary working groups involving paediatric and adult specialists (emergency physicians, stroke unit fellows, neurologists, interventional and diagnostic neuroradiologists, anaesthesiologists).⁴³⁻⁴⁶ All the protocols were organized in three diagnostic steps based on (a) the quick assessment of the patient at the point of care (either emergency departments, intensive care units, or in-hospital wards), (b) the transfer to the MRI suite where pre-determined hyperacute MRI stroke protocols are being performed, and (c) the determination of eligibility for recanalization therapy, according to the presence/absence of inclusion/exclusion criteria (see Attachment C).

A) INCLUSION CRITERIA FOR THROMBECTOMY

1. Persistent neurological deficit (PedNIHSS ≥ 6)
2. Onset of symptoms < 6 hours
3. Radiologically confirmed large vessel occlusion (anterior or posterior circulation)
4. Age ≥ 2 years

B) AFTER-THROMBECTOMY MANAGEMENT

1. Clinical monitoring of femoral artery access
2. Forced supine decubitus for at least 8 hours
3. Blood count control after 6 hours
4. Brain MRI after 24 hours
5. Removal of the compression bondage after 12 hours
6. Inr available for emergencies

ATTACHMENT D**: DIAGNOSTIC TESTS TO INDIVIDUATE ETIOLOGY
(AFTER ACUTE PHASE OR AFTER TREATMENT)**

Blood second-level tests:

- Antibodies: aCL, $\beta 2$ GPI, LAC, ANA, dsDNA, ANCA, ENA
- C3, C4
- Total cholesterol, triglycerides, ApoA1 e B, Lipoprotein A
- Homocysteine and other specific metabolic tests (MTHFR, folate, B12, lactate, sialo transferrin isoelectrofocusing, enzymatic test for Fabry disease)
- Pathological haemoglobin
- Serology for varicella
- Ammonium
- Screening for Thrombophilia
- Pregnancy test
- Anti Xa (if monitoring EBPM)
- Urine tox screen

Medical consultations:

- Cardiological (ECG, transthoracic echocardiogram and, if necessary, transoesophageal echocardiogram)
- Infectious disease specialist (consider LP if meningoencephalitis is suspected)

Others (if indicated):

- EEG
- VEP

REFERENCES

1. Sacco RL, Kasner SE, Broderick J et al (2013) An updated definition of stroke for the 21st century: a statement for health care professionals from the American Heart Association/American Stroke Association. *Stroke* 44:2064–2089.
2. Ferriero H J, Fullerton et al (2019) Management of stroke in neonates and children a scientific statement from the American Heart Association/American Stroke Association. on behalf of the American Heart Association Stroke Council and Council on Cardiovascular and Stroke Nursing. *Stroke* 50:e00–e00.
3. Dunbar M, Kirton A (2018) Perinatal stroke: mechanisms, management, and outcomes of early cerebrovascular brain injury. *Lancet Child Adolesc Health* 2(9):666–676.
4. deVeber GA, Kirton A, Booth FA et al (2017) Epidemiology and outcomes of arterial ischemic stroke in children: the Canadian Pediatric Ischemic Stroke Registry. *Pediatr Neurol* 69:58–70.
5. Gerstl L, Weinberger R, Heinen F et al (2019) Arterial ischemic stroke in infants, children, and adolescents: results of a Germany-wide surveillance study 2015–2017. *JNeuro* 266(12):2929–2941.
6. Rambaud T, Legris N, Bejot B et al (2020) Acute ischemic stroke in adolescents. *Neurology* 94:e1–e12.
7. Grunt S, Mazenauer L, Buerki SE et al (2015) Incidence and outcomes of symptomatic neonatal arterial ischemic stroke. *Pediatrics* 135(5):e1220–e1228.
8. Mallick A, Ganesan V, Kirkham F et al (2014) Childhood arterial ischemic stroke incidence, presenting features, and risk factors: a prospective population-based study. *Lancet Neurol* 13:35–43.
9. Fullerton H J, Wu YW, Zhao S, Johnston SC (2003) Risk of stroke in children: ethnic and gender disparities. *Neurology* 61:189–194.
10. Golomb MR, Fullerton HJ, Nowak-Gottl U,

- Deveber G (2009) Male predominance in childhood ischemic stroke: findings from the International Pediatric Stroke Study. *Stroke* 40:52–57
11. Martinez-Biarge M, Ferriero DM, Cowan FM (2019) Perinatal arterial is chemic stroke. Hand book of clinical neurology, Vol.162 (3rd series) *Neonatal Neurology*.
 12. Harteman JC, Groenendaal F, Kwee A et al (2012) Risk factors for perinatal arterial ischaemic stroke in full-term infants: a case-control study 97(6):F411–F416.
 13. Royal College of Pediatrics and Child Health and the Stroke Association Stroke in childhood (2017) Clinical guidelines for diagnosis, management and rehabilitation. NICE accredited. Royal college of pediatrics and child health. *Royal College of Pediatrics and Child Health Ed. London*.
 14. Gerstl L, Weinberger R, von Kries R et al (2018) Risk factors in childhood arterial ischaemic stroke: findings from a population - based study in Germany. *Eur J Paediatr Neurol* 22(3):380–386.
 15. Mackay MT, Wiznitzer M, Benedict SL et al (2011) Arterial ischemic stroke risk factors: the International Pediatric Stroke Study. *Ann Neurol* 69(1):130–140.
 16. Felling RJ, Rafay MF, Bernard T et al (2020) Predicting recovery and outcome after pediatric stroke: results from the international pediatric stroke study. *Ann Neurol* 87(6):840–852.
 17. Numis AL, Fox CK (2014) Arterial ischemic stroke in children: risk factors and etiologies. *Curr Neurol Neurosci Rep* 14(1):422.
 18. Mackay MT, Steinlin M (2019) Recent developments and new frontiers in childhood arterial ischemic stroke. *Int J Stroke* 14(1):32–43
 19. Fullerton HJ, Hills NK, Elkind M et al (2015) Infection, vaccination, and childhood arterial ischemic stroke: results of the VIPS study. *Neurology* 85:1459–1466.
 20. Elkind MS, Hills NK, Glaser CA et al (2016) Herpesvirus infections and childhood arterial ischemic stroke: results of the VIPS study. *Circulation* 133(8):732–741.
 21. DeVeber G, Kirkham F, Shannon K et al (2019) Recurrent stroke: the role of thrombophilia in a large international pediatric stroke population. *Haematologica* 104(8):1676–1681.
 22. Rodan L, McCrindle BW, Manlhiot C et al (2012) Stroke recurrence in children with congenital heart disease. *Ann Neurol* 72(1):103–111.
 23. Asakai H, Stojanovski B, Galati JC, Zannino D, Cardamone M, Hutchinson D, Cheung MMH, Mackay MT (2017) Risk factors for peri-procedural arterial ischaemic stroke in children with cardiac disease. *Pediatr Cardiol* 38:1385–1392.
 24. Kirton A, Armstrong-Wells J, Chang T et al (2011) Symptomatic neonatal arterial ischemic stroke: the international pediatric stroke study. *Pediatrics* 128:e1402–e1410.
 25. Yock-Corrales A, Babl FE, Mosley IT et al (2011) Can the FAST and ROSIER adult stroke recognition tools be applied to confirmed childhood arterial ischemic stroke? *BMC Pediatr* 21(11):93.
 26. Mackay MT, Churilov L, Donnan GA, Babl FE, Monagle P (2016) Performance of bedside stroke recognition tools in discriminating childhood stroke from mimics. *Neurology* 86(23):2154–2161.
 27. Mackay MT, Chua ZK, Lee M et al (2014) Stroke and nonstroke brain attacks in children. *Neurology* 82:1434–1440.
 28. Mackay MT, Yock-Corrales A, Churilov L et al (2016) Differentiating childhood stroke from mimics in the emergency Department. *Stroke* 47:2476–2481.
 29. Rafay MF, Pontigon AM, Chiang J et al (2009) Delay to diagnosis in acute pediatric arterial ischemic stroke. *Stroke* 40(1):58–64.
 30. Srinivasan J, Miller SP, Phan TG, Mackay MT (2009) Delayed recognition of initial stroke in children: need for increased awareness. *Pediatrics* 124(2):e227–e234.
 31. Mirsky DM, Beslow LA, Amlie-Lefond C et al (2017) Pathways for neuro imaging of childhood stroke. *Pediatr Neurol* 69:11–23.
 32. Chen J, Licht DJ, Smith SE et al (2009) Arterial spin labeling perfusion MRI in pediatric arterial ischemic stroke: initial experiences. *J Magn Reson Imaging* 29:282–290.
 33. Zuccoli G, Fitz C, Greene S, Cummings DD et al (2018) Imaging review of common and rare causes of stroke in children. *Top Magn Reson Imaging* 27(6):463–477.
 34. Zhang X, Li C, Li Q (2016) Magnetic resonance imaging in pediatric sickle cell anemia. *Exp Ther Med* 12(2):555–558.
 35. Pacheco JT, Siepmann T, Barlinn J et al (2018) Safety and efficacy of recanalization therapy in pediatric stroke: a systematic review and meta-analysis. *Eur J Paediatr Neurol* 22(6):1035–1041.
 36. Bhatia K, Kortman H, Blair C et al (2019) Mechanical thrombectomy in pediatric stroke: systematic review, individual patient data meta-analysis, and case series. *J Neurosurg Pediatr*. 1–14.
 37. Chan AK, Hovinga CA, Gill JC et al (2015) Thrombolysis in pediatric stroke study. *Stroke* 46(3):880–885.
 38. Amlie-Lefond C, Shaw DWW, Cooper A et

- al (2020) Risk of intracranial hemorrhage following intravenous TPA (tissue-type plasminogen activator) for acute stroke is low in children. *Stroke* 51(2):542-548
39. The National Institute of Neurological Disorders and Stroke rt-PA stroke study group. (1995). Tissue plasminogen activator for acute ischemic stroke. *N Engl J Med.* 333:1581-1587.
 40. Barry M, Hallam DK, Bernard TJ, Amlie-Lefond C (2019) What is the role of mechanical thrombectomy in childhood stroke? *Pediatr Neurol* 95:19-25
 41. Sporns PB, Straeter R, Minnerup J et al (2020) Does device selection impact recanalization rate and neurological outcome? An analysis of the Save Child S Study. *Stroke* 51(4):1182-1189
 42. Testai FD, Gorelick PB (2010) Inherited metabolic disorders and stroke part 1: Fabry disease and mitochondrial myopathy, encephalopathy, lactic acidosis, and stroke like episodes. *Arch Neurol* 67:19-24
 43. Harrar DB, Salussolia CL, Kapur K et al (2020) A stroke alert protocol decreases the time to diagnosis of brain attack symptoms in a pediatric emergency department. *J Pediatr* 216:136-141.e6
 44. Tabone L, Mediamolle N, Bellesme C et al (2017) Regional pediatric acute stroke protocol: initial experience during 3 years and 13 recanalization treatments in children. *Stroke* 48:2278-2281
 45. Rivkin MJ, Bernard TJ, Dowling MM, Amlie-Lefond C (2016) Guidelines for urgent management of stroke in children. *Pediatr Neurol* 56:8-17
 46. DeLaroche AM, Sivaswamy L, Farooqi A, Kannikeswaran N (2016) Pediatric stroke clinical pathway improves the time to diagnosis in an emergency department. *Pediatr Neurol* 65:39-44
 47. Shack M, Andrade A, Shah-Basak PP et al (2017) Acute Stroke Protocol study group. A pediatric institutional acute stroke protocol improves timely access to stroke treatment. *Dev Med Child Neurol.* 59:31-37
 48. Wharton JD, Barry MM, Lee CA et al (2020) Pediatric acute stroke protocol implementation and utilization over 7 years. *J Pediatr* 220:214-220
 49. Ladner TR, Mahdi J, Gindville MC et al (2015) Pediatric acute stroke protocol activation in a children's hospital emergency department. *Stroke* 46(8):2328-2331
 50. Beslow LA, Dowling MM, Hassanein SMA et al (2018) International pediatric stroke study investigators. Mortality after pediatric arterial ischemic stroke. *Pediatrics.* 141: e20174146
 51. Fullerton HJ, Wu YW, Sidney S, Johnston SC (2007) Risk of recurrent childhood arterial ischemic stroke in a population-based cohort: the importance of cerebrovascular imaging. *Pediatrics* 119(3):495-501
 52. Rivella C, Viterbori P (2021) Executive function following pediatric stroke. *A systematic review Child Neuropsychol* 27:209-231