

Innovative Method of Filling Collagen Particles in Difficult Cavity Wounds

Sushil Nepali¹, Ravi Kumar Chittoria², Jacob Antony Chakiath³, Kanav Gupta⁴,
Padmalakshmi Bharathi Mohan⁵

How to cite this article:

Sushil Nepali, Ravi Kumar Chittoria, Jacob Antony Chakiath *et al.* Innovative Method of Filling Collagen Particles in Difficult Cavity Wounds. *Ind. Jr. Med. & Health Sci.* 2024;11(1):19–21.

Abstract

Difficult cavity wounds have undermined and tunnelling morphology of their edges or a location which is difficult to access. Collagen dressing is an interactive dressing designed for wound management. Collagen particles are designed for cavity wound management. Filling the difficult cavity wound with collagen particles has some practical problems. To overcome these problems, we have used an innovative method of filling collagen particles in difficult cavity wound. Innovative method of filling collagen particles is simple and effective solution of the problems faced in managing difficult cavity wounds.

Keywords: Difficult cavity wounds; Collagen particles; Wound bed preparation.

Key message: Innovative method of filling collagen particles is simple and effective solution of the problems faced in managing difficult cavity wounds.

INTRODUCTION

Definition of a 'wound' is: loss of continuity of the skin or mucous membrane with associated tissue loss.¹ A wound is called as 'cavity wound' if the tissue loss is beyond the subcutaneous tissue and underlying tendons, muscle, or bone are exposed.¹ 'Difficult cavity wounds' have undermined and tunnelling morphology of their

edges or a location which is difficult to access. Collagen particles are designed specifically for cavity wound management.² Filling the cavity wound with collagen particles has a practical problem of spillage and difficult packing. Also significant amount of collagen particles get lost after application of secondary dressing over the collagen filled cavity wound. To overcome these problems we have used an innovative method of putting the collagen particles in cavity wound. Through this article, we share our experience of using this innovative method in a single case.

Author Affiliation: ¹Junior Resident, Department of General Surgery, ²Professor, Head of IT Wing and Telemedicine, ^{3,4}Senior Resident, ⁵Assistant Professor, Department of Plastic Surgery, JIPMER, Puducherry 605006, India.

Corresponding Author: Ravi Kumar Chittoria, Senior Professor, Head of IT Wing and Telemedicine, Department of Plastic Surgery and Telemedicine, JIPMER, Puducherry 605006, India.

E-mail: drchittoria@yahoo.com

Received on: 07.05.2024

Accepted on: 28.06.2024

MATERIALS AND METHODS

This is a case report of the use of Innovative Method of Filling Collagen Particles in Difficult Cavity Wounds in a electrical burn wound. This study was conducted in a JIPMER, tertiary care hospital in 2024, a case of 8 years old male child with electrical burns (35%). Informed written consent was taken from the patient. The patient



had a raw area with a cavity over the scalp following the electrical burn (Fig. 1). The patient was thoroughly investigated. Wound tissue culture was sent and appropriate antibiotic therapy was given. Regular cleaning and dressings were done for two weeks but the wound failed to show any healthy granulation tissue. To promote the healing, decision was made to give trial therapy of Innovative Method of Filling Collagen Particles (Fig. 2). Under all aseptic precautions, cleaning and draping was done. Collagen granules was sprayed uniformly over the raw area with a cavity along with collagen gel and a sterile dressing was applied (Fig. 3). The collagen granules with collagen gel was applied every time the dressing was changed (twice a week). The wound was assessed weekly by clinical examination.

RESULTS

With regular filling Collagen Particles in Difficult Cavity Wounds, the wound starts granulating and size also decreased. The wound area was measured using digital planimetry. Before the application of collagen granules, it was 5.7 cm² (Fig. 1). After three weeks, no ulcer or cavity is noted (Fig. 3). The wound also developed red healthy granulation tissue. (Fig. 3)



Fig. 1: Showing condition of raw area following electrical burn at the time of presentation




Fig. 2: Showing the collagen granules used



Fig. 3: Showing application of collagen granules and collagen gel over the raw area in scalp



Fig. 4: Showing the result of the application of collagen granules

DISCUSSION

Principles of wound management are summarized by acronym 'SWCR':⁴

1. Systemic analysis of patient and wound
2. Wound bed preparation
3. Clinical decision making
4. Repair, reconstruct and rehabilitate Like any other wound, principles of wound management remain same for cavity wound also.

However, approach of management differs.¹ Wound bed preparation is a critical part of wound management before proceeding to reconstruction and rehabilitation. There are four components of wound bed preparation which can be summarized by acronym 'TIMES'⁴

1. Tissue debridement (of non-viable and infected tissue)
2. Inflammation and infection control
3. Moisture balance
4. Edge of the wound- epithelial advancement
5. Surrounding tissue (control of oedema, inflammation and venous stasis)

Once the tissue debridement is completed the focus of wound management shifts to other components of wound bed preparation. Role of wound dressing is to maintain the adequate moisture and provide favourable wound environment for epithelial advancement.

Simultaneously wound dressing should also control local infection and inflammation, while preventing further infection from outside the wound.⁵

Collagen dressing is an interactive dressing designed for wound management. It acts as a sacrificial substrate to deal with elevated levels of matrix metalloproteinases (MMPs).⁶ This help the endogenous native collagen to continue normal wound healing.⁶ Collagen dressings are available in various forms: sheet, foam, gel, particles etc. Medicated collagen particles are effective tool of cavity wound management.² Some cavity wounds are difficult to access due to their location or morphology. Filling the cavity with collagen particles has some practical problems in such wounds:

1. Significant amount of particles gets spilled out of the margins of wound while filling.
2. Cavities located in antigravity direction are further difficult to fill.
3. Packing of particles in the cavity is also important to achieve contact of collagen with maximum wound surface area. Effective packing is difficult in such cavity wounds.
4. Once cavity is filled and packed with the collagen particles, it needs to be covered with secondary dressing. Significant amount of collagen gets lost in contact with the secondary dressing. Innovative method of collagen filling proposed in this article deals with all above problems. The

method is simple, effective and reproducible. It can be used for other cavity wounds like pressure sores, diabetic foot, burst abdomen etc. Limitation of this method is that it increases cost of the dressing. This study does not compare various options of cavity wound management. Randomized control trials for validation of effectiveness of this method are required

CONCLUSION

Innovative method of filling collagen particles is simple and effective solution of the problems faced in managing difficult cavity wounds.

REFERENCES

1. Smith N, Overland J, Greenwood J. Local management of deep cavity wounds—current and emerging therapies. *Dove Press*. 2015 Jan 1;2015(2):159-70.
2. Westgate S, Cutting KF, DeLuca G, Asaad K. Collagen dressings made easy. *Wounds UK*. 2012;8(1):1-4.
3. Crockett DJ. The Millard “crane flap” for acute hand injuries. *Hand*. 1970 Apr;2(2):156-9.
4. Chittoria RK, Kumar P, Bajaj SP, Singh AK, Gupta DK. General clinical guidelines for wound management: redefining acronym SWCR. *J Soc Wound Care Res*. 2014;7:2-7.
5. Vowden K, Vowden P. *Wound dressings: principles and practice*. Surgery (Oxford). 2017 Sep 1;35(9):489-94.
6. Brett D. A review of collagen and collagenbased wound dressings. *Wounds*. 2008 Dec 1;20(12):347-56.