



# Morphological Examination of the Hair of Different Wild Animals: A Prospective Study

Chaithanya Shajan<sup>1</sup>, Anita Yadav<sup>2</sup>, Tanushree Dheer<sup>3</sup>, Harshita Verma<sup>4</sup>

## How to cite this article:

Chaithanya Shajan, Anita Yadav, Tanushree Dheer *et al.* Morphological Examination of the Hair of Different Wild Animals: A Prospective Study. *Jr of Clin Forensic Sci.* 2024;2(1):11–15.

## Abstract

The present study was conducted to identify morphological and numerical features of hair samples of wild animals using light and scanning electron microscope to make a key for their identification. Hair samples were collected from different body regions of different species of animals, and animal hair structures were analyzed. Animals of the same species are more likely to have similar scale patterns. The guard hair diameter, color, shaft, cuticle pattern, the medulla, the scale patterns, and the root were examined using a light microscope. In the present study, different animal samples are collected from Mathikettan Shola National Park and Zoological Park in Thiruvananthapuram, Kerala. The samples were collected by using brushes and forceps, and each sample was packed separately in zip-lock bags to prevent mixing and preserve the integrity of the sample. Photographs of every sample were taken for visual reference, and they were taken to the laboratory for further examination. The analysis of the surface cuticular pattern, cross section, and medullary index provides information regarding the wild species of the animals.

**Keywords:** Animal Hair; Photography; Microscope; Medullary Index; Color.

## INTRODUCTION

Hair samples are the most compelling evidence at crime scenes involving wildlife. These are important for species identification in wildlife forensic studies.<sup>1</sup> The chemical and histological composition of hair has remained constant over

the years.<sup>2</sup> Hair color, texture, scale margin, scale spacing, and scale pattern showed significant differences.<sup>3</sup> The combination of these features is important when identifying species. Animal hair is called fur, while human hair is called hair.<sup>4</sup>

Macroscopic and microscopic anatomical features of animals are often used to identify hairs and to study how animals adapt to their environment. Each species has its own unique pattern.<sup>5</sup> Many scientists have made practical applications of hair identification in biology and forensics.<sup>6</sup> Hair testing can also be useful in determining whether the individual was poisoned or under the influence of drugs.<sup>7</sup> Hair identification provides excellent and up-to-date information quickly.

Wild animals have different types of hair, including feathers, bristles, spines, quills, and fur.<sup>8</sup> Each breed is developed for the animal's lifestyle and environment and serves a specific function.<sup>9</sup>

**Author's Affiliation:** <sup>1</sup>M.Sc Student, <sup>2</sup>Associate Professor, <sup>3,4</sup>Assistant Professor, Department of Forensic Science, Sanjeev Agrawal Global Educational University, Bhopal 462022, Madhya Pradesh, India.

**Correspondence:** Anita Yadav, Associate Professor, Department of Forensic Science, Sanjeev Agrawal Global Educational University, Bhopal 462022, Madhya Pradesh, India.

**E-mail:** anitakakas7@gmail.com

**Received on:** 01.08.2023

**Accepted on:** 15.12.2023



Creative Commons Non Commercial CC BY-NC: This article is distributed under the terms of the Creative Commons Attribution NonCommercial 4.0 License (<http://www.creativecommons.org/licenses/by-nc/4.0/>) which permits non-Commercial use, reproduction and distribution of the work without further permission provided the original work is attributed as specified on the Red Flower Publication and Open Access pages (<https://www.rfppl.co.in>)

Many animals undergo a moulting process in which they shed old, damaged hair and grow new hair.<sup>10</sup> It is possible to maintain the functionality and quality of the coat.<sup>11</sup> This extraordinary adaptability allows these organisms to live in their environment and perform their ecological duties.

Wild animals have unique adaptations for their hair.<sup>12</sup> In colder climates, some animals develop thicker and longer fur during the winter to provide additional insulation. This is a phenomenon known as winter fur.<sup>13</sup> They use its form and colour patterns for displays and camouflage; they use it for its insulating properties, for self-defence, and even as a sensory organ.<sup>14</sup> It is important to remember that the specific characteristics of wild animal hair can vary greatly between species and evolutionary histories.<sup>15</sup> Each species has its own adaptations and traits that make its hair suitable for its environment and lifestyle.

Wild animal hair can be used to study animal populations, track migration routes, and monitor wildlife welfare.<sup>16</sup> Genetic analysis of hair samples provides insight into the genetic diversity, relatedness, and overall health of animal populations, which can aid conservation efforts and wildlife management planning.<sup>17</sup>

Hair identification plays an important role in wildlife research and is of great importance in several fields such as wildlife conservation, ecological research, and forensic investigation. It is also important for species identification and non-invasive sampling.<sup>18</sup> It helps researchers understand and protect wildlife, study ecosystems, solve crime, and make informed decisions for the well-being of animals and people. Information gained from hair samples helps scientists understand species diversity, population dynamics, ecological interactions, health conditions, and the impact of human activity on wildlife populations. This knowledge is essential for developing effective conservation strategies and ensuring the long-term survival of wildlife and their habitats.<sup>19</sup>

The current study involves shredded hair and describes the cuticular configuration of the wild animals like Deer, Sambar deer, Chital, Black buck, African buffalo, Bison, Lion, Leopard, Wild Cat, Fox, Indian Hare.

Cortellini *et al.* collected and examined the samples from 12 different species of mammals. Hair from scalp was collected. Twelve individuals from each species were taken to give required samples. For microscopic examination each hair was kept on slide and covered with a cover slip and examined under 10x.<sup>20</sup>

---

## METHODOLOGY

---

The samples of shredded hair of wild animals were collected from Mathikettan Shola National Park (Idukki district, Kerala) and Zoological Park Thiruvananthapuram, Kerala, with the permission of Munnar Wild Life Warden. It was packed separately in zip-lock packets, photographs of each sample were taken, and it was sent to the laboratory for further analysis. A total of 12 animals were chosen for the study, namely deer, sambar deer, chital, black buck, African buffalo, Bison, Lion, Leopard, Wild Cat, Fox, and Indian Hare.

---

## MICROSCOPIC EXAMINATION

---

A few strands of hair were placed parallel on the slide, and two drops of glycerin were dropped onto the hair to hold the hair in place. A coverslip was placed over the hair and scanned along the length of the hair under a compound microscope at 100x and 400x magnification to observe the morphological features of the cuticle and medulla, as well as the pigment distribution in the cortex. Focus was on the core of the hair and its size, shape, and transparency were examined. Width of the medulla was measured and the details, such as the presence or absence of constrictions were noted.

Determination of wild animal species was based on hair medullary index which requires comparison of hair sample with reference sample to know the similarities and dissimilarities of different species. Animal hair typically has a continuous, textured medulla that makes up more than one-third of the hair shaft's overall diameter.

---

## RESULTS AND DISCUSSIONS

---

After analyzing hair samples of different species it showed that the hair index refers to the presence or absence of the medulla, the column of cells that runs down the middle of the hair shaft. The medullary index is the ratio of the pith diameter to the diameter of the entire hair shaft. The medullary index can vary greatly between animal species. Some species had hairs with large, continuous piths, while others have fragmented or non-existent piths. By comparing the medullary index of a hair sample with the medullary index range of different species, it is possible to determine the species to which the hair belongs. If the medulla index falls within a range characteristic of a particular species,

this indicates that the hair sample is likely of that species. The medullary Index helped to distinguish between different animal species based on medulla size, shape, and structure. The pith can vary greatly between species and even between individuals of the same species. The medulla can have a variety of structures; for example, it can be continuous, segmented, or in specific patterns within the hair bundle. The size and shape of the marks may also vary. These features can be compared to known reference specimens to determine the likely species. The medullary index of individuals of the same species showed slight changes when body region changes, but showed marked changes at the species level. Forensic analysis of hair centers on color and structure, determined through microscopic

magnification. The hair shaft has three forensically relevant layers: the cuticle, cortex and medulla. The Wild animals had a larger hair diameter than domestic ones. No relevant differences were observed between the wild investigated species; depending on the species, the scalp shape and the outline of their margins showed some peculiarities. The medullary and cuticular indices of individuals from the same species showed very small variations by changing the body regions.

In this study, most of the animals hair structure was rough, and the medullary index showed only a slight variation among the species. Cuticle had over lapping external scales, which helped in species identification.



Fig. 1 Hair sample of Deer, Leopard, Rhinoceros, Lion




Fig. 2: Value of medullary index of different animal species

Fig. 2 represented the value and variation of the medullary index. Sambar deer had the highest medullary index value and rhinoceros had the lowest one

**Table 1:** Structural details (microscopic and macroscopic) of the specimen

| Si. No | Animal Name     | Species       | Scientific name       | Colour of hair         | Texture of hair | Medullary index | Microscopic image Of the sample                                                       |
|--------|-----------------|---------------|-----------------------|------------------------|-----------------|-----------------|---------------------------------------------------------------------------------------|
| 1      | Chital          | Axis          | Axis Axis             | Brown                  | Rough           | 0.6             |    |
| 2      | Deer            | O.Virginiana  | Odocoileus virginiana | Brown or Grey          | Rough           | 0.7             |    |
| 3      | Sambar deer     | R.unicolar    | Russa unicolor        | Greyish and brown      | Rough           | 0.9             |    |
| 4      | Black buck      | Cervicapra    | Antilope Cervicapra   | Black or brown         | Smooth          | 0.5             |    |
| 5      | African Buffalo | Caffer        | Syncerus Caffer       | Black                  | Rough           | 0.4             |    |
| 6      | Leopard         | Pardus        | Panthera Pardus       | Yellowish brown        | Smooth          | 0.5             |    |
| 7      | Lion            | Leo           | Panthera Leo          | Golden yellow or brown | Rough           | 0.4             |   |
| 8      | Wild Cat        | Silvestris    | Felis Silvestris      | Brown or black         | Smooth          | 0.4             |  |
| 9      | Fox             | Vulpes        | Vulpes Vulpes         | Brown, White or grey   | Smooth          | 0.4             |  |
| 10     | Indian Hare     | L.nigricollis | Lepus Nigricollis     | White, Brown           | Smooth          | 0.3             |  |
| 11     | Bison           | B.bison       | Bison bison           | Black or brown         | Rough           | 0.4             |  |
| 12     | Rhinoceros      | Unicornis     | Rhinoceros Unicornis  | Brownish black         | Rough           | 0.2             |  |

## CONCLUSION

The present study was conducted to examine the morphological and numerical features of hair samples of several wild animal species using light

microscopy and to find the key to their identification. Samples of hair were obtained from different body regions of Deer (Cervidae), Sambar deer (unicolor), Axis deer, Black buck (Antilope cervicapra), African buffalo (Syncerus caffer), Leopard (Panthera pardus), Lion (Panthera Leo), Wild cat (Felis Silvestris), fox



(vulpes), Indian Hare (*Lepus Nigricollis*), bison, and Rhinoceros. The medullary index of individuals of the same species showed slight changes when body regions changed but showed marked changes at the species level. This study shows that there is a wide range of variation among all species with respect to the medullary index. In the present study, the medullary index value of the lion was distributed in multicellular rows having a fine grained transparent structure, whether or not wild cats were observed to have a different structure. The hair structure of Lion, Black buck, Sambar deer, Wild cat, Fox, and Indian hare is smooth, while that of bison, rhinoceros chital, and lion is rough.

From the analysis, the medullary index of each species is different. The study gives a clear picture that Sambar Deer have the highest medullary Index value and Rhinoceros have the lowest.

## REFERENCES

- Koppikar BR, Sabnis JH. Identification of hairs of some Indian mammals. *Journal of the Bombay Natural History Society*. 1976;73(1):5-20.
- Headquarters FB, Facts Q, Wanted M, Terrorists MW, Crimes WC, Assistance V, Schemes CF. Forensic hair comparison: background information for interpretation. *Forensic Science Communications*. 2009 Apr;11(2).
- Meyer W, Seger H, Hülmann G, Neurand K. A computer-assisted method for the determination of hair cuticula patterns in mammals. *Berliner und Munchener Tierarztliche Wochenschrift*. 1997 Mar 1;110(3):81-5.
- Dziurdzik B. Key to the identification of hairs of mammals from Poland. *Acta S zoologica cracoviensia*. 1973;18(4):73-113.
- Ohtaishi N, Sheng H, editors. *Deer of China: Biology and Management: Proceedings of the International Symposium on Deer of China, Held in Shanghai, China, 21-23 November 1992*. Elsevier; 1993.
- Species Identification. *Journal of Forensic Science and Medicine*. 2019 Jan 1;5(6):20-3.
- Needle DB, Gibson R, Hollingshead NA, Sidor IF, Marra NJ, Rothenheber D, Thachil AJ, Stanhope BJ, Stevens BA, Ellis JC, Spanswick S. Atypical dermatophytosis in 12 North American porcupines (*Erethizon dorsatum*) from the northeastern United States 2010–2017. *Pathogens*. 2019 Sep 30;8(7):171.
- Feder FH, Arias P. Comparative studies of the hair of Pudupudu and European red deer. *Anatomia, Histologia, Embryologia*. 1992 Mar 1;21(8):76-81.
- Hess WM, Flinders JT, Pritchett CL, Allen JV. Characterization of hair morphology in families Tayassuidae and Suidae with scanning electron microscopy. *Journal of Mammalogy*. 1985 Feb 26;66(9):75-84.
- Superina M, Aguiar JM. A reference list of common names for the edentates. *Edentata*. 2006 May;2006(10):33-44.
- Verma K, Joshi B. Different animal species hairs as biological tool for the forensic assessment of individual identification characteristics from animals of zoological park, Pragti Maidan, New Delhi, India. *Journal of forensic research*. 2012;3(11).
- Cortellini V, Carobbio A, Brescia G, Cerri N, Verzeletti A. A Comparative Study of Human and Animal Hairs: Microscopic Hair Comparison and Cytochrome: c: Oxidase I Species Identification. *Journal of Forensic Science and Medicine*. 2019 Jan 1;5(12):20-3.
- Johnson E, Hornby J. Age and seasonal coat changes in long haired and normal Fallow deer (Damadama). *Journal of Zoology*. 1980 Dec;192(13):501-9.
- Meyer W, Seger H, Hülmann G, Neurand K. A computer-assisted method for the determination of hair cuticula patterns in mammals. *Berliner und Munchener Tierarztliche Wochenschrift*. 1997 Mar 1;110(14):81-5.
- Chernova of. Architectonics of the medulla of guard hair and its importance for identification of taxa. In *Doklady Biological Sciences* 2001 Jan (Vol. 376, pp. 81-85). Kluwer Academic Publishers-Plenum Publishers.
- Sargis EJ, Campbell KK, Olson LE. Taxonomic boundaries and craniometric variation in the treeshrews (Scandentia, Tupaiidae) from the Palawan Faunal Region. *Journal of Mammalian Evolution*. 2014 Mar;21:111-23.
- Dziurdzik B. Klucz do oznaczania włosów ssaków Polski. Zakład Zoologii Systematycznej Doświadczalnej Polskiej Akademii Nauk; 1973.
- Ahmed YA, Ali S, Ghallab A. Hair histology as a tool for forensic identification of some domestic animal species. *EXCLI journal*. 2018;17:663.
- Mariacher A, Garofalo L, Fanelli R, Lorenzini R, Fico R. A combined morphological and molecular approach for hair identification to comply with the European ban on dog and cat fur trade. *PeerJ*. 2019 Nov 11;7:e7955.
- Cortellini V, Carobbio A, Brescia G, Cerri N, Verzeletti A. A Comparative Study of Human and Animal Hairs: Microscopic Hair Comparison and Cytochrome: c: Oxidase.