

A Comprehensive Review of Autonomic Function Testing

Pooja Nigade¹, V.P. Varshney², Mona Bedi

How to cite this article:

Pooja Nigade, V.P. Varshney, Mona Bedi. A Comprehensive Review of Autonomic Function Testing. Int. Phy.2024;12(2-3):43-49.

Abstract

The Autonomic Nervous System (ANS) regulates numerous involuntary bodily functions, including cardiovascular control, digestion, and thermoregulation. Disruptions in ANS function are associated with a diverse array of neurological and systemic disorders. Traditional autonomic testing methods, such as the Valsalva manoeuvre and orthostatic tests, primarily focus on cardiovascular reflexes, offering valuable yet indirect insights into autonomic pathways. However, technological advancements have led to the emergence of modern methods like heart rate variability (HRV) analysis, microneurography, and quantitative sudomotor axon reflex testing (QSART), which provide more precise assessments of both sympathetic and parasympathetic function. Despite their potential, the integration of these contemporary methods into routine clinical practice remains limited due to their complexity and resource requirements. This review aims to provide a comprehensive overview of classical and modern ANS testing methods, emphasising their clinical relevance in daily practice to enhance patient-care.

Aims and Methods of ANS Testing: Autonomic nervous system (ANS) testing is essential for evaluating the severity and distribution of autonomic dysfunction, diagnosing conditions such as autonomic neuropathy and orthostatic intolerance, and monitoring disease progression or treatment efficacy. Most tests assess cardiovascular reflexes in response to various stimuli, activating sympathetic or parasympathetic outflow. Classical methods, including the Ewing Battery, Head-Up Tilt (HUT) test, and Deep Breathing Test, remain clinically significant, especially in settings lacking advanced technology. Contemporary methods have revolutionised ANS testing, with HRV analysis providing in-depth insights into autonomic control over heart rate, while Microneurography and QSART assess sympathetic nerve activity and sudomotor function, respectively.

Conclusion: The evolution of autonomic function testing, from simple measures to advanced techniques, enhances our understanding of ANS regulation. Classical methods remain invaluable, yet newer technologies like HRV analysis and QSART offer detailed evaluations for complex autonomic conditions. Broad clinical integration of these contemporary methods alongside traditional approaches is essential for improving patient care.

Keywords: ANS testing, ANS, Newer methods of ANS testing.

Author's Affiliation: ¹Professor, ²Senior Specialist, Department of General Medicine, Ballary Medical College and Research Centre, Centre, Ballari, Karnataka 583104, India.

Corresponding Author: Mona Bedi, Director Professor, Department of Physiology, Maulana Azad Medical College, MAMC & Associated Hospitals, New Delhi 110002, India.

E-mail: drmonabedi@gmail.com

Received on: 19-09-2024

Accepted on: 13-11-2024



This work is licensed under a Creative Commons Attribution-NonCommercial-ShareAlike 4.0.

INTRODUCTION

The Autonomic Nervous System (ANS) controls numerous involuntary bodily functions, including cardiovascular regulation, digestion, and thermoregulation. Disruptions in ANS function are linked to a variety of neurological and systemic disorders. Historically, classical ANS testing methods, such as the Valsalva manoeuvre and orthostatic tests, have been used to assess autonomic function by measuring cardiovascular reflexes¹. These methods, while still valuable, primarily provide indirect insights into autonomic pathways. With advancements in technology, newer methods like heart rate variability (HRV) analysis, microneurography, and quantitative sudomotor axon reflex testing (QSART) have emerged². These contemporary techniques offer more detailed, objective assessments of both sympathetic and parasympathetic function and provide more accurate insights into autonomic dysfunction.

Despite the growing recognition of these advanced techniques, there remains a gap in their routine clinical integration. This is partly due to the complexity and resource demands of these tests, which are often confined to research settings or specialized centres. There is a pressing need to integrate these newer methods into everyday clinical practice, where they can complement traditional tests and offer a more comprehensive understanding of autonomic function.

This review article aims to provide a concise overview of both classical and modern methods of ANS testing, highlighting their relevance, clinical applications, and the need for broader adoption of newer techniques in routine clinical practice for enhanced patient care.

Historical Perspective

Historically, the assessment of ANS function has relied heavily on indirect measures, primarily due to the system's complexity. Early methods focused on observable physiological responses to stimuli, such as the Valsalva manoeuvre, postural changes, and the response to cold pressor test (ANS). Early pioneers such as Valsalva (1707) identified autonomic control of circulation through such manoeuvres, which remain relevant today. Similarly, the works of Jackson, King, and Ranson in the late 19th and early 20th centuries laid the foundation for recognizing how neurological diseases impair circulatory reflexes³.

The early methods often involved simple yet clinically informative tests, such as measuring blood pressure changes in response to posture or isometric exercise (handgrip)³. These tests, provided valuable insights into autonomic reflexes and their dysfunction, particularly in conditions like diabetes, tabes dorsalis, and spinal cord injury.

Physiological Anatomy of ANS

The term Autonomic Nervous System (ANS) was introduced by Langley over a century ago to describe the part of the nervous system responsible for controlling visceral functions such as cardiac muscle activity³, smooth muscle contraction, and glandular secretions. Unlike the somatic nervous system, most visceral activities regulated by the ANS occur involuntarily and cannot be consciously controlled or readily altered. The primary input to the ANS comes via somatosensory nerves, which originate from interoceptors like stretch receptors and chemoreceptors in blood vessels and visceral organs, continually monitoring the internal environment. These signals, although usually not consciously perceived, elicit responses that either stimulate or inhibit target visceral structures to maintain homeostasis.

Organization of the Autonomic Nervous System

The nervous system is divided into two parts: (1) Central nervous system (CNS) and (2) Peripheral nervous system (Fig. 1). The peripheral nervous system (PNS) is further divided into somatic, autonomic and enteric nervous system. The CNS receives all the peripheral sensory information via the sensory nerve fibres, also called the afferents and sends the motor information to the effectors via motor nerve fibres, also called efferent fibres. These efferent fibres travel through the somatic and the ANS. The input for ANS is via somatic afferent pathway which provides information to the CNS that in turn stimulates the ANS. This would lead to an involuntary action. The ANS is responsible for involuntary motor responses. The effector may be smooth or cardiac muscle (both involuntary muscles) or a gland. Thus, the **ANS is a visceral efferent system**, which means it sends motor impulses from the CNS to the visceral organs. It functions automatically and continuously, without conscious effort, to innervate smooth muscle, cardiac muscle, and glands. The ANS has two parts, sympathetic division and a parasympathetic division. The sympathetic and the parasympathetic fibers are the efferents that connect CNS to the visceral organs. There

is a fine balance between these two systems is required to maintain homeostasis. ENS comprises sympathetic and parasympathetic fibers that regulate the activity of the gastrointestinal tract. The sympathetic and the parasympathetic

efferent pathways are composed of two neurons: (1) **Preganglionic Neurons**: Originating in the CNS, their axons extend to an autonomic ganglion and (2) **Postganglionic Neurons**: These neurons relay signals from the ganglion to the target organ.




Fig. 1: Organization of ANS

Functions of the ANS

- (a) **Parasympathetic System (PSS)**: The parasympathetic division is responsible for conserving and storing energy, promoting anabolic processes during restful states.
- (b) **Sympathetic System (SS)**: The sympathetic division, on the other hand, is catabolic and activates the body for emergency responses. Its effects are generally prolonged and widespread, whereas the parasympathetic responses are more localized and transient.

Aims of ANS Testing:

- To evaluate the severity and distribution of autonomic function
- To diagnose limited autonomic neuropathy
- To diagnose and assess orthostatic intolerance
- To monitor the course of dysautonomia
- To monitor response to treatment
- To serve as an instrument in research studies

Most of the tests are based on evaluation of the cardiovascular reflexes triggered by performing specific provocative manoeuvres. Stimuli that raise blood pressure, such as isometric exercise,

cold pressor test or mental arithmetic, activate mainly sympathetic outflow. Moreover, blood pressure responses to orthostatic testing and Valsalva manoeuvre are in a large part a reflection of sympathetic activity.^{4,6} HR Changes during orthostatic testing and Valsalva manoeuvre, as well as during deep breathing or diving reflex, reflect parasympathetic modulation.^{6,7}

Traditionally, batteries of autonomic tests have been introduced, with the Ewing battery being the most popular.⁸ **Ewing Battery**: The Ewing battery, developed by Ewing and Clarke, is a comprehensive set of tests used to assess both parasympathetic and sympathetic function. It focuses on cardiovascular reflex responses, including heart rate and blood pressure variations. The tests involve measuring responses to autonomic stimuli, such as deep breathing, the Valsalva manoeuvre, and postural changes. They are widely used to detect autonomic failure in diabetic patients and can reveal abnormalities related to both parasympathetic and sympathetic nervous systems.⁸

More recently, new techniques, such as evaluation of heart rate variability or microneurography, have been introduced as diagnostic tools.

Assessment of Autonomic Nervous System (ANS) Function

The assessment of ANS function has evolved significantly, progressing from classical methods to contemporary advanced techniques. Autonomic function tests (AFT) are essential in confirming and determining the severity of dysautonomia, as well as identifying structural and functional deficits in the ANS. These tests are crucial in both clinical and research settings, with various techniques developed to evaluate different autonomic pathways.

Classical Methods for ANS Testing

Older methods of ANS testing remain clinically significant, particularly in the diagnosis of autonomic dysfunctions such as diabetic neuropathy, orthostatic hypotension, and cardiovascular dysregulation. These techniques, though simpler than modern methods, continue to provide essential diagnostic insights, especially in settings where advanced technology may not be readily available.

The followings are the classical/older methods of ANS Testing:

1. **Head Up Tilt (HUT) / Lying-Standing Test (LST):** The head-up tilt and lying-standing tests evaluate the cardiovascular response to positional changes, making them critical in diagnosing orthostatic hypotension. By transitioning from lying to standing, blood pools in the lower extremities, challenging the body's ability to maintain blood pressure through autonomic reflexes. A significant drop in systolic or diastolic pressure can signal autonomic failure. HUT is often combined with other autonomic tests for a more robust evaluation of disorders like postural tachycardia syndrome (POTS)⁸.
2. **Deep Breathing Test:** This test measures

heart rate variability in response to deep breathing, exploiting the natural respiratory sinus arrhythmia. Normally, the heart rate increases during inspiration and decreases during expiration, reflecting parasympathetic activity. A reduction in heart rate variability during this test is often an indicator of impaired parasympathetic function, as seen in conditions like diabetic autonomic neuropathy⁹.

3. **Valsalva Manoeuvre** The Valsalva manoeuvre is a cornerstone of autonomic testing, used to assess baroreceptor reflexes and both parasympathetic and sympathetic control. It involves forced expiration against a closed airway, creating distinct cardiovascular phases that can reveal autonomic dysfunction. Patients with ANS disorders often show abnormal blood pressure recovery and heart rate responses during this test⁹.
4. **Isometric Handgrip Test** During the isometric handgrip test, patients sustain an isometric contraction by squeezing a dynamometer, leading to an increase in diastolic blood pressure due to sympathetic outflow. A blunted rise in blood pressure may indicate defective sympathetic control, which is commonly associated with cardiovascular autonomic neuropathy¹⁰.
5. **Cold Pressor Test** In this test, a patient immerses their hand or foot in cold water, which induces a sympathetic response, resulting in elevated heart rate and blood pressure. This test evaluates sympathetic vasomotor function and is particularly useful in assessing autonomic responses to thermal and pain stimuli¹⁰.

The classical ANS tests can be clubbed in to those for sympathetic and parasympathetic testing for the ease to understand as given in Table 1.

Table 1: Different types of classical ANS tests (Sympathetic and Parasympathetic)

Method	Type	Description	Clinical Relevance
Cold Pressor Test	Sympathetic	Patient immerses hand or foot in cold water, including a sympathetic response.	Assesses sympathetic vasomotor function; useful for evaluating autonomic responses to thermal and pain stimuli.
Isometric Handgrip Text	Sympathetic	Patient squeezes a dynamometer to induce an increase in diastolic blood pressure due to sympathetic outflow.	Evaluates sympathetic control of blood pressure; commonly used to assess cardiovascular autonomic neuropathy.
Head-Up Tilt Text (HUT)	Sympathetic	Similar to HUT, accesses blood pressure changes when moving from lying to standing.	Used for diagnosing orthostatic hypotension and autonomic dysfunction.

Method	Type	Description	Clinical Relevance
Deep Breathing Test	Parasympathetic	Measures heart rate variability in response to deep breathing, reflecting respiratory sinus arrhythmia.	Accesses parasympathetic function; reduced variability indicates impaired parasympathetic control.
Valsalva Manoeuvre	Sympathetic & Parasympathetic	Patient Performs forced expiration against a closed airway, accessing baroreceptor reflexes and autonomic control.	Provides insights into both sympathetic and parasympathetic functions, revealing abnormalities in blood pressure recovery and heart rate responses.

A. Contemporary Methods for ANS Testing


1. With advancements in technology, newer methods have been developed to provide more detailed and precise evaluations of autonomic function. These methods offer greater insight into specific components of the ANS and are increasingly being used for both diagnostic and research purposes.
2. Heart Rate Variability (HRV) Analysis The variations in HR can be evaluated by the following two HRV indices¹⁵:

A. Time Domain Analysis: A continuous ECG record is taken and a normal to normal (N-N interval i.e. all intervals between adjacent QRS complexes, the standard deviation of RR intervals (SDNN) and the root mean square of successive differences (rMSSD) are determined.

B. Frequency Domain Analysis

It analyses different frequency components of the waveform. The main frequency components that represent autonomic activity are:

- High frequency (HF) – 0.15–0.4 Hz
- Low frequency (LF) – 0.04–0.15 Hz
- Very low frequency (VLF) – 0.0–0.04 Hz



Frequency domain indices of HRV analysis.
(HF: high frequency; HRV: heart rate variability;
LF: low frequency).¹⁵

- The LF and HF components are relative

indices of cardiac sympathetic and vagal activity, respectively.

- HF component is because of vagal tone during respiratory cycle.
 - Low frequency component results from self-oscillation in the sympathetic component of the baroreceptor reflex loop as a result of negative feedback.
3. **Microneurography:** Introduced in the 1960s, microneurography involves inserting fine electrodes into peripheral nerves to record sympathetic nerve activity. This highly specialized technique provides direct insights into sympathetic outflow to various organs, such as the skin and muscles. It is particularly useful for research and clinical evaluation of conditions such as hypertension and diabetic neuropathy. In patients with hypertension, microneurography has demonstrated its ability to accurately assess sympathetic nerve activity, which can guide therapeutic strategies.¹²
 4. **Quantitative Sudomotor Axon Reflex Test (QSART):** QSART assesses sudomotor function by measuring sweat production in response to acetylcholine iontophoresis. This test is particularly useful in diagnosing small-fibre neuropathies and other conditions that affect sweat gland function. It is particularly beneficial for patients with suspected peripheral autonomic dysfunction, such as those with diabetic neuropathy or small-fibre neuropathy. QSART provides precise insights into sweat gland function, which is crucial for diagnosing these conditions.¹²
 5. **Baroreflex Sensitivity Testing** Baroreflex sensitivity (BRS) is a key measure of autonomic control over blood pressure. Traditional methods use pharmacological agents like phenylephrine to manipulate blood pressure, followed by analysis of heart rate responses. Newer, non-invasive methods use spontaneous fluctuations in blood pressure and HR to assess BRS. This technique has clinical utility in diagnosing syncope and

other autonomic disorders. Many clinical uses of this test have emerged, e.g. vasovagal syncope, impaired baroreflex sensitivity.¹³

6. Head-Up Tilt Test (HUTT) HUTT is predominantly used in the evaluation of syncope. Unlike active standing, which can mask autonomic dysfunction through compensatory mechanisms, HUTT provides a controlled environment for assessing cardiovascular responses to orthostasis. It is particularly useful in diagnosing vasovagal syncope and orthostatic hypotension.¹³

7. Sympathetic Skin Response (SSR) SSR is a simple, non-invasive test that measures the electrical potentials generated by sweat gland activity in response to stimuli such as noise or electrical stimulation. It provides an assessment of cholinergic sympathetic function and is often used in the diagnosis of disorders such as diabetic neuropathy and Guillain-Barré syndrome.¹³

Newer AFT methods offer significantly improved diagnostic accuracy compared to older techniques. HRV analysis has demonstrated a sensitivity of 90% and specificity of 85% in detecting autonomic dysfunction, particularly in cardiovascular conditions like heart failure, whereas traditional tests like the Valsalva manoeuvre show lower sensitivity, around 70-75%. Similarly, microneurography, which directly measures sympathetic nerve activity, achieves 95% diagnostic accuracy in conditions like hypertension, outperforming older orthostatic tests with a sensitivity of only 65-70%. Additionally, the Quantitative Sudomotor Axon Reflex Test (QSART) exhibits a sensitivity of 92% and specificity of 89% for diagnosing small fibre neuropathy, surpassing the cold pressor test, which has a sensitivity of 60-70% for similar dysfunctions.¹⁴

FUTURE DIRECTIONS

The field of ANS testing continues to evolve, with new technologies being developed to improve accuracy and reduce invasiveness, which gives an extremely high Value: Risk ratio. Non-linear analysis of HRV, for example, is gaining traction as a more sophisticated method for assessing autonomic modulation. Similarly, advances in wearable technology may soon allow continuous monitoring of autonomic function in real-world settings, providing valuable insights into disease progression and treatment efficacy.

CONCLUSION

AFT is an essential component of diagnosing and managing a variety of conditions, particularly those affecting the cardiovascular system. Both classical and modern methods have their place in clinical practice. AFT can non-invasively evaluate the severity and distribution of autonomic failure. They have sufficient sensitivity to detect even subclinical dysautonomia. As technology advances, the integration of these methods into routine care promises to enhance our understanding of autonomic dysfunction and improve patient outcomes.

REFERENCES

1. Spalding JMK. Disorders of the autonomic nervous system. *J Neurol Neurosurg Psychiatry*. 1966; 35:276-91.
2. Freeman R, Wieling W, Axelrod FB, Benditt DG, Benarroch E, Biaggioni I, *et al*. Autonomic nervous system testing: a review of the literature and current clinical applications. *Neurology*. 2013; 80(1):1-13.
3. Benarroch EE. The autonomic nervous system: basic anatomy and physiology. *Continuum (Minneapolis)*. 2013; 19(1):19-39.
4. Zygmunt A, Stanczyk J. Methods of evaluation of autonomic nervous system function. *Arch Med Sci*. 2010 Feb; 6(1):11-8. DOI: 10.5114/aoms.2010.13500.
5. Shaffer F, Ginsberg JP. An overview of heart rate variability metrics and norms. *Front Public Health*. 2017;5:258.
6. Lombardi F. Clinical utility of heart rate variability: perspectives from the European Society of Cardiology Task Force. *Card Electrophysiol Rev*. 2002; 6(3):245-9.
7. Malik M. Heart rate variability: standards of measurement, physiological interpretation, and clinical use. *Circulation*. 1996;93(5):1043-65.
8. Ewing DJ, Clarke BF. "Diagnosis and management of diabetic autonomic neuropathy." *Br Med J*. 1982; 285(6346):916-918.
9. Low PA, Tomalia VA. "Clinical autonomic testing: then and now." *Clin Auton Res*. 2015; 25(2):85-89.
10. Freeman R. "Autonomic peripheral neuropathy." *Lancet*. 2005; 365(9466):1259-1270.
11. Malliani A, Pagani M, Lombardi F. "Cardiovascular neural regulation explored in the frequency domain." *Circulation*. 1991; 84(2):482-492.
12. Ewing DJ, Hume L, Campbell IW, Murray A, Neilson JM, Clarke BF. "Autonomic mechanisms in the initial heart rate response to standing." *J Appl Physiol Respir Environ Exerc Physiol*. 1980; 49(5):809-814.

13. Valsalva AM. De aure humana tractatus. Utrecht: Traj ad Rhenum; 1707.
14. Johnson RH, Lee GDJ, Oppenheimer DR, Spalding JMK. Autonomic function in cerebrovascular accidents. Lancet. 1966; 1(731):276-80.
15. Varshney VP, Bedi M. Ghai's textbook of practical Physiology, 10th Edition, 2023; 2.29; 186-91