

# Role of Hybrid Reconstructive Ladder in Adult Scald Burns

Lakshmi Sriram<sup>1</sup>, Neljo Thomas<sup>2</sup>, Ravi Kumar Chittoria<sup>3</sup>

## How to cite this article:

Lakshmi Sriram, Neljo Thomas, Ravi Kumar Chittoria/Role of Hybrid Reconstructive Ladder in Adult Scald Burns/Indian Journal of Medical & Health Sciences. 2022;9(2):47-50.

## Abstract

The reconstructive ladder discusses progressively more difficult soft tissue coverage methods for wound coverage and healing. Regenerative medicine modalities can be added to conventional reconstructive methods, making it a hybrid reconstruction ladder, which may make it easier to achieve wound coverage at a lower 'rung'. We have used the hybrid reconstruction ladder in an adult patient with scald burns with significant clinical improvement.

**Keywords:** Hybrid reconstructive ladder; Scald burns; Adult.

## INTRODUCTION

A burn is defined as a thermally induced traumatic injury to organic tissue. The most often reported thermal injuries in children under the age of 16 were scalds and contact burns.<sup>1</sup>

Burn injuries can have a significant impact on physical and psychological health, as well as health related morbidity and mortality. Burn victims are more likely to have depression and other linked psychiatric issues, and severe burns can

leave patients with substantial scars and painful contractures.<sup>2</sup> The term "reconstructive ladder" was created by plastic and reconstructive surgeons to describe the several levels of increasingly complex treatment for soft tissue disorders. At the bottom of the ladder, the surgeon would handle a clinical reconstructive problem by applying the most straight forward reconstruction approach.

As a more complex or effective solution was needed for a specific reconstruction challenge, the reconstructive surgeon would move up the ladder. Complex wound patterns have sparked efforts to create novel tissue regeneration techniques. These procedures, also referred to as hybrid reconstructions, combine regenerative medicine applications with traditional reconstruction methods. By limiting the disability and morbidity brought on by standard reconstruction, the hybrid reconstruction model helps to maximise function.<sup>5</sup>

## MATERIALS AND METHODS

This study was conducted in the Department of

**Author Affiliation:** <sup>1</sup>Junior Resident, Department of General Surgery, <sup>2</sup>Senior Resident, Department of Plastic Surgery, <sup>3</sup>Professor & Head of IT Wing and Telemedicine, Department of Plastic Surgery & Telemedicine, Jawaharlal Institute of Postgraduate Medical Education and Research, Pondicherry 605006, India.

**Corresponding Author:** Ravi Kumar Chittoria, Professor & Head of IT Wing and Telemedicine, Department of Plastic Surgery & Telemedicine, Jawaharlal Institute of Postgraduate Medical Education and Research, Pondicherry 605006, India.

E-mail: [drchittoria@yahoo.com](mailto:drchittoria@yahoo.com)

Received on: 30.08.2022

Accepted on: 12.09.2022

Plastic Surgery, in a tertiary care centre in South India after getting the department ethical committee approval. Written and informed consent was taken from the patient for proposed treatment plan and clinical photography. The subject was a thirty three

old female with alleged history of accidental scald burns over both thighs and abdomen. Hydro jet debridement, Vitamin D3 therapy, Autologous platelet rich plasma, Dry collagen scaffold dressing, Low level laser therapy (Fig. 1-6),



Fig. 1: Patient at presentation



Fig. 2: Hydrojet debridement



Fig. 3: Vitamin D3 with Cantella extract



Fig. 4: Activated platelet rich plasma therapy



Fig. 5: Low level laser therapy



Fig. 6: Heterografting with dry collagen sheet

Negative pressure wound therapy (Fig. 7), were used in the preparation of wound bed followed by split thickness skin grafting (Fig. 8).



Fig. 7: Negative pressure wound therapy

## RESULTS

The wound healed well at the time of discharge. There was good graft take. There were no side effects noted.

## DISCUSSION

Burn injuries, particularly severe burns, are accompanied by an immune and inflammatory response, metabolic changes and distributive shock that can lead to multiple organ failure. Scald injuries tend to appear more superficial initially, due to rapid dilution of the source and energy.<sup>3</sup>

Skin grafting, bone reconstruction, nerve repair or reconstruction, and vascular repair are some of the standard treatment modalities for injuries with significant composite tissue loss.<sup>5,6</sup> These traditional treatments could reduce a patient's functional ability and increase donor site morbidity.<sup>7</sup> This has led to increased use of regenerative medicine techniques to promote tissue regeneration and improve reconstructive and clinical outcomes.

Autologous platelet rich plasma is a treatment that contains fibrin and high concentrations of growth factors with the potential to improve the healing of chronic wounds. But the overall quality of evidence of autologous PRP for treating chronic wounds is low.<sup>8</sup>

Low level lasers that affect biological systems without using heat include those made of Krypton, Argon, He, Ne, and ruby. When the tissue chromophores are influenced by laser energy, the cytochromes in the mitochondria absorb the



Fig. 8: Split thicknes skin grafting



Fig. 9: Patient at discharge

laser radiation and convert them into energy by the cell (ATP), and created energy induces protein synthesis and acceleration or stimulation of cell proliferation.<sup>11</sup> The interaction of light with biological tissues is influenced by various factors, including wavelength, laser dose, and the tissue's optical characteristics. The structure, water content, thermal conductivity, heat capacity, density, and capacity to absorb, disperse, or reflect the released energy are examples of tissue qualities.<sup>9</sup>

Negative pressure wound therapy, also called vacuum assisted wound closure, refers to wound dressing systems that continuously or intermittently apply sub atmospheric pressure to the system, which provides a positive pressure to the surface of a wound. NPWT has become a popular treatment

modality for the management of many acute and chronic wounds.<sup>10</sup>

In our study too, regenerative techniques proved to be a great adjunct to standard reconstruction procedures in wound healing and overall clinical outcome.

## REFERENCES

1. Brassolatti, P., de Andrade, A.L.M., Bossini, P.S. et al. Evaluation of the low-level laser therapy application parameters for skin burn treatment in experimental model: a systematic review. *Lasers Med Sci* 33, 1159–1169 (2018).
2. Zanni GR. Thermal burns and scalds: clinical complications in the elderly. *Consult Pharm.* 2012 Jan;27(1):16-22.
3. Belmonte Torras JA, Marín de la Cruz D, Gornés Benajam MB, Gubern Pi L, Guinot Madrdejos A. Quemaduras por agua caliente sanitaria [Tap-water scald burns]. *An Pediatr (Barc)*. 2004 Nov;61(5):413-7. Spanish.
4. Latifi R, El-Hennawy H, El-Menyar A, et al. The therapeutic challenges of degloving soft-tissue injuries. *J Emerg Trauma Shock*. 2014;7:228–32.
5. Gottlieb LJ, Krieger LM. From the Reconstructive Ladder to the Reconstructive Elevator. *Plastic and Reconstructive Surgery*. 1994;93(7):1503.
6. Vedder NB, Wei FC, Mardini S, eds. Problem analysis in reconstructive surgery.
7. Reconstructive ladders, elevators, and surgical judgment. In: *Flaps and Reconstructive Surgery*. 2017;2nd ed. Toronto: Elsevier; 1–5.
8. Stansbury LG, Lalliss SJ, Branstetter JG, Bagg MR, Holcomb JB. Amputations in U. S. military personnel in the current conflicts in Afghanistan and Iraq. *J. Orthop. Trauma* 22(1), 43–46 (2008).
9. Mohapatra, Devi Prasad M.Ch.; Thiruvoth, Friji Meethale M.Ch. Reconstruction 2.0: Restructuring the Reconstructive Ladder, *Plastic and Reconstructive Surgery: March 2021 -Volume 147 - Issue 3 - p 572e-573e* Martínez-Zapata MJ, Martí-Carvajal AJ, Solà I, Expósito JA, Bolibar I, Rodríguez L, Garcia J, Zaror C. Autologous platelet-rich plasma for treating chronic wounds. *Cochrane Database Syst Rev*. 2016 May 25;2016(5):CD006899.
10. Saurabgupta et al. Effect of Low Level Laser Therapy (LLLT) On The Severity of Post-Burn Immature Scars: A Randomized Control Study. *International Journal of Clinical And Diagnostic Research* Volume 8, Issue 1, Jan-Feb 2020.
11. Hopkins JT, McLoda TA, Seegmiller JG, David Baxter G. Low-Level Laser Therapy Facilitates superficial wound healing in humans: a triple-blind, sham-controlled study. *J Athl Train*. 2004;39:223–229.