

Role of Application of Topical Nitroglycerine in Management of Flap Surgery

Nishad K.¹, Neljo Thomas², Ravi Kumar Chittoria³, Barath Kumar Singh⁴,
Jacob Antony Chakiath⁵

How to cite this article:

Nishad K., Neljo Thomas, Ravi Kumar Chittoria, *et al.*/Role of Application of Topical Nitroglycerine in Management of Flap Surgery/Indian J Prev Med. 2023;11(1):23-25.

Abstract

Flaps have been used for a long time for closure of soft tissue defects by plastic surgeons and form an important part of the surgeon's armoury, however 2% –22% of flap procedures led to necrosis in an observational study.¹ The search for an ideal intervention for ensuring flap survival still continues. Topical nitroglycerine has been found effective in prevention of flap necrosis in few studies.²⁻⁴ In this article we share our experience with prevention of flap necrosis using topical nitroglycerine.

Keywords: Topical nitroglycerine; Flap necrosis.

INTRODUCTION

Flap surgery has been an important part of the plastic surgery practice but an important complication is flap necrosis. Interventions have been used in various forms to prevent the flap necrosis by improving the blood supply to the area by producing vasodilatation. The ideal method is to do a delayed flap surgery so that the choke vessels open up. However, most often patients don't prefer a second surgery. Nitroglycerine is a vasodilator and acts by releasing nitric oxide and thus relaxes the vascular smooth muscle.⁵ Another

known advantage is its antithrombotic capabilities. By increasing the calibre of the vasculature and ensuring the patency of the vasculature it increases blood flow to the area of application.²

METHODOLOGY

This study was conducted in a tertiary care centre in the department of plastic surgery during the period of April to July 2020. There were two patients who underwent flap cover and we used topical nitroglycerine for prevention of flap necrosis. The cost of topical preparation of nitroglycerine was 34 rupees for 30gm.

Case 1

50 year female with h/o trauma and suffered injury to left distal forearm with loss of soft tissue over the lateral aspect, underwent hypogastric flap cover. (Fig. 1), topical NTG (Fig. 2) was applied at a low dose (4.5 mg) with repeated application (3 times daily) for 3 days 4 and continued till 7th post-operative day.

Author Affiliation: ^{1,2,4,5}Senior Resident, ³Professor, Department of Plastic Surgery, Jawaharlal Institute of Postgraduate Medical Education and Research, Pondicherry 605006, India.

Corresponding Author: Ravi Kumar Chittoria, Professor, Department of Plastic Surgery, Jawaharlal Institute of Postgraduate Medical Education and Research, Pondicherry 605006, India.

E-mail: drchittoria@yahoo.com

Received on: 13.12.2022

Accepted on: 15.01.2023



Fig. 1: Hypogastric flap



Fig. 2: Nitroglycerine ointment for local application

RESULT

There was no areas of flap necrosis seen at the end of flap division (Fig. 3a, b).



Fig. 3a: Flap after division



Fig. 3b: Flap after division

DISCUSSION

Successful use of topical NTG application to prevent flap necrosis was first reported in an animal model by Rohrich in 1984⁶ and then in the human subjects by Schauer in 1986.⁷ Nitroglycerine, or glyceryl nitrate, is an organic nitrate.⁶ It is a direct vasodilator when applied topically, with solution, ointment, and transdermal patch available. It releases nitric oxide (NO) and activates intracellular cyclic guanosine monophosphate (cGMP) and inhibition of myosin light chain kinase, and therefore causes relaxation of vascular smooth muscle.⁵ The increase in calibre of arteries and veins increases blood flow to the area of application² and has various results in reducing flap necrosis rates in animal models.^{3,8,9}

It causes improvement in flap survival by releasing prostacyclin, which decreases thrombosis in cultured human endothelial cells.⁶ Neovascularization was found to be the highest on day 7 after surgery in the rat model by Aral *et al.*⁸ Therefore, repeated NTG application daily may be needed for at least 7 days and may also be needed due to its short half-life, which is less than a day.¹⁰ It is detected 30 min after adequate plasma levels are attained and sustained for 4–6 h, with the therapeutic range in humans being greater than 1mg/ml⁶. Repeated application can be used as an alternative to an occlusive dressing.¹⁰

The possible complications that can arise are of hypotension, headache, and dizziness. However, in our study no patients developed any side effects. The advantage of topical nitroglycerine is that it is easy to apply and use.

CONCLUSION

In our study we have come to the conclusion that topical nitroglycerine is useful in prevention of flap necrosis, however ours is a small study and needs large scale RCT to bring the intervention to practice.

DECLARATIONS

Acknowledgment

Authors' contributions: All authors made contributions to the article

Availability of data and materials: Not applicable.

Financial support and sponsorship: None

Conflicts of interest: None.

Consent for publication: Not applicable.

Copyright: © The Author(s) 2020.

REFERENCES

1. Matsen CB, Mehrara B, Eaton A, *et al.* Skin flap necrosis after mastectomy with reconstruction: a prospective study. *Ann Surg Oncol* 2016; 23 (1):257–64.
2. Gdalevitch P, Van Laeken N, Bahng S, *et al.* Effects of nitroglycerin ointment on mastectomy flap necrosis in immediate breast reconstruction: a randomized controlled trial. *Plast Reconstr Surg* 2015; 135 (6):1530–9.
3. Rohrich RJ, Cherry GW, Spira M. Enhancement of skin flap survival using nitroglycerin ointment. *Plast Reconstr Surg* 1984; 73 (6):943–8.
4. Yun MH, Yoon ES, Lee BIL, Park SH. The effect of low-dose nitroglycerin ointment on skin flap necrosis in breast reconstruction after skin-sparing or nipple-sparing mastectomy. *Arch Plast Surg* 2017; 44 (6):509–15. doi: 10.5999/aps.2017.00934.
5. Vargas CR, Iorio ML, Lee BT. A systematic review of topical vasodilators for the treatment of intraoperative vasospasm in reconstructive microsurgery. *Plast Reconstr Surg* 2015; 136 (2):411–22
6. Levin RI, Jaffe EA, Weksler BB, Tack-Goldman K. Nitroglycerin stimulates synthesis of prostacyclin by cultured human endothelial cells. *J Clin Invest* 1981; 67 (3):762–9.
7. Scheuer S, Hanna MK. Effect of nitroglycerin ointment on penile skin flap survival in hypospadias repair experimental and clinical studies. *Urology* 1986; 27 (5):438–40.
8. Aral M, Tuncer S, Şencan A, Elmas Ç, Ayhan S. The effect of thrombolytic, anticoagulant, and vasodilator agents on the survival of random pattern skin flap. *J Reconstr Microsurg* 2015; 31 (7):487–92.
9. De SáJZ, de Aguiar JLA, Cruz AF, Schuler ARP, de Lima JRA, Marques OM. Efeitos da nitroglicerina tópica na autoneurização de retalhos pré-fabricados por implante vascular em ratos. *Acta Cir Bras* 2012; 27 (12):905–11
10. Dünn GL, Brodland DG, Griego RD, Huether MJ, Fazio MJ, Zitelli JA. A single postoperative application of nitroglycerin ointment does not increase survival of cutaneous flaps and grafts. *Dermatologic Surg* 2000; 26 (5):425–7.

