

Total Parenteral Nutrition

Barath Kumar Singh P¹, Ravi Kumar Chittoria²

How to cite this article:

Barath Kumar Singh. P, Ravi Kumar Chittoria/Total Parenteral Nutrition/Indian J Trauma Emerg Pediatr.2022;14(4):123-129.

Abstract

Total parenteral nutrition is a medication used to manage and treat malnourishment. It is in the nutrition class of drugs. Parenteral nutrition (TPN) is when the intravenous administered nutrition can serve as the main source of nutrition and also as a supplementary nutrition if the patient is not having adequate nutrition. This article highlights the composition, initiation, indications, contraindications, complications of total parenteral nutrition as a valuable agent in managing malnourishment.

Keywords: Total Parenteral Nutrition; Initiation; Composition; Indication; Monitoring; Complication.

INTRODUCTION

Parenteral nutrition support refers to the provision of calories, amino acids, electrolytes, vitamins, minerals, trace elements, and fluids via a parenteral route. Access, prescribing, monitoring, and complications of parenteral nutrition are discussed. Parenteral nutrition is a mixture of solutions that contain dextrose, amino acids, electrolytes, vitamins, minerals and trace elements.^{1,2} The initiation, composition, indications, contraindications, monitoring, complications of parenteral nutrition are discussed in this review article.

Author Affiliation: ¹Senior Resident, Department of Plastic Surgery, ²Professor & Registrar, Department of Plastic Surgery & Telemedicine, Jawaharlal Institute of Postgraduate Medical Education and Research, Pondicherry 605006, India.

Corresponding Author: Ravi Kumar Chittoria, Professor & Registrar, Department of Plastic Surgery & Telemedicine, Jawaharlal Institute of Postgraduate Medical Education and Research, Pondicherry 605006, India.

E-mail: drchittoria@yahoo.com

Received on: 07.11.2022

Accepted on: 12.12.2022

MATERIALS AND METHODS

This study was conducted in the Department of Plastic Surgery in a tertiary care institute. It is a review article based on the study of literature available in Scopus, PubMed, Google scholar & open sources. Inclusion criteria for selecting articles was to include only articles published at least in peer reviewed indexed journals about Parenteral Nutrition.

RESULTS

Based on the inclusion criteria 38 articles were studied to discuss Total Parenteral nutrition under following headings:

1. Initiation of Parenteral Nutrition
2. Composition of Parenteral Nutrition
3. Indications and Contraindications
4. Vitamins and Trace elements
5. Monitoring
6. Complications

DISCUSSION

Initiation of Parenteral Nutrition

In order to initiate parenteral nutrition, appropriate access must be obtained and the prescription (i.e., composition and infusion rate) must be determined. Guidelines suggest that when tolerance to enteral nutrition is evident, parenteral nutrition should be weaned and discontinued when >60 percent of the patient needs are met enterally.¹

Our practice is to wean parenteral nutrition proportionate to the amount of enteral nutrition being delivered, thus meeting but not exceeding the daily goals. For ease of calculation, most standard Parenteral Nutrition solutions contain approximately 1 kcal/ml.

Venous Access

Parenteral nutrition given for more than a few days must be delivered via a central venous catheter, so called central parenteral nutrition (CPN) because its high osmotic load is not tolerated by peripheral veins.^{2,3} Parenteral nutrition may be given via a peripheral vein if it is significantly more dilute (<900 mOsm), so called peripheral parenteral nutrition (PPN). Because of the ease of obtaining central access in the hospital, the large volume usually required to administer PPN, and the lack of benefit from short term parenteral nutrition, PPN is rarely indicated.

Choosing among a peripherally inserted central catheter (PICC) or a tunnelled central venous catheter (TCVC); also referred to as TICC, eg, Hickman catheter, Groshong catheter, or implanted infusion port) should be individualized and depend upon duration of need for Parenteral Nutrition, local expertise, ability of the facility or individual to take care of the device, and presence of other risk factors for catheter related blood stream infections (CRBSI)

The American Society for Parenteral and Enteral Nutrition (ASPEN) has issued clinical guidelines to describe best practices in the selection and care of central venous access devices (CVADs) for the infusion of home parenteral nutrition in adult patients.⁴ ASPEN guidelines use an arbitrary cutoff of 30 days to distinguish short term from long term Parenteral Nutrition use.

- The patients in whom short term administration of PN is desirable or intended that PN be delivered through a PICC. Alternatively, while not preferable, PN may be administered through a subclavian,

internal jugular, or femoral central venous catheter if PN is only needed for very short periods (eg, a few days) or if a PICC is not feasible or reasonable. Tradition teaches that the femoral site is least desirable due to an increased risk of CRBSIs⁵, and is usually not preferred.

- For long term administration of PN, A.S.P.E.N. guidelines indicated a preference for atunnelled central venous catheter (TCVC). The rationale for this choice is based upon expert opinion and observational studies that suggest infection rates may be lower with TCVC.⁶⁻⁹ In fact, one systematic review suggested that rates were no different between PN delivered through a PICC compared with a TCVC.^{6,7}
- When possible, a single lumen central venous catheter should be used for the infusion of parenteral nutrition. If a multiple lumen central venous catheter is used it should have one port dedicated solely for the infusion of parenteral nutrition. In addition, catheter manipulations should be minimized. These precautions may decrease the infectious complications associated with parenteral nutrition.^{10,11} For patients who have an existing CVC, a new CVC is not typically required unless there has been septicaemia during the life of the existing line.
- The Institute for Safe Medication Practices (ISMP) has listed Parenteral nutrition as a "high alert medication," in the same category of complexity and potential for error or harm as chemotherapy. Guidelines that describe safe practices for parenteral nutrition have been published by professional organizations.¹¹⁻¹⁵ These recommend that parenteral nutrition be prescribed by a multidisciplinary team of physicians, nutritionists, pharmacists, and nurses.

Parenteral nutrition should not be prescribed by clinicians nor compounded by pharmacists without specific training because it is complicated and requires advanced knowledge about issues such as nutrient metabolism and solute compatibility.^{14,15} Composition of Parenteral nutrition

Parenteral nutrition is an admixture of solutions containing dextrose, amino acids, electrolytes, vitamins, minerals, and trace elements. Lipid emulsion may be infused separately or added to the mixture. However, mixing all three, a so called total nutrient admixture, or 3-in-1 parenteral nutrition, is favoured by most experts.¹⁶ The exact composition

and infusion rate are tailored to the nutritional and fluid needs of each patient.

Dextrose

Dextrose-containing stock solutions are available in a variety of concentrations, most commonly 40, 50, and 70 percent. The percentage of calories that is contributed by dextrose is titrated according to individual factors, such as severity of illness, the caloric needs of the patient, and the patient's ability to tolerate fluid volume. The caloric contribution of dextrose in medical solutions is 3.4 kcal/gm, which differs from dietary carbohydrate (4 kcal/gm). The reason for the difference is that water contributes to the weight of the dextrose hydrate that is used to prepare parenteral nutrition.

Amino Acids and Electrolytes

- Amino acid solutions contain most essential and nonessential amino acids. The exceptions are arginine and glutamine. While these are thought to be conditionally essential in critical illness, due to blockade of their metabolic pathways in catabolism, research does not support their supplementation. The caloric contribution of amino acids is approximately 4 kcal/gm. The buffer used in amino acid solution contains electrolytes, usually in small quantities.
- Amino acid solutions with large amounts of electrolytes have been used infrequently. Premixed parenteral nutrition with a variety of available electrolyte content are available. It is occasionally necessary to use amino acid solutions with further electrolyte restriction (usually these are 15 percent amino acid solutions), particularly in patients with renal failure when electrolytes, particularly phosphate levels, are difficult to control.
- The amino acid stock solutions come in concentrations ranging from 5.5 to 15 percent. Higher concentrations are useful for minimizing volume and electrolytes delivered to patients.
- Branched chain amino acid-enriched parenteral nutrition to standard parenteral nutrition found a trend toward decreased mortality among patients who received branched chain amino acids.¹⁷

Lipids

Lipids are provided as an emulsion that may be infused separately or added to the mixture [total nutrient admixture (TNA) or three-in-one]. Most

lipid emulsion consists of long-chain omega-6 triglycerides derived from soybean and safflower oils and then emulsified using egg phospholipids and glycerine. Mixtures of several types of lipids like refined olive, soybean, and fish oil emulsions have been approved and are also available for use.¹⁸⁻²²

- Omega-6 fatty acids, omega-3 fatty acids are metabolized to mediators that are noninflammatory and anti-inflammatory. Manipulation of the inflammatory process has been of strong interest in the nutrition support, as it is the cause of disease related wasting. omega-3 fatty acid enriched parenteral nutrition was associated with a 40 percent lower risk of infection and 56 percent lower risk of sepsis compared with patients receiving standard parenteral nutrition without omega-3 fatty acid enrichment.²³ Hospital and ICU length of stay was also lower in the fatty acid enriched group.
- The caloric contribution of a typical lipid emulsion is 10 kcal/g or 2 kcal/mL in 20 percent emulsion and 11 kcal/g or 1.1 kcal/mL in 10 percent emulsion. Dietary fat provides 9 kcal/g, but there is a contribution of calories from the emulsifiers used to create the lipid suspension for intravenous administration. Intravenous fat emulsions should be given with care to patients with prior allergy to eggs. Egg phospholipid is used to emulsify the triglycerides in intravenous fat emulsions and allergic cross-reactions have been rarely reported.
- Cautious initiation of Intravenous lipid emulsions (ILE) has been recommended when serum triglycerides exceed 200mg/dL (eg, infuse lipid emulsion three times weekly). Severe hypertriglyceridemia and other severe disorders of lipid metabolism are a contraindication to lipid emulsion infusion (eg, serum triglycerides >1000 mg/dL due to pathologic hypertriglyceridemia, lipid necrosis, acute pancreatitis accompanied by hypertriglyceridemia, disorders of fat metabolism).

Peripheral Parenteral Nutrition

Peripheral parenteral nutrition (PPN) is prescribed when short-term parenteral nutrition is required. It is a type of parenteral nutrition that can be delivered for short periods through a peripheral intravenous catheter because it has an osmolality lower than that of conventional parenteral nutrition (<900 mOsm). To deliver adequate nutrients, either

a large volume and/or a high fat formulation must be used. Frequent replacement of intravenous access is usually necessary. Despite an osmolarity lower than conventional parenteral nutrition, PPN is still quite hyperosmolar and irritating to the peripheral veins causing thrombosis and stenosis of the veins.

Indications and Contraindications

The indications and contraindications for parenteral nutrition are described.

Indications

- Chronic intestinal obstruction as in intestinal cancer
- Bowel pseudo-obstruction with food intolerance.
- TPN can also be used to rest the bowel in cases of GI fistulas with high flow
- When an infant's gastrointestinal system is immature or has a congenital gastrointestinal malformation
- When there is a post-operative bowel anastomosis leak
- When the patient is unable to maintain nutritional status due to severe diarrhoea or vomiting
- Small bowel obstruction
- Hypercatabolic states due to sepsis, polytrauma, and major fractures
- An anticipated period of nothing by mouth (NPO) status greater than seven days as in patients with inflammatory bowel disease exacerbations as well as critically ill patients

Contraindication

- Infants with less than 8 cm of the small bowel
- Irreversibly decerebrate patients
- Patients with critical cardiovascular instability or metabolic instabilities; such instabilities require correction before administering intravenous nutrition.
- When gastrointestinal feeding is possible
- When the nutritional status is good, only short-term TPN is needed
- The lack of a therapeutic goal, as TPN should not be used to prolong life when death is unescapable.

Vitamins and trace elements

Patients receiving parenteral nutrition must receive adequate vitamins and minerals to prevent deficiencies. For most patients, a unit dose of a standard multivitamin and multi-trace element solution will suffice to provide minimum daily requirements. There are also concerns about clearance of some vitamins in patients. Monitor levels of vitamins and minerals if patients are on parenteral nutrition for extended periods.²⁴

Therapeutic Vitamin and Mineral Supplementation

It is not our practice to supplement vitamins and minerals above the recommended daily amounts, unless a deficiency is documented or strongly suspected. Rather, a deficiency occurs only when a lack of a substance causes an undesired health outcome that is reversed or prevented by supplementation.

- Levels of vitamin D, which is protein bound, drop in critical illness. But even though there is a correlation between vitamin D levels and ICU mortality, supplementation of vitamin D has no effect.^{25,26} The decrease in vitamin D levels is due to the decrease in carrier protein levels, much in the same way a decrease in albumin lowers calcium levels.
- Antioxidants on critically ill patients when given as single nutrients (eg, selenium) or as a combination of nutrients (selenium, copper, zinc, vitamin A, vitamin C, vitamin E, and N-acetylcysteine) is beneficial. The trials administered the antioxidants in various ways, including as a separate intravenous infusion, as a component of parenteral nutrition, as a component of enteral nutrition, and orally. Critically ill patients who received vitamins and trace elements, either alone or in combination, had a lower mortality rate than patients who did not receive vitamins or trace elements.²⁷
- Given their safety, it seems reasonable to provide standard doses of vitamins and trace elements to most critically ill patients, regardless of the type of nutrition support that they are receiving. The optimal mixture of vitamins and trace elements is yet to be determined.

Glutamine

Glutamine is a precursor for nucleotide synthesis and an important fuel source for rapidly dividing cells, such as gastrointestinal epithelia. Despite

evidence that indicates that parenteral glutamine may be beneficial to patients who are receiving parenteral nutrition,²⁸⁻³⁰ Glutamine supplementation in critically ill patients associated with increased 28-day mortality in those receiving glutamine enriched enteral nutrition was reported.³¹

Monitoring

Routine monitoring of parenteral nutrition includes measurement of fluid intake and output, as well as selected laboratory studies. It is reasonable to measure serum electrolytes, glucose, calcium, magnesium, and phosphate daily until they are stable, or more frequently when the patient is at high risk for or exhibiting refeeding syndrome.³²

- Intake and output 12-hour charts
- Urine sugar monitoring every 8 hours
- Serum electrolytes: daily sodium, potassium, bicarbonate, calcium, and chloride values
- Serum creatinine and blood urea daily values
- Serum protein levels twice daily
- Liver function tests twice daily

It is similarly reasonable to measure amino transferases, bilirubin, and triglyceride at least once each week during treatment. More frequent measurements should be performed in the period immediately after parenteral nutrition is initiated, or after changes in the composition. Triglyceride levels should be measured at baseline and once PN has been initiated, then monthly or more frequently if elevated. We measure trace elements, including iron, approximately every three months, or if deficiency is suspected. Frequency of monitoring should be adjusted to the individual patient's acuity, stability, and risks for deficiencies.

Complications

Patients who receive parenteral nutrition support are at risk for infection, adverse metabolic complication, and complications related to venous access.

Venous Access

Parenteral nutrition requires venous access, which is associated with potential complications include bleeding, vascular injury, pneumothorax, venous thrombosis, arrhythmia, and air embolism.

Bloodstream Infection

Access to the central venous system is necessary when Parenteral Nutrition is given. Parenteral Nutrition is typically administered through a

peripherally inserted centralvenous catheter (PICC) or a tunnelled central venous catheter (TCVC). Patients receiving parenteral nutrition are at slightly increased risk of acquiring a bloodstream infection (bacterial and fungal); approximately one episode per 100 inpatient Parenteral Nutrition days.^{32,33}

Among patients receiving parenteral nutrition, factors that are independently associated with bloodstream infection include poor patient hygiene, insertion of the central venous catheter under emergent circumstances, and to a lesser extent, the severity of illness and duration of central venous catheterization.³⁴ Conversely, proper hand hygiene and maximal barrier precautions during insertion of the central venous catheter are associated with fewer bloodstream infections.³⁴

Metabolic Effects

Parenteral nutrition is associated with metabolic complications, including hyperglycemia, serum electrolyte alterations, macro or micro-nutrient excess or deficiency, refeeding syndrome,³⁵ Wernicke's encephalopathy,³⁶ and hepatic dysfunction.

Routine monitoring of serum glucose, electrolytes, and volume status may minimize the impact of such complications.

- Hyperglycaemia is particularly common among patients who receive parenteral nutrition. The incidence of hyperglycaemia was approximately two times greater for patients who received parenteral nutrition than patients who received enteral nutrition.³⁷
- Refeeding syndrome is a potentially fatal condition resulting from rapid changes in fluids and electrolytes when malnourished patients are given oral, enteral, or parenteral feedings.³⁸ Patients with ongoing electrolyte losses like diarrhoea, vomiting, fistulas are at increased risk of refeeding syndrome. It is defined primarily by manifestations of severe hypophosphatemia including respiratory failure, cardiovascular collapse, rhabdomyolysis, seizures, and delirium, hypokalaemia, hypomagnesemia, and Wernicke's encephalopathy also occur.
- Emergent hypertriglyceridemia or worsening of existing hypertriglyceridemia may result from the infusion of lipid emulsion. Concentrations up to 400 mg/dL are acceptable during therapy without altering the rate of lipids during therapy.

Doses of lipid emulsion should be reduced when triglyceride concentrations rise above 400 mg/dL. Levels >1000 mg/dL are a contraindication to infusion. It is also reasonable to attempt decreasing the proportion of calories provided by dextrose in this hypertriglyceridemia.

CONCLUSION

Parenteral nutrition should be prescribed by a multidisciplinary team with advanced knowledge about how to tailor the composition and infusion rate to the needs of each patient, rather than by untrained individual clinicians. Guidelines exist that describe safe practices for parenteral nutrition. For critically ill patients receiving parenteral nutrition, we suggest the inclusion of multivitamins and trace elements. They are typically already included as a component of the parenteral nutrition, in the multivitamin, and multi-trace element solutions that are usually given in unit doses. The optimal mixture of nutrients has not been determined. Routine monitoring of parenteral nutrition includes measurement of fluid intake and output, as well as selected laboratory studies. Patients who receive parenteral nutrition support are at risk for infection, adverse metabolic effects, and complications related to venous access.

REFERENCES

1. Compher C, Bingham AL, McCall M, et al. Guidelines for the provision of nutrition support therapy in the adult critically ill patient: The American Society for Parenteral and Enteral Nutrition. *JPEN J Parenter Enteral Nutr* 2022; 46:12.
2. Kuwahara T, Asanami S, Tamura T, Kaneda S. Effects of pH and osmolality on phlebotic potential of infusion solutions for peripheral parenteral nutrition. *J Toxicol Sci* 1998;23:77.
3. Kuwahara T, Asanami S, Tamura T, Kubo S. Dilution is effective in reducing infusion phlebitis in peripheral parenteral nutrition: an experimental study in rabbits. *Nutrition* 1998; 14:186.
4. Kovacevich DS, Corrigan M, Ross VM, et al. American Society for Parenteral and Enteral Nutrition Guidelines for the Selection and Care of Central Venous Access Devices for Adult Home Parenteral Nutrition Administration. *JPEN J Parenter Enteral Nutr* 2019;43:15.
5. Safdar N, Kluger DM, Maki DG. A review of risk factors for catheter related bloodstream infection caused by percutaneously inserted, noncuffed central venous catheters: implications for preventive strategies. *Medicine (Baltimore)* 2002; 81:466.
6. Hon K, Bihari S, Holt A, et al. Rate of Catheter-Related Bloodstream Infections Between Tunneled Central Venous Catheters Versus Peripherally Inserted Central Catheters in Adult Home Parenteral Nutrition: A Meta-analysis. *JPEN J Parenter Enteral Nutr* 2019;43:41.
7. Cotogni P, Pittiruti M, Barbero C, et al. Catheter-related complications in cancer patients on home parenteral nutrition: a prospective study of over 51,000 catheter days. *JPEN J Parenter Enteral Nutr* 2013; 37:375.
8. Buchman AL, Opilla M, Kwasny M, et al. Risk factors for the development of catheter-related bloodstream infections in patients receiving home parenteral nutrition. *JPEN J Parenter Enteral Nutr* 2014; 38:744.
9. Bech LF, Drustrup L, Nygaard L, et al. Environmental Risk Factors for Developing Catheter-Related Bloodstream Infection in Home Parenteral Nutrition Patients: A 6-Year Follow-up Study. *JPEN J Parenter Enteral Nutr* 2016; 40:989.
10. O'Grady NP, Alexander M, Dellinger EP, et al. Guidelines for the prevention of intravascular catheter-related infections. Centers for Disease Control and Prevention. *MMWR Recomm Rep* 2002; 51:1.
11. Mirtallo J, Canada T, Johnson D, et al. Safe practices for parenteral nutrition. *JPEN J Parenter Enteral Nutr* 2004; 28:S39.
12. Naylor CJ, Griffiths RD, Fernandez RS. Does a multidisciplinary total parenteral nutrition team improve patient outcomes? A systematic review. *JPEN J Parenter Enteral Nutr* 2004;28:251.
13. Ayers P, Adams S, Boullata J, et al. A.S.P.E.N. parenteral nutrition safety consensus recommendations. *JPEN J Parenter Enteral Nutr* 2014; 38:296.
14. Boullata JI, Gilbert K, Sacks G, et al. A.S.P.E.N. clinical guidelines: parenteral nutrition ordering, order review, compounding, labeling, and dispensing. *JPEN J Parenter Enteral Nutr* 2014; 38:334.
15. Guenter P, Boullata JI, Ayers P, et al. Standardized Competencies for Parenteral Nutrition Prescribing: The American Society for Parenteral and Enteral Nutrition Model. *Nutr Clin Pract* 2015; 30:570.
16. Slattery E, Rumore MM, Douglas JS, Seres DS.

- 3-in-1 vs 2-in-1 parenteral nutrition in adults: a review. *Nutr Clin Pract* 2014; 29:631.
17. Manzanares W, Dhaliwal R, Jurewitsch B, et al. Parenteral fish oil lipid emulsions in the critically ill: a systematic review and meta-analysis. *JPEN J Parenter Enteral Nutr* 2014;38:20.
 18. Edmunds CE, Brody RA, Parrott JS, et al. The effects of different IV fat emulsions on clinical outcomes in critically ill patients. *Crit Care Med* 2014; 42:1168.
 19. Hall TC, Bilku DK, Al-Leswas D, et al. A randomized controlled trial investigating the effects of parenteral fish oil on survival outcomes in critically ill patients with sepsis: a pilot study. *JPEN J Parenter Enteral Nutr* 2015; 39:301.
 20. Abbasoglu O, Hardy G, Manzanares W, Pontes-Arruda A. Fish Oil-Containing Lipid Emulsions in Adult Parenteral Nutrition: A Review of the Evidence. *JPEN J Parenter Enteral Nutr* 2019; 43:458.
 21. Pradelli L, Mayer K, Klek S, et al. ω -3 Fatty-Acid Enriched Parenteral Nutrition in Hospitalized Patients: Systematic Review With Meta-Analysis and Trial Sequential Analysis. *JPEN J Parenter Enteral Nutr* 2020; 44:44.
 22. Mirtallo JM, Ayers P, Boullata J, et al. ASPEN Lipid Injectable Emulsion Safety Recommendations, Part 1: Background and Adult Considerations. *Nutr Clin Pract* 2020;35:769.
 23. Zhang YP, Wan YD, Sun TW, et al. Association between vitamin D deficiency and mortality in critically ill adult patients: a meta-analysis of cohort studies. *Crit Care* 2014;18:684.
 24. Amrein K, Schnedl C, Holl A, et al. Effect of high-dose vitamin D3 on hospital length of stay in critically ill patients with vitamin D deficiency: the VITdAL-ICU randomized clinical trial. *JAMA* 2014; 312:1520.
 25. Heyland DK, Dhaliwal R, Drover JW, et al. Canadian clinical practice guidelines for nutrition support in mechanically ventilated, critically ill adult patients. *JPEN J Parenter Enteral Nutr* 2003; 27:355.
 26. Grau T, Bonet A, Miñambres E, et al. The effect of L-alanyl-L-glutamine dipeptide-supplemented total parenteral nutrition on infectious morbidity and insulin sensitivity in critically ill patients. *Crit Care Med* 2011; 39:1263.
 27. Heyland D, Muscedere J, Wischmeyer PE, et al. A randomized trial of glutamine and antioxidants in critically ill patients. *N Engl J Med* 2013; 368:1489.
 28. Jones JS, Tidwell B, Travis J, et al. Nutritional support of the hospitalized patient: a team approach. *J Miss State Med Assoc* 1995; 36:91.
 29. Fonseca G, Burgermaster M, Larson E, Seres DS. The Relationship Between Parenteral Nutrition and Central Line-Associated Bloodstream Infections. *JPEN J Parenter Enteral Nutr* 2017; :148607116688437.
 30. Yilmaz G, Koksali I, Aydin K, et al. Risk factors of catheter-related bloodstream infections in parenteral nutrition catheterization. *JPEN J Parenter Enteral Nutr* 2007; 31:284.
 31. da Silva JSV, Seres DS, Sabino K, et al. ASPEN Consensus Recommendations for Refeeding Syndrome. *Nutr Clin Pract* 2020; 35:178.
 32. Mattioli S, Miglioli M, Montagna P, et al. Wernicke's encephalopathy during total parenteral nutrition: observation in one case. *JPEN J Parenter Enteral Nutr* 1988; 12:626.
 33. Petrov MS, Zagainov VE. Influence of enteral versus parenteral nutrition on blood glucose control in acute pancreatitis: a systematic review. *Clin Nutr* 2007; 26:514.
 34. Mehanna HM, Moledina J, Travis J. Refeeding syndrome: what it is, and how to prevent and treat it. *BMJ* 2008; 336:1495.