

REVIEW ARTICLE

Vigilance, Not Panic: Understanding the Emerging COVID-19 Variant NB.1.8.1

Durre Aden¹, Nadeem Ahmad², Sufian Zaheer³

HOW TO CITE THIS ARTICLE:

Durre Aden, Nadeem Ahmad, Sufian Zaheer. Vigilance, Not Panic: Understanding the Emerging COVID-19 Variant NB.1.8.1. Ind Jr of Path: Res and Practice 2026; 15(2) 133-139.

ABSTRACT

The emergence of the NB.1.8.1 variant of COVID-19 in India marks another critical juncture in the ongoing trajectory of the pandemic. A progeny of the Omicron BA.2.86 lineage, NB.1.8.1 harbours multiple mutations in the spike protein, particularly in the receptor-binding domain(RBD), like F486P, E484K, N501Y, which raise concerns about increased infectivity and potential immune escape. Although it has not been designated a Variant of Concern by the World Health Organisation (WHO), its genetic profile warrants close monitoring. Clinically, the variant primarily presents with mild upper respiratory and systemic symptoms; however, high-risk individuals remain vulnerable to complications. Current diagnostic modalities, including RT-PCR and antigen-based assays, remain effective, while genomic surveillance is essential for accurate variant detection. Early reports suggest no increase in severity; however, high transmissibility of the NB.1.8.1 variant could still place significant strain on healthcare systems. Vaccination continues to provide strong protection against severe illness, though booster uptake remains suboptimal, particularly among the elderly and immunocompromised patients. While reinfection is possible, hybrid immunity and T-cell-mediated response offer a degree of protection. As NB.1.8.1 spreads, India must respond with scientific vigilance rather than complacency by strengthening surveillance, updating vaccines, reinforcing public health measures, and maintaining community awareness. Adaptive, evidence-based strategies are urgently required to address the dynamic mutational landscape of the SARS-CoV-2 virus.

KEYWORDS

- COVID-19 • NB.1.8.1 COVID-19 Variants • severe acute respiratory syndrome
- Coronavirus 2 • SARS-CoV-2

AUTHOR'S AFFILIATION:

¹ Assistant Professor, Department of Pathology, AIIMS, New Delhi, India.

² Associate Professor, Department of microbiology, University College of Medical Sciences, Delhi, India.

³ Professor, Department of Pathology, Vardhman Mahavir Medical College and Safdarjung Hospital, New Delhi, India.

CORRESPONDING AUTHOR:

Sufian Zaheer, Professor, Department of Pathology, Vardhman Mahavir Medical College and Safdarjung Hospital, New Delhi, India.

E-mail: sufianzaheer@gmail.com

➤ Received: 28-03-2026 ➤ Accepted: 29-04-2026



Creative Commons Non Commercial CC BY-NC: This article is distributed under the terms of the Creative Commons Attribution NonCommercial 4.0 License (<http://www.creativecommons.org/licenses/by-nc/4.0/>) which permits non-Commercial use, reproduction and distribution of the work without further permission provided the original work is attributed as specified on the Red Flower Publication and Open Access pages (<https://www.rfppl.co.in>)

INTRODUCTION

The recent detection of the SARS-CoV-2 variants NB.1.8.1 and LF.7 in India has led to public concern, especially as cases rise across parts of Southeast Asia. However, it is important to assess this development with evidence-based reasoning. As the pandemic evolves, the appearance of new viral lineages demands attention due to their chances of transmission, severity of infection, vaccine effectiveness, and the burden of long-term complications following infection.¹⁻³

A study from the Indian SARS-CoV-2 Genomics Consortium (INSACOG), NB.1.8.1 was detected in Tamil Nadu in April 2025, while four cases of LF.7 were identified in Gujarat in May. Both are subvariants of Omicron, likely deriving from the currently dominant JN.1 lineage. Recent reports indicate that JN.1 comprises over half (53%) of sequenced SARS-CoV-2 samples in India, followed by BA.2-related variants (26%) and other Omicron sublineages (20%).⁴

As of May 23, 2025, the WHO has categorised NB.1.8.1 and LF.7 as Variants Under Monitoring (VUM), suggesting it may possess notable genetic changes but do not yet demonstrate a clear epidemiological impact.⁵ Early data from the WHO suggests that NB.1.8.1 poses a low to intermediate public health risk. While it carries spike mutations (A435S, V445H, T478I) which might be associated with increased transmission and immune evasion, current evidence does not indicate greater severity compared to previous variants.⁶

Several states have reported recent increases in COVID-19 cases. For instance, in May 2025, Delhi reported 23 new cases, Andhra Pradesh four, and Telangana one. Bengaluru registered a positive case in a nine-month-old child. Kerala reported 273 new infections during the same month.^{7,8} Despite these numbers, most infections appear mild. Clinicians such as Dr. Aviral Mathur and Dr. Dhruv Chauhan have stated that the dominant strain, JN.1, is generally associated with mild symptoms and does not warrant panic, though adherence to standard hygiene practices remains advisable.^{9,10}

In the United States, NB.1.8 has also been identified. Experts from the CDC and FDA, including Drs. Natalie Thornburg and Jerry Weir, have noted that SARS-CoV-2 now

appears to exhibit seasonal fluctuations, with new dominant variants often emerging from existing lineages.¹¹⁻¹³

Genetic Features and Evolutionary Context of NB.1.8.1

NB.1.8.1 is a sublineage of Omicron BA.2.86 and contains several spike protein mutations. Of particular note are changes in the RBD that influence interaction with the ACE2 receptor and neutralising antibodies.^{14,15} Notable mutations include F486P, E484K, and N501Y were previously linked to increased binding affinity and immune evasion in earlier variants.¹⁶

The emergence of NB.1.8.1 is part of a broader trend observed with other Omicron sublineages, such as LB.1 (also called the "FLiRT" variant), which saw a spike in emergency department visits in early 2024.¹⁷ NB.1.8.1 evolved from XDV 1.5.1 and features additional spike protein mutations compared to LP 8.1 and JN.1.¹⁸ Preliminary studies from China suggest NB.1.8.1 exhibits enhanced binding affinity to cells, which may contribute to its higher transmissibility.¹⁹

Although not classified as a Variant of Concern [VOC], the variant's ability to partially escape immunity, particularly in those with older vaccine-induced protection, is a matter for ongoing study. Enhanced genomic surveillance remains critical.⁴

Clinical Presentation

Current reports suggest that symptoms of NB.1.8.1 largely resemble previous Omicron infections, with mild respiratory complaints like sore throat, nasal congestion, mild fever, fatigue, and occasional gastrointestinal issues such as nausea or mild diarrhoea.²⁰⁻²⁵ Loss of smell and taste has become less common. These symptoms have been predominantly mild, especially in individuals with prior infections or vaccination.²⁶⁻²⁹

No significant increase in hospitalisations or mortality has been attributed to NB.1.8.1 or LF.7. Nevertheless, older adults, those with chronic conditions, and immunocompromised individuals remain vulnerable. The variant's potential to escape immune responses may reduce the effectiveness of older vaccine-induced immunity.³⁰⁻³¹

Disease Severity and Hospital Trends

At present, NB.1.8.1 does not appear to cause serious disease like earlier Omicron strains. Hospitalisation rates remain low. However, widespread transmission could increase total case numbers, indirectly leading to more hospital admissions. Coinciding respiratory infections and low booster coverage may further strain healthcare systems.^{2,32-35}

India's public health infrastructure, including the Integrated Disease Surveillance Programme (IDSP) and the Indian Council of Medical Research (ICMR), has maintained active monitoring. The Directorate General of Health Services has reported no need for hospitalisation in most current cases.^{36,37} States like Delhi have issued advisories to ensure hospital readiness and emphasised continued testing, isolation, and genome sequencing.³⁸

Diagnostics

RT-PCR remains the most reliable test for COVID-19 diagnosis. Most commercial assays can detect NB.1.8.1 as it does not exhibit deletions in commonly targeted genomic regions. While rapid antigen tests are still in use, their sensitivity is variable. Broader sequencing, especially in clinical clusters and among international travellers, is necessary for tracking variant spread.³⁹⁻⁴⁰

Public Health Recommendations

Fundamental prevention measures continue to be relevant: mask-wearing in closed or crowded spaces, regular hand hygiene, and isolating when symptomatic. States are advised to maintain screening and isolation protocols, particularly in institutions and transport hubs.⁴¹⁻⁴⁷

Population Immunity and Reinfection Risks

India's population exhibits variable immunity. Some individuals have hybrid immunity (infection plus vaccination), while others may have waning protection due to time since last dose or infection.⁴⁸⁻⁵¹ Reinfections are expected, particularly among those previously infected with pre-Omicron strains. T-cell-mediated immunity remains relatively preserved, even as antibody levels decline.⁵²⁻⁵⁵

While memory B-cell responses to Omicron have diminished, T-cell responses across CD4+ and CD8+ subsets remain robust. This may protect against severe outcomes despite

reduced neutralising antibody levels.⁵⁴⁻⁵⁷

Vaccine Effectiveness

India has achieved high coverage for primary and first booster doses, but uptake of the second booster, particularly among the elderly, is lower.⁵⁸⁻⁵⁹ Studies indicate that vaccines such as Covishield and Covaxin continue to protect against severe illness, though their ability to prevent infection is reduced due to ongoing mutations.⁶⁰⁻⁶² Updated mRNA vaccines and nasal formulations tailored to Omicron sublineages may offer additional benefits.⁶³⁻⁶⁵ Adaptive policy responses, including flexible booster schedules and improved surveillance, should be prioritised.⁶⁶⁻⁶⁷ Vaccination has played a critical role in decreasing the hospital stay and long COVID.⁶⁸⁻⁷⁰

The NB.1.8.1 variant is another reminder that SARS-CoV-2 continues to evolve. Although current data does not suggest a significant public health threat, continuous monitoring is essential.⁷¹⁻⁷³ Strengthened genomic surveillance, updated vaccines, and renewed emphasis on preventive measures will be key to managing its spread. While no immediate cause for alarm exists, appropriate preparedness grounded in scientific evidence remains essential for mitigating the impact of future surges.

CONCLUSION

As India reports fresh cases of the new COVID-19 variant NB.1.8.1, the public health narrative again finds itself at a familiar crossroads by balancing vigilance with preparedness. This variant, marked by mutations that may enhance immune escape and mimic symptoms of earlier strains, explains the ongoing mutation of this virus under immune and environmental pressures. While the threat level remains moderate, complacency could prove detrimental.

A science-driven, proactive approach is essential. Strengthening genomic surveillance linked to clinical outcomes, reinstating masking and testing in high-risk areas, and expediting booster vaccinations, especially for vulnerable groups, is critical. Rapid approval and deployment of updated vaccines targeting new variants must be prioritised. Simultaneously, consistent public health messaging and community adherence to COVID-appropriate behaviour remain key. This evolving situation

calls not for panic but for preparedness, guided by scientific vigilance and coordinated action across healthcare, policy, and public domains.

REFERENCES

1. <https://timesofindia.indiatimes.com/toireporter/author-toi-news-desk-479263436.cms>. [Accessed on: May 20, 2025]
2. Aden D., Zaheer S., Kumar R., Raj S., Khan T., Varshney S. Beyond COVID-19 and SARS-CoV-2, cardiovascular outcomes of "long COVID" from a pathological perspective - a look back and road ahead. *Pathol Res Pract*. 2022 Nov; 239: 154144. doi: 10.1016/j.prp.2022.154144
3. SARS-CoV-2 variants of concern as of 25 April 2025 <https://www.ecdc.europa.eu/en/covid-19/variants-concern> [Accessed on: May 20, 2025]
4. <https://dbtindia.gov.in/insacog> [Accessed on May 25, 2025]
5. WHO TAG-VE Risk Evaluation for SARS-CoV-2 Variant Under Monitoring: NB.1.8.1.1.8.1. https://cdn.who.int/media/docs/default-source/documents/epp/tracking-sars-cov-2/23052025_nb.1.8.1.1.8.1.ire.pdf?sfvrsn=7b14df58_4. [Accessed on: May 22, 2025]
6. Tracking SARS-CoV-2 variants. <https://www.who.int/activities/tracking-SARS-CoV-2-variants>. [Accessed on: May 25, 2025]
7. <https://www.timesnownews.com/india/273-cases-in-kerala-in-may-23-in-delhi-covid-19-cases-surge-across-india-experts-say-rise-not-a-worry-article-151709665>. [Accessed on May 20, 2025]
8. <https://news.abplive.com/health/covid-update-nb181-lf7-variants-found-in-india-amid-surge-scare-1-death-reported-in-thane-1774705>. [Accessed on: May 25, 2025]
9. No need to panic over new Covid variant, precautions still key: Doctors. <https://www.tribuneindia.com/news/delhi/no-need-to-panic-over-new-covid-variant-precautions-still-key-doctors>. [Accessed on: May 26, 2025]
10. Should you be worried about Covid variant JN.1? Check what doctors are saying. <https://economictimes.indiatimes.com/news/new-updates/should-you-be-worry-about-jn-1-heres-what-doctors-are-saying-regarding-new-covid-variant/articleshow/121390309.cms?from=mdr>. [Accessed on: May 26, 2025]
11. Park G.W., Reija B., Tamin A., Hicks H., Flanders M.H., Metz J.M., *et al.* Comparative analysis of environmental persistence of SARS-CoV-2 variants and seasonal coronaviruses. *Appl Environ Microbiol*. 2025 Apr 23; 91(4): e0168824. Doi: 10.1128/aem.01688-24
12. <https://economictimes.indiatimes.com/news/international/global-trends/new-covid-variant-nb-1-8-1-detected-at-us-airports-amid-rise-in-cases-check-for-symptoms-and-preventive>. [Accessed on: May 24, 2025]
13. New Covid variant NB.1.8.1 detected at US airports amid rise in cases. Check for symptoms and preventive measures
14. New Covid variant NB.1.8.1 detected at US airports amid rise in cases. Check for symptoms and preventive measures. <https://economictimes.indiatimes.com/news/international/global-trends/new-covid-variant-nb-1-8-1-detected-at-us-airports-amid-rise-in-cases-check-for-symptoms-and-preventive-measures/articleshow/121380180.cms?From=mdr>. [Accessed on: May 23, 2025]
15. https://cdn.who.int/media/docs/default-source/documents/epp/tracking-sars-cov-2/23052025_nb.1.8.1.1.ire.pdf?sfvrsn=7b14df58_4. [Accessed on: May 24, 2025]
16. Xue S., Han Y., Wu F., Wang Q. Mutations in the SARS-CoV-2 spike receptor binding domain and their delicate balance between ACE2 affinity and antibody evasion. *Protein Cell*. 2024 May 28;15(6):403-418. doi: 10.1093/procel/pwae007.
17. Zabidi N.Z., Liew H.L., Farouk I.A., Puniyamurti A., Yip A.J.W., Wijesinghe V.N., *et al.* Evolution of SARS-CoV-2 Variants: Implications on Immune Escape, Vaccination, Therapeutic and Diagnostic Strategies. *Viruses*. 2023 Apr 10; 15(4): 944. doi: 10.3390/v15040944.
18. Aden D., Zaheer S. Investigating the FLiRT variants of COVID-19: Is it an emerging concern? *Pathol Res Pract*. 2024 Oct;262:155542. doi: 10.1016/j.prp.2024.155542.
19. World Health Organization Tracking SARS-CoV-2 Variants Available online: <https://www.who.int/activities/tracking-SARS-CoV-2-variants> [Accessed on: May 24, 2025]
20. Pather S., Charpentier N., van den Ouweland F., Rizzi R., Finlayson A., Salisch N., *et al.* Benefit-Risk Assessment of Vaccines by Technology Working Group (BRAVATO; ex-V3SWG). A Brighton Collaboration standardized template with key considerations for a benefit-risk assessment for the Comirnaty COVID-19 mRNA vaccine. *Vaccine*. 2024 Sep 17; 42(22): 126165. doi: 10.1016/j.vaccine.2024.126165.
21. New Covid variant NB.1.8.1 detected at US airports amid rise in cases. Check for symptoms and preventive measures. [https://m.economictimes.com/news/international/global-trends/new-covid-](https://m.economictimes.com/news/international/global-trends/new-covid-variant-nb-1-8-1-detected-at-us-airports-amid-rise-in-cases-check-for-symptoms-and-preventive-measures)

- variant-nb-1-8-1-detected-at-us-airports-amid-rise-in-cases-check-for-symptoms-and-preventive-measures/articleshow/121380180.cms. [Accessed on: May 25, 2025]
22. <https://m.economictimes.com/news/india/new-covid-nb-1-8-1-and-lf-7-subvariants-detected-in-india-as-cases-tick-up-in-pockets/articleshow/121377974.cms>. [Accessed on: May 24, 2025]
23. Jamie Ducharme. What to Know About the 'FLiRT' Variants of COVID-19. <https://time.com/6972143/covid-19-FLiRT-variants-kp-2/>. [Accessed on: May 24, 2025]
24. Carabelli A.M., Peacock T.P., Thorne L.G., Harvey W.T., Hughes J.; COVID-19 Genomics UK Consortium; Peacock S.J., Barclay W.S., de Silva T.I., Towers G.J., Robertson D.L. SARS-CoV-2 variant biology: immune escape, transmission and fitness. *Nat Rev Microbiol.* 2023 Mar; 21(3): 162-177. doi: 10.1038/s41579-022-00841-7.
25. <https://www.onlymyhealth.com/symptoms-of-new-covid-variants-nb181-and-lf7-12977831770>. [Accessed on: May 26, 2025]
26. <https://www.cdc.gov/covid/signs-symptoms/index.html>. [Accessed on: May 24, 2025]
27. Koyama A.K., Siegel D.A., Oyegun E., Hampton W., Maddox N., Koumans E.H.. Symptoms Reported With New Onset of Loss of Taste or Smell in Individuals With and Without SARS-CoV-2 Infection. *JAMA Otolaryngol Head Neck Surg.* 2021 Oct 1;147(10):911-914. doi: 10.1001/jamaoto.2021.2239.
28. Adly H.M., Saleh S.A.K., Garout M.A., Abdulkhaliq A.A., Khafagy A.A., Saati A.A. Post COVID-19 Symptoms Among Infected Vaccinated Individuals: A Cross-Sectional Study in Saudi Arabia. *J Epidemiol Glob Health.* 2023 Dec; 13(4): 740-750. doi: 10.1007/s44197-023-00146-9
29. <https://m.economictimes.com/news/new-updates/nb-1-8-1-lf-7-variants-of-covid-detected-in-india-check-if-its-deadly-symptoms-preventive-measures-other-details-covid-news/articleshow/121391901>. [Accessed on: May 26, 2025]
30. <https://www.livemint.com/news/india/most-covid-19-infections-mild-health-ministry-urges-vigilance-as-new-variant-jn-1-detected-all-you-need-to-know-11748137753450.html>. [Accessed on: May 24, 2025]
31. Lamba S., Ganesan S., Daroch N., Paul K., Joshi S.G., Sreenivas D. *et al.* SARS-CoV-2 infection dynamics and genomic surveillance to detect variants in wastewater - a longitudinal study in Bengaluru, India. *Lancet Reg Health Southeast Asia.* 2023 Apr; 11: 100151. doi: 10.1016/j.lansea.2023.100151.
32. Jian F., Yisimayi A., Song W., Wang J., Xu Y., Chen X *et al.* Humoral immunogenicity comparison of XBB and JN.1 in human infections. *bioRxiv* 2024.04.19.590276. doi: 10.1101/2024.04.19.590276.
33. India detects cases of new NB.1.8.1 and LF.7 COVID strains; experts say no need to panic. <https://www.newindianexpress.com/nation/2025/May/25/india-detects-cases-of-new-nb181-and-lf7-covid-strains-experts-say-no-need-to-panic>. [Accessed on: May 26, 2025].
34. <https://www.cdc.gov/covid/long-term-effects/index.html>. [Accessed on: May 26, 2025]
35. Centers for Disease Control and Prevention (CDC). *Seasonal Influenza and COVID-19: Considerations for Health Systems.* <https://www.cdc.gov>. [Accessed on: May 26, 2025].
36. World Health Organization (WHO). *Risk Communication and Community Engagement for Influenza and COVID-19.* <https://www.who.int>. [Accessed on: May 26, 2025].
37. <https://www.thehindu.com/sci-tech/health/whats-behind-the-recent-covid-19-spike-in-india/article69605844>. [Accessed on: May 25, 2025].
38. <https://idsp.mohfw.gov.in/>. [Accessed on: May 26, 2025].
39. <https://www.aninews.in/news/national/general-news/delhi-government-issues-advisory-to-hospitals-amid-rising-covid-19-cases20250523222229> Refresher training for staff should be conducted. [Accessed on: May 26, 2025].
40. <https://www.cdc.gov/coronavirus/2019-ncov/symptoms-testing/testing.html>. [Accessed on: May 26, 2025].
41. Shahid S., Parveen S., Vyas N., Abuhasaballah H.H.A., Annadurai S., El Assadi S.A. *et al.* Tracking of SARS-CoV-2 Variants of Concern by RT-PCR Assay in the Post-Pandemic Era. *New Emirates Medical Journal.* 2024; 5: Volume 5, 0250-6882. doi: 10.2174/0102506882268552230922112521.
42. Lerner A.M., Folkers G.K., Fauci A.S. Preventing the Spread of SARS-CoV-2 With Masks and Other "Low-tech" Interventions. *JAMA.* 2020; 324(19): 1935-1936. doi:10.1001/jama.2020.21946
43. Aden D., Zaheer S., Raj S. Challenges faced in the cancer diagnosis and management-COVID-19 pandemic and beyond-Lessons for future. *Heliyon.* 2022 Dec 5; 8(12). doi: 10.1016/j.heliyon.2022.e12091
44. <https://www.cdc.gov/clean-hands/about/hand-sanitizer.html>. [Accessed on: May 26, 2025]

45. <https://www.nhs.uk/conditions/covid-19/covid-19-symptoms-and-what-to-do/> Avoid unnecessary gatherings, especially for vulnerable individuals during periods of increased viral activity. [Accessed on: May 26, 2025]
46. <https://www.cdc.gov/infection-control/hcp/isolation-precautions/precautions.htm>. [Accessed on: May 26, 2025]
47. Aden D., Zaheer S., Kumar R, Ranga S. Monkeypox (Mpox) outbreak during COVID-19 pandemic-Past and the future. *J Med Virol.* 2023 Apr; 95(4):doi: 10.1002/jmv.28701.
48. <https://www.who.int/emergencies/diseases/novel-coronavirus-2019/advice-for-public>. [Accessed on: May 26, 2025]
49. Menegale F., Manica M., Zardini A., *et al.* Evaluation of Waning of SARS-CoV-2 Vaccine-Induced Immunity: A Systematic Review and Meta-analysis. *JAMA Netw Open.* 2023; 6(5): e2310650. doi:10.1001/jamanetworkopen.2023.10650
50. Shete A.M., Patil D.Y., Sahay R.R., Sapkal G.N., Deshpande G.R., Yadav P.D. Waning natural and vaccine-induced immunity leading to reinfection with SARS-CoV-2 Omicron variant. *Hum Vaccin Immunother.* 2022 Nov 30; 18(6): 2127289. doi: 10.1080/21645515.2022.2127289.
51. Khoury D.S., Schlub T.E., Cromer D., Steain M., Fong Y., Gilbert P.B., *et al.* Correlates of Protection, Thresholds of Protection, and Immunobridging among Persons with SARS-CoV-2 Infection. *Emerg Infect Dis.* 2023 Feb; 29(2): 381-388. doi: 10.3201/eid2902.221422
52. Murhekar M.V., Bhatnagar T., Thangaraj J.W.V., Saravanakumar V., Kumar M.S., Selvaraju S., ICMR Sero surveillance Group. SARS-CoV-2 seroprevalence among the general population and healthcare workers in India, December 2020-January 2021. *Int J Infect Dis.* 2021 Jul; 108: 145-155. doi: 10.1016/j.ijid.2021.05.040.
53. Khan, K., Lustig, G., Römer, C. *et al.* Evolution and neutralization escape of the SARS-CoV-2 BA.2.86 subvariant. *Nat Commun* 14, 8078 (2023). <https://doi.org/10.1038/s41467-023-43703-3>
54. Bergeri I., Whelan M.G., Ware H., Subissi L., Nardone A., Lewis H.C. Global SARS-CoV-2 seroprevalence from January 2020 to April 2022: A systematic review and meta-analysis of standardized population-based studies. *PLoS Med.* 2022 Nov 10; 19(11): e1004107. doi: 10.1371/journal.pmed.1004107.
55. Tarke A., Coelho C.H., Zhang Z., Dan J.M., Yu E.D., *et al.* SARS-CoV-2 vaccination induces immunological T cell memory able to cross-recognize variants from Alpha to Omicron. *Cell.* 2022 Mar 3; 185(5): 847-859.e11. doi: 10.1016/j.cell.2022.01.015.
56. Keeton, R., Tincho, M.B., Ngomti, A. *et al.* T cell responses to SARS-CoV-2 spike cross-recognize Omicron. *Nature.* 2022; 603: 488-492. doi: 10.1038/s41586-022-04460-3
57. Lau J.J., Cheng SMS, Leung K., Lee C.K., Hachim A., Tsang LCH, Real-world COVID-19 vaccine effectiveness against the Omicron BA.2 variant in a SARS-CoV-2 infection-naïve population. *Nat Med.* 2023 Feb; 29(2): 348-357. doi: 10.1038/s41591-023-02219-5.
58. Mukhopadhyay L., Gupta N., Yadav P.D., Aggarwal N. Neutralization assays for SARS-CoV-2: Implications for assessment of protective efficacy of COVID-19 vaccines. *Indian J Med Res.* 2022 Jan; 155(1): 105-122. doi: 10.4103/ijmr.ijmr_2544_21.
59. Sharma N., Basu S., Lalwani H., Rao S., Malik M., Garg S. COVID-19 Booster Dose Coverage and Hesitancy among Older Adults in an Urban Slum and Resettlement Colony in Delhi, India. *Vaccines (Basel).* 2023 Jun 29; 11(7): 1177. doi: 10.3390/vaccines11071177
60. Rizvi A.A., Singh A. Vaccination coverage among older adults: A population-based study in India. *Bull World Health Organ.* 2022 Jun 1; 100(6): 375-384. doi: 10.2471/BLT.21.287390.
61. Das S., Kar S.S., Samanta S., Banerjee J., Giri B., Dash S.K. Immunogenic and reactogenic efficacy of Covaxin and Covishield: A comparative review. *Immunol Res.* 2022 Jun; 70(3): 289-315. doi: 10.1007/s12026-022-09265-0. Epub 2022 Feb 22.
62. Mahumud R.A., Ali M.A., Kundu S., Rahman M.A., Kamara J.K., Renzaho A.M.N. Effectiveness of COVID-19 Vaccines against Delta Variant (B.1.617.2): A Meta-Analysis. *Vaccines (Basel).* 2022 Feb 11; 10(2): 277. doi: 10.3390/vaccines10020277.
63. Zhou Z., Zhu Y., Chu M. Role of COVID-19 Vaccines in SARS-CoV-2 Variants. *Front Immunol.* 2022 May 20; 13: 898192. doi: 10.3389/fimmu.2022.898192
64. Shanti Pather, Nicola Charpentier, Frank van den Ouweland, Ruben Rizzi, Andrew Finlayson, *et al.* A Brighton Collaboration standardized template with key considerations for a benefit-risk assessment for the Comirnaty COVID-19 mRNA Vaccine, Volume 42, Issue 22, 2024
65. Ella R., Reddy S., Jogdand H., Sarangi V., Ganneru B., Prasad S. *et al.* Safety and immunogenicity of an inactivated SARS-CoV-2 vaccine, BBV152: interim results from a double-blind, randomised, multicentre, phase 2 trial, and 3-month follow-up of a double-blind, randomised phase 1 trial. *Lancet Infect*

- Dis. 2021 Jul; 21(7): 950-961. doi: 10.1016/S1473-3099(21)00070-0.
66. Tobias J., Steinberger P., Wilkinson J., Klais G., Kundi M., Wiedermann U. SARS-CoV-2 Vaccines: The Advantage of Mucosal Vaccine Delivery and Local Immunity. *Vaccines* (Basel). 2024 Jul 18; 12(7): 795. doi: 10.3390/vaccines12070795.
 67. **Ministry of Health and Family Welfare, Government of India.** *Operational Guidelines for COVID-19 Preparedness and Response in India*. 2024 <https://www.mohfw.gov.in>. [Accessed on: May 26, 2025].
 68. **Centers for Disease Control and Prevention (CDC).** (2024). *Interim Clinical Considerations for Use of COVID-19 Vaccines*. <https://www.cdc.gov/vaccines/covid-19/clinical-considerations/interim-considerations-us.html>. [Accessed on: May 26, 2025].
 69. Yang, J., Rai, K.K., Alfred, T. *et al.* The impact of COVID vaccination on incidence of long COVID and healthcare resource utilisation in a primary care cohort in England, 2021–2022. *BMC Infect Dis* 25, 214 (2025). doi: 10.1186/s12879-024-10097-6
 70. Català M., Mercadé-Besora N., Kolde R., Trinh N.T.H., Roel E., Burn E., Rathod-Mistry T., *et al.* The effectiveness of COVID-19 vaccines to prevent long COVID symptoms: staggered cohort study of data from the UK, Spain, and Estonia. *Lancet Respir Med*. 2024 Mar; 12(3): 225-236. doi: 10.1016/S2213-2600(23)00414-9.
 71. Centres for Disease Control and Prevention. February 1, 2024. COVID Data Tracker. Centres for Disease Control and Prevention. May 11, 2024. [Accessed on: May 26, 2025]
 72. Soukhovolsky V., Kovalev A., Pitt A, Shulman K., Tarasova O., Kessel B. The Cyclicity of coronavirus cases: “Waves” and the “weekend effect”. *Chaos Solitons Fractals*. 2021 Mar; 144: 110718. doi: 10.1016/j.chaos.2021.110718. Epub 2021 Jan 28.
 73. Filip R., Gheorghita Puscaselu R, Anchidin-Norocel L., Dimian M., Savage W.K. Global Challenges to Public Health Care Systems during the COVID-19 Pandemic: A Review of Pandemic Measures and Problems. *J Pers Med*. 2022 Aug 7; 12(8): 1295. doi: 10.3390/jpm12081295