

Management of Tetralogy of Fallot with Major Aortopulmonary Collateral Arteries: Presurgical Endovascular Coil Embolization Experience

Dharmendra Kumar Srivastava¹, Navneet Srivastava², Abhishek Chauhan³, A P Jain⁴

How to cite this article:

Dharmendra Kumar Srivastava¹, Navneet Srivastava², Abhishek Chauhan *et al.* Management of Tetralogy of Fallot with Major Aortopulmonary Collateral Arteries: Presurgical Endovascular Coil Embolization Experience. J Cardiovasc Med Surg.2024;10(1-2):15-19.

Abstract

Tetralogy of Fallot with major aortopulmonary collateral arteries is a complex and heterogeneous condition varying degrees of severity, while MAPCA present in 20-25% with TOF and pulmonary atresia. Our institutional approach to this lesion emphasizes presurgical endovascular multiple coil embolization of significant bilateral MAPCAs along with complete intracardiac repair in the same sitting. The presurgical endovascular coil embolization of all MAPCAs of more than 3mm and complete intracardiac repair were done in the same sitting. The multiple Boston scientific pushable coils of diameters ranging. The presence of MAPCAs is a major determining factor in the prognosis and management of TOF. The role of the cardiac radiologist is not only in reporting critical components regarding collateral arteries and underlying cardiac structural disease but also in the management of TOF with MAPCAs along with cardiac surgeons.

Keywords: Aortopulmonary collateral; Coil; Endovascular; Tetralogy of fallot.

INTRODUCTION

Tetralogy of Fallot (TOF) with major aortopulmonary collateral arteries (MAPCAs) is a complex and heterogeneous condition varying

degrees of severity, while MAPCA present in 20-25% with TOF and pulmonary atresia. Our institutional approach to this lesion emphasizes presurgical endovascular multiple coil embolization of significant bilateral MAPCAs along with complete intracardiac repair in the same sitting.

Embryologic formation: Flowchart depicts the normal pulmonary vasculature development with regression of the segmental arteries as the pulmonary artery develops from the sixth branchial arch. PA pulmonary artery, PDA patent ductus arteriosus, RV right ventricle.^{1,2} (Fig. 1 & 2)

Author's Affiliation: ^{1,4}Professor, Department of Cardiovascular and Thoracic Surgery, ²Associate Professor, Department of Cardiovascular and Thoracic Surgery, ³Additional Professor, Department of Radio diagnosis, Dr. Ram Manohar Lohia Institute of Medical Sciences, Lucknow 226010, Uttar Pradesh, India.

Corresponding Author: Dharmendra Kumar Srivastava, Professor, Department of Cardiovascular Thoracic Surgery, Dr. Ram Manohar Lohia Institute of Medical Sciences, Lucknow. Uttar Pradesh-226010, India.

E-mail: drdks01@gmail.com

Received on: 24.02.2024

Accepted on: 25.06.2024

MATERIAL & METHODS

A 14-years-old boy presented with symptoms of Dyspnoea on exertion, bluish discoloration, signs



This work is licensed under a Creative Commons Attribution-NonCommercial-ShareAlike 4.0.

of cyanosis, and clubbing with oxygen saturation on room air 85-89%. Echocardiography and cardiac computerized tomography revealed ventricular septal defect (VSD) of 18mm in size, overriding of the aorta, severe right ventricular outlet obstruction, multiple MAPCAs of 4-6 mm in size, Diameters of main, right and left pulmonary arteries 11mm, 8mm and 7.6 mm respectively, right and left coronaries to appear normal. (Fig. 3 & 4) The presurgical endovascular coil embolization of all MAPCAs of more than 3mm and complete intracardiac repair were done in the same sitting. The multiple Boston scientific pushable coils of diameters ranging from 2-6 mm were used through a trans femoral route under digital subtraction angiography. (Fig. 5 & 6) The complete intracardiac

repair includes Dacron patch under moderate hypothermic cardiopulmonary bypass with cold blood cardioplegic arrest with topical cooling.

RESULTS

The cross-clamp and cardiopulmonary bypass times were 126 minutes and 178 minutes respectively. without any intracardiac blood flooding and complication. The patient was extubated within 6 hours of surgery and ICU stays were 6 days. At follow-up RVOT was found to be of good caliber with mild pulmonary valvular stenosis, no residual VSD, good LV systolic function and patient is on diuretics and beta blockers.




Fig. 1: Flowchart shows the development of collateral vessels with regard to the formation of ductus arteriosus. MAPCAS = Major Aortopulmonary Collateral Arteries, PA-VSD = Pulmonary Atresia with Ventricular Septal Defect, PDA = Patent Ductusarteriosus, TOF = Tetralogy of Fallot.




Fig. 2: Diagram representing the common sites of major aortopulmonary collateral arteries origin and classification. D4-D6 = Dorsal Vertebrae 4 to 6, IMA = Internal Mammary Artery.



Fig. 3: CT image shows dilated MAPCAs

DISCUSSION

With the development of medical technology and computed tomography imaging technology, cardiac surgeon gradually has advanced understanding about the MAPCAs, and their treatment has become an important part of intracardiac correction of TOF.^{3,4} The disadvantages of ligation of MAPCAs are surgical field is relatively fixed, creates big wounds, easily damages surrounding tissue, and

consumes time. Transcatheter occlusion, which preferably solves the disadvantages of ligating, is becoming a common method for treating collaterals in recent years.⁵ The occlusion can be performed before, during, or after operation. Most research on the management of MAPCAs before surgical correction of TOF; however, occlusion of MAPCAs before surgical correction could lead to a further decrease in arterial oxygen saturation, and the patient needed surgical correction immediately after transcatheter closure of MAPCAs.




Fig. 4: Descending thoracic aortic angiography image shows the MAPCAS.

Boston Scientific Pushable Coil Portfolio
Versatile and Cost-Effective

Six Shapes (L to R): Complex Helical, VortX™, VortX Diamond, Straight, Figure 8, Multi-Loop (2D Helical)

Coil Length	Coil Diameter									
	2 mm	3 mm	4 mm	5 mm	6 mm	7 mm	8 mm	9 mm	10 mm	11 mm
0-3cm										
4-6cm										
7-9cm										

0.018" Pushable Coils (Blue) 0.035" Pushable Coils (Yellow)

- Include highly thrombogenic, dense Dacron® fibers designed for fast and complete occlusion
- Have MRI conditional compatibility (up to 3.0 Tesla)
- Are recommended for use with continuous flush setup

© 2015 Boston Scientific Corporation. All Rights Reserved.

Boston Scientific Pushable Coil Portfolio
Shape Comparisons & Benefits

Shape	Complex Helical	Figure 8	VortX	Straight	Multi-Loop (2D Helical)	VortX Diamond
Coil Type	Framing	Filling	Filling	Straight	Framing	Filling
Benefit	Provide structure around the vessel wall	Short length fills small vessel area	Stackable shape helps to fill vessels tightly	Fill small accessory vessels prone to spasm	Provide structure around the vessel wall	Center coil adheres to vessel wall while sides taper down and fill area
Best for	The first and last coil placed in a vessel	Small vessels (20mm), such as the right gastric	Filling a segment of a vessel proximal to a framing coil	Tiny GI bleeds where vessel diameter < 2mm	The first and last coil placed in a vessel	Filling a segment of a vessel proximal to a framing coil

© 2015 Boston Scientific Corporation. All Rights Reserved.

Fig. 5: Boston coils of different shapes and sizes (Source: <https://www.bostonscientific.com/content/dam/bostonscientific/pi/product-catalog/2023-EMEA-PI-Product-Catalogue-Eluvia-Ranger-Update%282%29.pdf>)

CONCLUSION

The presence of MAPCAs is a major determining factor in the prognosis and management of TOF

The role of the cardiac radiologist is not only in reporting critical components regarding collateral arteries and underlying cardiac structural disease but also in the management of TOF with MAPCAs along with cardiac surgeons.



Fig. 6: Coils in aortopulmonary Collaterals

REFERENCES

1. Rabinovitch M, Herrera de Leon V, Castaneda AR, Reid L. Growth and development of the pulmonary vascular bed in patients with tetralogy of Fallot with or without pulmonary atresia. *Circulation*. 1981; 64:1234-1249.
2. Jefferson K, Rees S, Somerville J. Systemic arterial supply to the lungs in pulmonary atresia and its relation to pulmonary artery development. *Br Heart J*. 1972; 34:418-427.
3. Lapierre C., Dubois J., Rypens F., Raboisson M.-J., Déry J. Tetralogy of fallot: preoperative assessment with MR and CT imaging. *Diagnostic and Interventional Imaging*. 2016;97(5):531-541.
4. Perry S. B., Radtke W., Fellows K. Ekeane J. F., Lock J. E. Coil embolization to occlude aortopulmonary collateral vessels and shunts in patients with congenital heart disease. *Journal of the American College of Cardiology*. 1989; 13(1):100-108.
5. Alex A, Ayyappam A, et al major aortopulmonary collateral arteries. *Radiology Cardiothoracic Imaging* 2022;4(1) 1-12.

