

On Earth Harmony: Algae Crafting Environment Revival

Devanshi Trivedi¹, Vivek Pandya², Kinjal Upadhyay³

How to cite this article:

Devanshi Trivedi, Vivek Pandya, Kinjal Upadhyay. On Earth Harmony: Algae Crafting Environment Revival. J Microbiol Relat Res. 2024;10(1):29-37.

Abstract

In the intricate dance of nature, algae emerge not just as the silent architects of vibrant underwater realms, but as the unsung heroes with the power to breathe life into the natural world. In this comprehensive analysis we go through how algae jack into – carbon's void by sequestering carbon of CO₂ using photosynthesis and mitigates greenhouse emissions, purifies tainted waters by using wastes from wastewater like nutrients, sparks a green revolution as it produces biofuels and being used as food for nutrient enrichment as well as monitors environment as biosensors, thus concluding into algae's ability to transcending conventional boundaries of environmental stewardship. The more emphasis will be on illuminating on major benefits like biofuels, carbon sequestration and wastewater treatment which might be sweet in the citrus.

Keywords: Microalgae; Application; Wastewater treatment; Carbon sequestration; biofuels.

INTRODUCTION

The environment is deteriorating and different pollution is emerging as severe problems around the globe. Thus, rendering it to fundamental concern to ecotoxicologist, environmental biologist, eco-chemists, pathologist and many other fields of research. Therefore, advanced

methods redefining approaches meeting the demands of growing world; called 'green agenda'. The Algae being diverse photosynthetic group of eukaryotes diversified in many species and genera like Chlorophyta, dinoflagellates, diatoms, etc. they have emerged as valuable environmental applicators becoming the sustainable solutions across various sectors. Their ability of absorbing nutrients, in wastewater treatment, aiding water purification, forming biofuels and many more. Emerging environmental contaminants present in wastewater are drawing significant awareness as they exhibit several bad qualities such as high polarity, the ability to be bioaccumulated by aquatic organisms and resistance to biodegradation. They harm the aquatic ecosystem and human health as well. The most common contaminants include not only pharmaceuticals products but also several personal care products, perfluorinated compounds, gasoline additives, surfactants, organometallic compounds, disinfection by-products, brominated and organophosphate flame retardants, endocrine-disrupting compounds,

Author Affiliation: ^{1,2}Student, ³Assistant Professor, Department of Biochemistry and Biotechnology, St. Xaviers College (Autonomous), Ahmedabad 380009, Gujarat, India.

Corresponding Author: Kinjal Upadhyay, Assistant Professor, Department of Biochemistry and Biotechnology, St. Xaviers College (Autonomous), Ahmedabad 380009, Gujarat, India.

E-mail: kinjal.upadhyay@sxca.edu.in

Received on: 21.04.2024

Accepted on: 09.06.2024



This work is licensed under a Creative Commons
Attribution-NonCommercial-ShareAlike 4.0.

nanoparticles and plasticizers (Müller *et al.*, 2007; Morin-Crini *et al.*, 2022). Algae-based technologies have demonstrated greater efficiency in removing emerging contaminants (Morin-Crini *et al.*, 2022) at both laboratory scales and in real wastewater.

Bioremediation is basically the use of living biological organisms in order to remove the contaminants. Algae can be used in bioremediating industrial effluent including heavy metal, dye, nitrogen, phosphorus, etc. removal.

Heavy metal are the class with atomic density $>4000 \text{ kg m}^3$ (Vardhan *et al.*, 2019). They show accumulation and thus are having non-biodegradable properties, toxicity and thus be a critical problem worldwide. A heavy metal being pollutant also causes diseases like nervous system disorder, gastrointestinal and kidney dysfunction, vascular damage, skin lesion, immune system dysfunction and cancer even at low concentrations (Edelstein *et al.*, 2018).

Heavy metals have three categories: 1) radionuclides 2) precious metals 3) toxic metals (Pavithra *et al.*, 2020; Kafil *et al.*, 2022). The heavy metals can be removed by Biosorption and bioaccumulation (Ahmed *et al.*, 2022).

- First method, bioaccumulation works by transporting metal ions across cell membrane using passive and active transport. Method occurs only in live cells.

- Second one, biosorption involves the sorption of metals onto the cell surface which will be helped by Exopolysaccharides (Ahmed *et al.*, 2022).

Applications of algae:

Throughout the last several decades, the world frugality has expanded at a dramatic pace. Extreme population expansion is a major issue, with the world's population cast to reach 8.5 billion in 2030, 9.7 billion in 2050 and 10.9 billion by 2100 (United Nations, 2019). As a consequence of the growing global population, large quantities of energy and coffers have been consumed and pollution situations are high. The necessity of anticipating and preparing for these heads has been honored and appreciated by numerous transnational associations. Promoting a "green frugality" is one similar plan. The green frugality conception was developed during the 2012 United Nations Conference on Sustainable Development in Rio de Janeiro and it's grounded on the idea that environmental protection helps both the frugality and society. The thing of this conception is to empower growers and manufacturers to produce further environmentally friendly product and consumption systems grounded on exercise and recovering for sustainable development (Loiseau *et al.*, 2016).




Fig. 1: Applications of algae Adopted from Kandasamy *et al.*, 2022

Nutrients bioremediation by algae

External and agrarian waste water contains a large quantum of nutrients like nitrogen, phosphorus and other minerals. The predominant forms in which they do in wastewater are ammonium ions, nitrite, nitrate and orthophosphate. Phosphate enters the algal cell laboriously through a symporter with H or Na ions furnishing the driving force. Algae also hydrolyze organic phosphorus composites with membrane-bound as well as free phosphatases, releasing bioavailable phosphorus that's latterly taken up by the algal cells (Bolan *et al.*, 2004). Among inorganic nitrogen sources, algae preferentially take up ammonium because of its further stoutly favorable assimilation and direct protein objectification process (Bolan *et al.*, 2004). Algae uptake ammonium by a group of membrane transporter proteins belonging to the ammonium transporter family. On the other hand, nitrate and nitrite are reduced to ammonium, by nitrate reductase and nitrite reductase, independently, for intracellular uptake, which is energy ferocious.

Removal of contaminants by algae

Algae as being able to tolerate toxic elements and require heavy metals as like zinc, molybdenum, manganese, iron, cobalt, copper and boron as trace elements thus they can grow and metabolize whereas some like cadmium, chromium, lead, arsenic, mercury are harmful to them.

This called hormesis, is a process of requiring toxic heavy metals for growth. Precious metals Ag and Au can be recovered while removing toxic radioactive elements from water and these algae can tolerate toxicity via mechanisms like as gene regulation, heavy metal immobilization, chelation, exclusion and the production of different enzymes that decrease the toxicity of these substances. (Monteiro *et al.*, 2012; Tripathi and Poluri, 2021; Manikandan *et al.*, 2022)

Heavy metal control by algae majorly follows biosorption and bioaccumulation, it can form organometallic compound with heavy metals in cytoplasm and separating vacuoles. The biosorption is metal ion binding to dead or inactive algal cells via different interaction like electrostatic, chelation, ionic, etc. (Srimongkol *et al.*, 2022).

Instead bioaccumulation involve live algae where first the semi biosorption takes place with metal ion binding functional groups like -OH, -COOH, amine, etc. on wall of cell. Then second phase, intracellular uptake happens, ingesting these

toxic elements playing crucial role in detoxification. (Srimongkol *et al.*, 2022).

Some studies show dead algae being better in wastewater treatment as for live algae show restricted sorption due to the live cell poisoning; also live cells have complex intracellular mechanism for the uptake of contaminant which makes absorption complex. On the other hand dead ones will act as assemblers of polymers which absorbs metals at extracellular level. (Salam, 2019).

Algal species from family *Phaeophyceae* in which alginate serves mainly for heavy metal binding which directs the influence on biosorption. A weird matter of fact is some contaminant specifically boost rates of other contaminants; for example, sulfamethazine removal gets increase by presence of sulfamethoxazole (Xiong, 2019). Co-metabolism reported in *Chlamydomonas Mexicana* removes ciprofloxacin after adding sodium acetate, here breakdown via enzyme takes place (Xiong, 2019). Temperature effects as higher temperature increases rate of entire process (Vijayaraghavan and Yun, 2007).

Studies show that by product formed can may be proven more toxic than primary compound, one such case is textile water remediation using *Oscillatoria tenuis*, *Chlorella pyrenoidosa* and *Chlorella vulgaris* degrade azo dyes into simple aromatic amines being carcinogenic and persistent (Fazal, 2018). Thus, before using algae systems, the study should be made for effects of environmental factors on monitoring and growth of algae for successful application.




Fig. 2: Percentage removal of Nitrogen and phosphorus by different algal species

Pharmaceutical contaminants can be removed based on their physiological properties using microalgae. A study by De Godos *et al.*, 2012, showed ability of *Chlorella vulgaris* for removing more than 50% tetracycline via adsorption as interaction established between positively charged molecules on cell surface. Not only this but also florfenicol, trimethoprim, sulfamethoxazole and carbamazepine be bioaccumulated by *Chlorella sp.* (Song *et al.*, 2019). An algae *Scenedesmus dimorphus* biodegrades 85% of 17 α -estradiol in about 7 days (Zhang *et al.*, 2014). The biodegradability is determined by how complex structured the compound is, *i.e.* more complex the substrate then

less efficiency of degradability is seen and vice versa.

Now when studied for domestic effluent, (Wang *et al.* 2010) reported that sewage sludge (nutrient rich) cultivation of microalgae was more convenient and better than primary and secondary treated wastewater when parameters like COD, PO₄-P, NO₃-N and NH₄-N. Zheng *et al.*, 2018, mixed piggery wastewater with brewery wastewater for determining optimal C/N ratio as to know nutrient Removal and microalgal growth; at an optimum C/N ratio of 7.9, removal efficiency of all the above mentioned parameters reached 93%, 90%, 96% and 100%.




Fig 3: Percentage removal of dye using different algal species




Fig 4: Percentage removal of Cadmium Data Adopted from Bilal *et al.*, 2018




Fig 5: Percentage removal of Ibuprofen and Carbamazepine. Data Adopted from Abdelfattah *et al.*, 2023




Fig 6: Percentage removal of different pharmaceutical contaminants by *Nannochloris sp.* Data Adopted from Abdelfattah *et al.*, 2023




Fig 7: Percentage removal of different dyes by *Chlorella* sp. Data Adopted from Abdel Fattah *et al.*, 2023

Carbon sequestration

As said by IEA in 2021 that CO₂ amount is to be rebound and increase by 4.8% as 2019 reaches peaks of fossil fuel rebound with increasing demand, which pose the problem of increased green house pollution. Algae being magically active creatures can utilize this CO₂ via photosynthesis and recycle them in form of bioenergy which makes using microalgae a sustainable and environment friendly way (Brilman *et al.*, 2013).

CO₂ assimilation by photosynthesis involves both light dependent and light independent reactions; moreover this CO₂ is converted to carbohydrate

precursors and biomass which later synthesizes major biomolecules like lipids, proteins and nucleic acids (Zeng *et al.*, 2011; Gg and Liu, 2021).

A report stated, cost of manufacturing *Chlorella* Biomass is around \$4.87 per kg having consumed 0.96kWh per kg of energy (Valdovinos-García *et al.*, 2020). Moreover around 2.2 kilotons of CO₂/ year is sequestered using 4000m³ microalgae growth in pond systems when natural sunlight is available (Stewart and Hessami, 2005).

According to some other studies, algae might reduce CO₂ emissions by 50% (Stepan *et al.*, 2002).




Fig 8: Represents CO₂ sequestration process Adopted from Zhaom *et al.*, 2020

Biofuels

Biofuels are the fuels coming to rescue in the form of renewable energy as the threat of losing these petroleum based fossil fuels we are using now. Biofuels include ethanol, biodiesel, biohydrogen.

Microalgae have a capacity as a patron similar to that of a land factory, which has been critical to mortal survival for a long time in history considering terms of significant sources of food, drug, erecting accoutrements and energy force. Different photosynthetic unicellular microalgae are arising as new sources of renewable energy that can fulfill the demands of the conditioning. Microalgae lipids can be employed as a raw material for biodiesel conflation and remaining biomass rich in carbohydrates can be used to produce bioethanol or biogas. They generally accumulate lipids between 20 - 50% of their dry weight. Yet some species can be as high as 80% under certain condition (Chisti, 2007). These neutral lipids, substantially in the form of triacylglycerols (mark up to 90 - 95%), which can be converted to adipose acid methyl esters (FAMES) and converted biodiesel. Algae have advantages over first generation biofuels made from sugar, bounce and vegetable oil because of their high growth rates and productivity, capability to grow on a non-arable land using wastewater, capability to use water pollutants and CO₂ and capability to produce a variety of high-value natural product of biodiesel from microalgae comprises two different ways: 1) lipid birth from microalgal cells and 2) transesterification of lipids using alcohol and a catalyst (Mondal *et al.*, 2017). In a primary study of biomass product and ammonium junking in *Synechococcus sp.* VDW (accession number MH393765) insulated from natural seawater in Thailand (Tinpranee *et al.*, 2018), we discovered that at optimum conditions (original pH 7.4, inoculum size of 0.17 (OD730) and ammonium of 10.5 mg L⁻¹), maximum ammonium junking and biomass productivity were 95 and 34 mg L⁻¹ d⁻¹, independently. A review study by Pancha *et al.*, 2019, showed colorful microalgae lipid content cultivated in colorful wastewater ranging from 18-79% w/w of biomass. Meanwhile, Chinnasamy *et al.* (2010) set up that > 96% nutrient in treated wastewater containing 85-90% carpet assiduity backwaters with 10-15% external sewage which could be removed by a range of native algal isolates. Biomass product eventuality and lipid content of 92-17.8 tons/h/time and 6.82%, independently. Biohydrogen product from

Table 1: Percentage of Co₂ sequestered by different algal species

Algal species	% Co ₂ sequestered
Cyanothrix caldarium	100%
Scenedesmus sp.	80%
Cylindrocapsa littorale	60%
Euglena gracilis	50%
Eudorina spp.	20%
Nannochloris sp.	18%
Tetraselmis sp.	15%

Data Adopted from Sanyal *et al.*, 2019

microalgae can take place through in different routes but generally involves turmoil biohydrogen product (e.g., dark turmoil biohydrogen product, print turmoil biohydrogen product and print-dark combined turmoil biohydrogen product) and photosynthesis biohydrogen product (e.g., direct natural photolysis biohydrogen product, circular natural photolysis biohydrogen product) (Wang *et al.*, 2021). Batista *et al.*, 2015, reported that can grow in civic wastewater and also the biomass can be converted into biohydrogen through dark turmoil by *Enterobacter aerogenes* producing 56.8 ml H₂ per gvs (Ruiz- Marin *et al.*, 2020)

The Biogas is the end product of anaerobic digestion. Generally, anaerobic digestion is conducted by two processes: 1) simple sugar is instigated by fermentative bacteria and converted into alcohols through anaerobic condensation and 2) metanogenic microorganisms use these composites and synthesize biomethane (Danquah *et al.*, 2011; Choudhary *et al.*, 2016) revealed that native colleges PA6 has good nutrient junking capability from pastoral wastewater with a theoretical methane eventuality of over to 0.79 m³ VS per kg. Kinnunen and Rintala, 2016, showed that the biomethane eventuality of *C. Vulgaris* and mixed culture of native algae species (dominating by *Scenedesmus* sp.) varied between 154 and 252 L VS per kg of methanol. Depending on culture media including synthetic medium, wastewater (castrated and non-sterilized) and digestate from anaerobic digestion of pulp and paper biosludge (castrated and non-sterilized).

CONCLUSION

Algae as a boon can thus be used to clear up the environment by mitigating CO₂, reduce heavy metals by adsorption, clearing wastewater and still many others like making biofuels and being the

green gold of earth.

Conflict of Interest

There are no conflicts to declare

Acknowledgements

The authors are thankful to St. Xavier's College (Autonomous), Ahmedabad, India for providing a financial assistance under the scheme of Research Project Seed Grant having award number SXCA/2021-2022/PR-01.

REFERENCES

1. Abdelfattah, A., Ali, S. S., Ramadan, H., El-Aswar, E. I., Eltawab, R., Ho, S., Elsamahy, T., Li, S., El-Sheekh, M. M., Schagerl, M., Kornaros, M., & Sun, J. (2023). Microalgae-based wastewater treatment: Mechanisms, challenges, recent advances and future prospects. *Environmental Science and Ecotechnology*, 13, 100205. <https://doi.org/10.1016/j.esec.2022.100205>
2. Ahmad, S. F., Mofijur, M., Parisa, T. A., Islam, N., Kusumo, F., Inayat, A., et al. (2022). Progress and challenges of contaminate removal from wastewater using microalgae biomass. *Chemosphere* 286, 131656. doi:10.1016/j.chemosphere.2021.131656
3. Anand, U., Dey, S., Parial, D., Federici, S., Ducoli, S., Bolan, N., Dey, A., & Bontempi, E. (2023). Algae and bacteria consortia for wastewater decontamination and transformation into biodiesel, bioethanol, biohydrogen, biofertilizers and animal feed: a review. *Environmental Chemistry Letters*, 21(3), 1585–1609 <https://doi.org/10.1007/s10311-023-01562-w>
4. Batista, A. P., Ambrosano, L., Graça, S., Sousa, C., Marques, P. A., Ribeiro, B., et al. (2015). Combining urban wastewater treatment with biohydrogen production-an integrated microalgae-based approach. *Bioresour. Technol.* 184, 230–235. doi:10.1016/j.biortech.2014.10.064
5. Bilal, M., Rasheed, T., Sosa-Hernández, J. E., Raza, A., Nabeel, F., & Iqbal, H. M. (2018). Biosorption: An Interplay between Marine Algae and Potentially Toxic Elements – A Review. *Marine Drugs*, 16(2), 65. doi: 10.3390/md16020065. PMID: 29463058; PMCID: PMC5852493.
6. Bolan NS, Saggarr S, Luo JF, Bhandral R, Singh J (2004) Gaseous emissions of nitrogen from grazed pastures. In: DL Sparks (Ed.) Processes, measurements and modelling, environmental implications and mitigation, *Advances in Agronomy*, Vol. 84. Elsevier academic pressinc. [https://doi.org/10.1016/S0065-2113\(04\)840021](https://doi.org/10.1016/S0065-2113(04)840021)
7. Brilman, W., Alba, L. G. and Veneman, R. (2013). Capturing atmospheric CO₂ using supported amine sorbents for microalgae cultivation. *Biomass Bioenergy* 53, 39–47. doi:10.1016/j.biombioe.2013.02.042
8. C. Song, Y. Wei, Y. Qiu, Y. Qi, Y. Li, Y. Kitamura, Biodegradability and mechanism of florfenicol via *Chlorella* sp. UTEX1602 and L38: experimental study, *Bioresour. Technol.* 272 (2019) 529e534. doi:10.1016/j.biortech.2018.10.080
9. Chinnasamy, S., Bhatnagar, A., Hunt, R. W. and Das, K. C. (2010). Microalgae cultivation in a wastewater dominated by carpet mill effluents for biofuel applications. *Bioresour. Technol.* 101 (9), 3097–3105. doi:10.1016/j.biortech.2009.12.026
10. Chisti, Y. (2007). Biodiesel from microalgae. *Biotechnol. Adv.* 25, 294–306. doi:10.1016/j.biotechadv.2007.02.001
11. Choudhary, P., Prajapati, S. K. and Malik, A. (2016). Screening native microalgal consortia for biomass production and nutrient removal from rural wastewater for bioenergy applications. *Ecol. Eng.* 91, 221–230. doi:10.1016/j.ecoleng.2015.11.056
12. Danquah, M., Liu, B. and Harun, R. (2011). "Analysis of process configurations for bioethanol production from microalgal biomass," in *Progress in biomass and bioenergy production*. Editor S. S. Shaukat (London: IntechOpen).
13. Fazal T, Mushtaq A, Rehman F, Khan AU, Rashid N, Farooq W, Xu J (2018) Bioremediation of textile wastewater and successive biodiesel production using microalgae. *Renew Sustain Energy Rev* 82:3107–3126. <https://doi.org/10.1016/j.rser.2017.10.029>
14. H. Zheng, M. Liu, Q. Lu, X. Wu, Y. Ma, Y. Cheng, M. Addy, Y. Liu, R. Ruan, Balancing carbon/nitrogen ratio to improve nutrients removal and algal biomass production in piggery and brewery wastewater, *Bioresour. Technol.* 249 (2018) 479e486. doi:10.1016/j.biortech.2017.10.057.
15. I. de Godos, R. Muñoz, B. Guieysse, Tetracycline removal during wastewater treatment in high-rate algal ponds, *J. Hazard Mater.* 229 (2012) 446e449. <http://dx.doi.org/10.1016/j.jhazmat.2012.05.106>
16. Kandasamy, S., Zhang, B., He, Z., Bhuvanendran, N., EL-Seesy, A. I., Wang, Q., Narayanan, M., Thangavel, P., & Dar, M. A. (2022). Microalgae as a multipotential role in

- commercial applications: Current scenario and future perspectives. *Fuel*, 308, 122053. <https://doi.org/10.1016/j.fuel.2021.122053>
17. Kinnunen, V. and Rintala, J. (2016). The effect of low-temperature pretreatment on the solubilization and biomethane potential of microalgae biomass grown in synthetic and wastewater media. *Bioresour. Technol.* 221, 78–84. doi:10.1016/j.biortech.2016.09.017
 18. L. Wang, M. Min, Y. Li, P. Chen, Y. Chen, Y. Liu, Y. Wang, R. Ruan, Cultivation of green algae *Chlorella* sp. in different wastewater from municipal wastewater treatment plant, *Appl. Biochem. Biotechnol.* 162 (2010) 1174e1186. doi: 10.1007/s12010-009-8866-7
 19. Loiseau, E., Saikku, L., Antikainen, R., Droste, N., Hansjürgens, B., Pitkänen, K., et al. (2016). Green economy and related concepts: An overview. *J. Clean. Prod.* 139, 361–371. doi:10.1016/j.jclepro.2016.08.024
 20. Manikandan A, Suresh Babu P, Shyamalagowri S, Kamaraj M, Muthu- kumaran P, Aravind J (2022) Emerging role of microalgae in heavy metal bioremediation. *J Basic Microbiol* 62(3–4):330–347. <https://doi.org/10.1002/jobm.202100363>
 21. Mondal, M., Goswami, S., Ghosh, A., Oinam, G., Tiwari, O. N., Das, P., et al. (2017). Production of biodiesel from microalgae through biological carbon capture: A review. *3 Biotech.* 7 (2), 99. doi:10.1007/s13205-017-0727-4
 22. Monteiro CM, Castro PM, Malcata FX (2012) Metal uptake by micro- algae: underlying mechanisms and practical applications. *Bio-technol Prog* 28(2):299–311. <https://doi.org/10.1002/btpr.1504>
 23. Morin-Crini N, Lichtfouse E, Fourmentin M, Ribeiro ARL, Nout-sopoulos C, Mapelli F, Crini G (2022) Removal of emerging contaminants from wastewater using advanced treatments a review. *Environ Chem Lett* 20:1–43. <https://doi.org/10.1007/s10311-021-01379-5>
 24. Müller K, Magesan GN, Bolan NS (2007) A critical review of the influence of effluent irrigation on the fate of pesticides in soil. *AgrEcosyst Environ* 116–93:(4–2)120. <https://doi.org/10.1016/j.agee.2006.08.016>
 25. Pancha, I., Chokshi, K. and Mishra, S. (2019). "Industrial wastewater-based microalgal biorefinery: A dual strategy to remediate waste and produce microalgal bioproducts," in *Application of microalgae in wastewater treatment*. Editors S. Gupta and F. Bux (Cham: Springer), 173–193.
 26. Ruiz-Marin, A., Canedo-López, Y. and Chávez-Fuentes, P. (2020). Biohydrogen production by *Chlorella vulgaris* and *Scenedesmus obliquus* immobilized cultivated in artificial wastewater under different light quality. *Amb. Express* 10, 191. doi:10.1186/s13568-020-01129-w
 27. Salam, K. A. (2019). *Biofuel Research Journal*, 6(2), 948–961. <https://doi.org/10.18331/brj2019.6.2.2>
 28. Sanyal, D., Kodgire, S., Desai, D., Saxena, N., Singh, S., & Dasgupta, S. (2019). Commitment for a Cleaner India: Utilization of CO₂ and Sewage Wastewater by Green Algae *Scenedesmus* sp. Under Laboratory Conditions. In *Smart innovation, systems and technologies* (pp. 277–286). https://doi.org/10.1007/978-981-32-9578-0_25
 29. Srimongkol, P., Sangtanoo, P., Songserm, P., Watsuntorn, W., & Karnchanatat, A. (2022). Microalgae-based wastewater treatment for developing economic and environmental sustainability: Current status and future prospects. *Frontiers in Bioengineering and Biotechnology*, 10. <https://doi.org/10.3389/fbioe.2022.904046>
 30. Stepan, D. J., Shockey, R. E., Moe, T. A. and Dorn, R. (2002). Carbon dioxide sequestering using microalgal systems. Pittsburgh, PA: U.S: Department of Energy. <https://doi.org/10.2172/882000>
 31. Stewart, C. and Hessami, M. A. (2005). A study of methods of carbon dioxide capture and sequestration-the sustainability of a photosynthetic bioreactor approach. *Energy Convers. Manag.* 46, 403–420. doi:10.1016/j.enconman.2004.03.009
 32. Tinpranee, N., Incharoensakdi, A. and Phunpruch, S. (2018). Screening cyanobacteria from marine coastal waters of Thailand for biohydrogen production. *J. Appl. Phycol.* 30, 3471–3481. doi:10.1007/s10811-018-1490-6
 33. Tripathi S, Poluri KM (2021) Heavy metal detoxification mechanisms by microalgae: insights from transcriptomics analysis. *Environ Pollut* 285:117443. <https://doi.org/10.1016/j.envpol.2021.117443>
 34. Valdovinos-García, E. M., Barajas-Fernández, J., Olán-Acosta, M. D. L. Á., Petriz- Prieto, M. A., Guzmán-López, A. and Bravo-Sánchez, M. G. (2020). Techno- economic study of CO₂ captu_{re} of a thermoelectric plant using microalgae (*Chlorella vulgaris*) for production of feedstock for bioenergy. *Energies* 13 (2), 413. doi:10.3390/en13020413
 35. Vijayaraghavan K, Yun YS (2007) Utilization of fermentation waste (*Corynebacterium glutamicum*) for biosorption of reactive black 5 from aqueous solution. *J Hazard Mater* 141(1):45–52. <https://doi.org/10.1016/j.jhazmat.2006.06.081>
 36. Wang, K., Khoo, S. K., Kit, W. C., Anurita, S., Wei-Hsin, C., Jo-Shu, C., et al. (2021).

- Microalgae: The future supply house of biohydrogen and biogas. *Front. Energy Res.* 9. doi:10.3389/fenrg.2021.660399
37. Xiong J-Q (2019) Combined effects of sulfamethazine and sulfamethoxazole on a freshwater microalga, *Scenedesmus obliquus*: toxicity, biodegradation and metabolic fate. *J Hazard Mater* 370:138–146. <https://doi.org/10.1016/j.jhazmat.2018.07.049>
38. Y. Zhang, M.Y. Habteselassie, E.P. Resurreccion, V. Mantripragada, S. Peng, S. Bauer, L.M. Colosi, Evaluating removal of steroid estrogens by a model alga as a possible sustainability benefit of hypothetical integrated algae cultivation and wastewater treatment systems, *ACS Sustain. Chem. Eng.* 2 (2014) 2544e2553. <https://doi.org/10.1021/sc5004538>
39. Zeng, X., Danquah, M. K., Chen, X. D. and Yinghua, L. (2011). Microalgae bioengineering: From CO₂ fixation to biofuel production. *Renew. Sustain. Energy Rev.* 15, 3252–3260. doi:10.1016/j.rser.2011.04.014
40. Zhang, S. and Liu, Z. (2021). Advances in the biological fixation of carbon dioxide by microalgae. *J. Chem. Technol. Biotechnol.* 96 (6), 1475–1495. doi:10.1002/jctb. 6714
41. Zhao, B., & Su, Y. (2020). Macro assessment of microalgae-based CO₂ sequestration: Environmental and energy effects. *Algal Research*, 51, 102066. <https://doi.org/10.1016/j.algal.2020.102066>

