

Eco-Alchemy - Algae Cultivation and Its Application in Bioplastic Production

Vivek Pandya¹, Devanshi Trivedi², Kinjal Upadhyay³

How to cite this article:

Vivek Pandya, Devanshi Trivedi, Kinjal Upadhyay. Eco-Alchemy - Algae Cultivation and Its Application in Bioplastic Production. J Microbiol Relat Res. 2024;10(1):39–45.

Abstract

In this world of rapidly growing techs and steadfast alternatives which rather damage environment than make it flourishing, we need the actual green alternative which will efficiently supersede – the intoxication by petro fuels, waste-treatment machinery, green house effects and many more. This search of alternative has made its way to sleek end through the cultivation of algae and its benefits. As little amounts will do no good to the seeking benefits the major ways found by scientists soaring through early methods like dilution isolation method and use of stencil & gelatine to create patterns to more modern ways of open pond systems to photobioreactors. The review includes various efficient ways of algal cultivation such as use of photobioreactors, open raceways, two-phase hybrid systems and 3D triangular systems by discussing its limitations plus advantages. The photo bioreactors though providing more efficiency than open pond system they still are not much used. The greed for improved systems made discovery of the two-phase hybrid systems which obliterates disadvantage of both systems which are discussed. Though the thirst of improving led to the more different designs in the way of trials like 3D triangular systems which will be honoured in this review. We also go through a brief glimpse of algae's role in replacing synthetic plastics with its bioplastic production as it shows very versatile group of polymers and high degradability.

Keywords: Microalgae; Cultivation-methods; Open system; Closed system; Bioplastic.

INTRODUCTION

The world is now going towards green impact as by alternating the use of non-renewable and polluting sources with the more renewable and environmental friendly source. This involves biofuels like bioethanol, biohydrogen, bio-methane using many of the biological species like plants – i.e. their seed, leaves, fruits, etc. and microbes like yeast, *Zymomonasmobilis*, *Bacillus licheniformis* and algae.

Author Affiliation: ^{1,2}Student, ³Assistant Professor, Department of Biochemistry and Biotechnology, St. Xaviers College (Autonomous), Ahmedabad 380009, Gujarat, India.

Corresponding Author: Kinjal Upadhyay, Assistant Professor, Department of Biochemistry and Biotechnology, St. Xaviers College (Autonomous), Ahmedabad 380009, Gujarat, India.

E-mail: kinjal.upadhyay@sxca.edu.in

Received on: 21.01.2024 **Accepted on:** 19.04.2024



This work is licensed under a Creative Commons
Attribution-NonCommercial-ShareAlike 4.0.

Algae are the photosynthetic organisms that are multipurpose for many of the needs for extracting exemplified pigments and vitamins for supplements to treating the waste of industrial lethargy and providing gold standards of eco-fuels in the form of bio-ethanol and biohydrogen, While also using the environmental CO₂ released from petroleum burnt through vehicles in order to make organic carbon for its growth and in turn purifying the environment.

The Algae is divided into groups like green algae, brown algae, red algae, blue green algae, diatoms, golden algae, dinoflagellates, euglenophytes. Out of these mainly blue-green and green algae are used as they are very rich in vitamins and nutrients, also for biofuels too; Others like red and brown algae are majorly used for bioplastic production.

As such of importance are algae that it holds bills of hundreds of millions in dollar for researching its cultivation, application and products. Almost \$200 million were spent in researching and pilot project development by US department of energy (DOE). An investment such high as \$600 million is made by an oil company alone in oil production efforts from algae and related advertisements. This led to now blooming many of the experiments revolving around algae cultivation and its fruiting capacity of forming biofuels (Hall and Banemann, 2011)

Salt and wastewater tolerance being a spotlight of algae reduces freshwater use while sometimes wastewater can be considered a hosting nutrient resource in certain circumstances (Schenk *et al.*, 2008). The CO₂ sequestration being cleaning environment property of algae it will not only reduce environmental CO₂ but also produce the precious O₂ which accounts 70-80% of the oxygen breath by us (Schenk *et al.*, 2008; Hall, 2011).

There has been report by IMechE, 2009, which told increased CO₂ concentration will increase algae growth rate till other nutrients are in abundance (IMechE, 2009).

The cultivation of algae majorly be done via open system and closed system in which open systems consist of the raceway ponds and other totally open systems and closed systems includes photo-bioreactors.

Cultivation of algae

The algae harvesting is as old as at least 2500 years ago which have been utilized for food and medicine, yet the cultivation began only 300 years ago (Tseng, 1981). As early mentions reverting to

1950's stated their first roles in fuel production (Morton, 1998) and thus began the cultivation from lab to pilot scale.

In algae, they depend on strain specific cultivation criteria so as to increase the yield. Factors include temperature, mixing, hydrodynamics mass transfer, gas transfer, light cycle, water intensity, water quality, pH levels and many others like carbon nutrients availability, cell density/fragility/growth inhabitation. Thus, there has been more special interest lying in its cultivation system designs.

Algae can be grown in various designed systems from low on tech ponds to highly technical and controlled system of bioreactors. The designs firmly differ in terms of its economic requirements to environmental impact to factors in operating systems.

Major there are two systems that cultivation follows: 1) Open systems and 2) Closed systems

Open systems

The open ponds are basically the natural pits created with specific dimensions. They can either be simple ponds of desired diameter and depths whereas others called raceway ponds have desired diameter and depth plus circulating paddlewheels (Gundula Proksch, 2013).

Basically the depth of 15 to 20cm or 6 to 8 inches is ideal whereas diameter can vary (Gundula Proksch, 2013). At lab scale you can also use flasks, beakers and trays to cultivate this green gold so as they can behave as the mimics.

Advantages: (Schenk *et al.*, 2008)

- There is always a low construction cost.
- They idealise low economic margins.

Disadvantages: (Schenk *et al.*, 2008)

- Large land area.
- Low light utilization.
- High contamination.
- Weather changes.

Solutions: (Gundula Proksch, 2013)

These problems can be overcome by using a transparent or translucent barrier which can be covered on the area and thus can be turned into a greenhouse.

Closed system

The algae need around 1000-10,000 lux of light intensity of indirect sunlight to optimize

photosynthetic reaction as direct sunlight can cause photo inhibition and photo bleaching (Schenk *et*

al., 2008). This can insist that algae can be grown in 3D space instead of growing just on surface like




Fig. 1: Open system

Adopted from (Santos *et al.*, 2020)

normal plants (Gundula Proksch, 2013).

Vertical closed system increase productivity as the light will diffuse better. They are made of two components: A feeding vessel through which CO₂ and nutrients can be introduced while second is a solar array through which the mixed solution is pumped being exposed to sunlight (Gundula Proksch, 2013). The system increases yield and

contamination is eliminated (Gundula Proksch, 2013).

Ther low cost photobioreactor can be made using poly-ethylene bags which can be hanged through racks such that all sides of it can be exposed to the sunlight. Algae can be mixed into solution and pumped mechanically using series of bags. Systems like this can sometime require additional




Fig. 2: Closed systems for algal cultivation

adopted from (Masojídek & Torzillo, 2008)

construction in order to cut through the weather fluctuations (Kizililsoley and Helvacioğlu, 2008).

The 10 times more costly and producing 5 to 10 times higher yields of areal footprint than conventional methods is the closed photobioreactor. Here minimal volume of water is used for maximizing absorption of nutrients and energy under controlled environment. The surface-to-volume ratio provides light saturations and proper mixing ensures even CO₂ distribution and prevents algal cells from settling down. Glass tubes will efficiently be used to grow algae (Schenk *et al.*, 2008). A German company Algomed operates

world's largest photo-bioreactor in Klötze and this consists of 500km of glass tubes in about 12,000m² green house being able to produce up to 100t algal biomass per year (Gundula Proksch, 2013).

Two-stage Hybrid system

This system is developed as a result of research carried out to reduce disadvantages of both open pond system and photobioreactor and make a setup that falls somewhere in between having advantages of both (Olaizola, 2000; Schenk *et al.*, 2008; Su *et al.*, 2011). This system is made up of open raceways connected to the photobioreactor




Fig. 3: Two stage hybrid system
Adopted from (Aziz *et al.*, 2020)

and the flow happens from photobioreactor to open raceways pond (Narala *et al.*, 2016).

Comparison between the different systems

A *Tetraselmis* sp. M8 is used; The photobioreactor as well as the open raceway pond cultivation

results three main growth cycles whereas the fourth harvest seemed possible at 28th and 29th day. The harvest of only that biomass that is rich in lipids is done. If compared the Two-stage system, the hybrid system produced nearly six cycles and harvests. These results suggested that the average of growth




Fig. 4: Comparison between different systems
Data adopted from (Gundula Proksch, 2019)

was higher in the Hybrid system compared to both individual methods (Narala *et al.*, 2016).

This happened because hybrid system has provided biomass growth and lipid induction independently as it has two distinct stages. Hybrid system involved higher growth rates and low culture dormancy thanks to the different cultivation system for biomass production and lipid induction. This becomes advantageous as algae can either increase biomass or it can biosynthesize lipid at a single time (Lim *et al.*, 2012; Li *et al.*, 2014; Sharma *et al.*, 2014).

The major disadvantage mostly in open pond system, i.e. contaminating predators or other algae, is overcome by hybrid systems. The issue is solved here as the photobioreactor holds the algae in closed environment for a much of their growth and the open pond systems (prone to contamination) will gain the biomass only few days before harvesting (Moheimani and Borowitzka, 2006; Wang *et al.*, 2013).

Role of algae in bioplastic production

Plastics that are produced from biomass or environment friendly resources like food crops are known as bioplastic and have the same function as petroleum base bioplastic (Mekonnen *et al.*, 2013). Algae makes a preferable biomass source for the production of bioplastic, which has increased level of lipid content (Khoo *et al.*, 2020; Yew *et al.*, 2020). The biomass of algae consists of protein and carbohydrate-based polymers which can be used as one of the bioplastic constituents. Some of the examples of protein-based polymers are starch, cellulose, PHA (Polyhydroxyalkanoate, PHB (Polyhydroxy butyrate), PLA (Polylactic acid), PE (Polyethylene), PVC (Polyvinyl chloride) from algal biomass which are utilized to develop biodegradable plastics (Karan *et al.*, 2019). Among above mentioned protein-based polymers PHA is a polymer that is

most recommended for the production of bioplastic because it can be degraded by enzymatic action (Rasul *et al.*, 2017). Algal biomass can also be blended with other materials in order to prolonged life span, also to provide a better mechanical properties. The materials which can be blended with the algal biomass are petroleum plastics, natural products (cellulose or starch) or polymers, for the production of bioplastic (Rahman & Miller, 2017).

Table 1: Data represents bioplastic products generated by different microalgal species

Algal Species	Product	Bioplastic content (mg ml ⁻¹)
<i>Microcystis aeruginosa</i>	PHB	0.49±0.5
<i>Haematococcus pluvialis</i>	PHB	0.39±0.42
<i>Botryococcus brauni</i>	PHB	247±0.42
<i>Chroococcoides turgidus</i>	PHB	0.1

Data adopted from (Arora *et al.*, 2023)

As shown in Fig 5 the different type of blended materials are used for manufacturing of improved quality of bioplastic. There have been some researchers who manufactured the bioplastics (A. Pownall, 2020).

The few of the bioplastic manufacturing examples by researchers and designers are: Eric Klarenbeek and Maartje Dros, Dutch designers made an algae based bioplastic that can replace the currently used traditional plastics, also establishing lab called Algae Lab for algae cultivation and starch production which can act as raw material for bioplastic production; Austėja Platukyte made biodegradable product mixing agar and algae coated with calcium carbonate having potential to replace petroleum-based plastics; Ari Jonsson mixed water and dried red algae creating plastic bottles that can alternate traditional plastic bottles, these new plastic bottles can retain the form when full and easily biodegrade




Fig. 5: Bioplastic production

Adopted from (Chong *et al.*, 2022)

when empty; thus, the algae based bioplastic can be promising in replacing the traditional bioplastic very efficiently (Arora *et al.*, 2023).

CONCLUSION

Algae being a strong beneficiary can be grown in bulk amounts via the discussed methods out of which bioreactor are more efficient but less used while the hybrid system is very well negating limits of both these systems. Algae being good bioplastic source can also maintain eco-friendly approach.

Conflicts of Interest

There are no conflicts to declare.

Acknowledgements

The authors are thankful to St. Xavier's College (Autonomous), Ahmedabad, India for providing a financial assistance under the scheme of Research Project Seed Grant having award number SXCA/2021-2022/PR-01.

REFERENCES

1. A. Pownall, 10 bioplastic projects made from algae, corn starch and other natural materials. <https://www.dezeen.com/2018/10/09/bioplastic-projects-algae-corn-starch-beetle-shells/>, 2018. (Accessed 20 August 2020).
2. Arora, Y. K., Sharma, S., & Sharma, V. (2023). Microalgae in Bioplastic Production: A Comprehensive review. *Arabian Journal for Science and Engineering*, 48(6), 7225–7241. <https://doi.org/10.1007/s13369-023-07871-0>
3. Aziz, M. A., Kassim, K. A., Shokravi, Z., Jakarni, F. M., Liu, H., Zaini, N., Tan, L. S., Islam, A. B. M. S., & Shokravi, H. (2020). Two-stage cultivation strategy for simultaneous increases in growth rate and lipid content of microalgae: A review. *Renewable & Sustainable Energy Reviews*, 119, 109621. <https://doi.org/10.1016/j.rser.2019.109621>
4. Benemann, J. (1980). Overview: Algae oil to biofuels. ResearchGate. <https://www.researchgate.net/publication/229039925>
5. Chia, W. Y., Tang, D. Y. Y., Khoo, K. S., Lup, A. N. K., & Chew, K. W. (2020). Nature's fight against plastic pollution: Algae for plastic biodegradation and bioplastics production. *Environmental Science and Ecotechnology*, 4, 100065. <https://doi.org/10.1016/j.ese.2020.100065>
6. Chong, J. W. R., Tan, X., Khoo, K. S., Ng, H. S., Jonglertjunya, W., Yew, G. Y., & Show, P. L. (2022). Microalgae-based bioplastics: Future solution towards mitigation of plastic wastes. *Environmental Research*, 206, 112620. <https://doi.org/10.1016/j.envres.2021.112620>
7. Hall, J. (2011) The most important Organism? Ecology Global Network [online]. Ecology Communications Group, Inc. Available from: <http://www.ecology.com/> 2011/09/12/important-organism/[assessed Sep. 15, 2011]
8. IMechE Report, Institution of Mechanical Engineers (2009) Geo- Engineering Giving Us the Time to Act? Institution of Mechanical Engineers, London.
9. Karan, H., Funk, C., Grabert, M., Oey, M., & Hankamer, B. (2019). Green bioplastics as part of a circular bioeconomy. *Trends in Plant Science*, 24(3), 237–249. <https://doi.org/10.1016/j.tplants.2018.11.010>
10. Khoo, K. S., Chia, W. Y., Tang, D. Y. Y., Show, P. L., Chew, K. W., & Chen, W. (2020). Nanomaterials utilization in biomass for biofuel and bioenergy production. *Energies*, 13(4), 892. <https://doi.org/10.3390/en13040892>
11. Kizililsoley, M. & Helvacioğlu, S. (2008) Microalgae Growth Technology Systems. Soley Institute. Presentation at IIMSAM, New York, NY, 2008
12. Li, Y., Naghdi, F. G., Garg, S., Adarme-Vega, T. C., Thurecht, K. J., Ghafor, W. A., et al. (2014). A comparative study: the impact of different lipid extraction methods on current microalgal lipid research. *Microb. Cell Fact.* 13, 14. doi:10.1186/1475-2859-13-14
13. Lim, D. K., Garg, S., Timmins, M., Zhang, E. S., Thomas-Hall, S. R., Schuhmann, H., et al. (2012). Isolation and evaluation of oil-producing microalgae from sub-tropical coastal and brackish waters. *PLoS ONE* 7:e40751. doi:10.1371/journal.pone.0040751
14. Masojídek, J., & Torzillo, G. (2008). Mass cultivation of freshwater microalgae. In *Elsevier eBooks* (pp. 2226–2235). <https://doi.org/10.1016/b978-008045405-4.00830-2>
15. Mekonnen, T. H., Mussone, P., Khalil, H., & Bressler, D. C. (2013). Progress in bio-based plastics and plasticizing modifications. *Journal of Materials Chemistry. A, Materials for Energy and Sustainability*, 1(43), 13379. <https://doi.org/10.1039/c3ta12555f>
16. Moheimani, N. R., and Borowitzka, M. A. (2006). The long-term culture of coccolithophore *Pleurochrysis carterae* (Haptophyta) in outdoor raceway ponds. *J. Appl. Phycol.* 18, 703–712. doi:10.1007/s10811-006-9075-1
17. Morton, S. L. Modern Uses of Cultivated

- Algae. (1998) Ethnobotanical Leaflets [online]. Southern Illinois University Carbondale. Available from: <http://www.ethnoleaflets.com/leaflets/algae.htm> [assessed Sep. 3, 2011]
18. Narala, R. R., Garg, S., Sharma, K., Thomas-Hall, S. R., Deme, M., Li, Y., & Schenk, P. M. (2016). Comparison of microalgae cultivation in photobioreactor, open raceway pond, and a Two-Stage hybrid system. *Frontiers in Energy Research*, 4. <https://doi.org/10.3389/fenrg.2016.00029>
19. Olaizola, M. (2000). Commercial production of astaxanthin from *Haematococcus pluvialis* using 25,000-liter outdoor photo bioreactors. *J. Appl. Phycol.* 12, 499–506. doi:10.1023/A:1008159127672
20. Proksch, G. (2013). Growing Sustainability - Integrating Algae Cultivation into the Built Environment. Research Gate. <https://www.researchgate.net/publication/337292607>
21. Rahman, A., & Miller, C. D. (2017). Microalgae as a source of bioplastics. In *Elsevier eBooks* (pp. 121–138). <https://doi.org/10.1016/b978-0-444-63784-0.00006-0>
22. Rasul, I., Azeem, F., Siddique, M. H., Muzammil, S., Rasul, A., Munawar, A., Afzal, M., Ali, M., & Nadeem, H. (2017). Algae Biotechnology. In *Elsevier eBooks* (pp. 301–334). <https://doi.org/10.1016/b978-0-12-812360-7.00008-2>
23. Santos, M. G. B. D., Duarte, R. L., Maciel, A. M., Abreu, M., Reis, A., & De Mendonça, H. V. (2020). Microalgae biomass production for biofuels in Brazilian scenario: A critical review. *Bioenergy Research*, 14(1), 23–42. <https://doi.org/10.1007/s12155-020-10180-1>
24. Schenk, P. M., Thomas-Hall, S. R., Stephens, E., Marx, U. C., Mussgnug, J. H., Posten, C., Kruse, O. & Hankamer, B. (2008). Second generation biofuels: High-Efficiency microalgae for biodiesel production. *Bioenergy Research*, 1(1), 20–43. <https://doi.org/10.1007/s12155-008-9008-8>
25. Sharma, K., Li, Y., and Schenk, P. M. (2014). UV-C mediated lipid induction and settling, a step change towards economical microalgae biodiesel production. *Green Chem.* 16, 3539–3548. doi:10.1039/C4GC00552J
26. Su, C.-H., Chien, L.-J., Gomes, J., Lin, Y.-S., Yu, Y.-K., Liou, J.-S., et al. (2011). Factors affecting lipid accumulation by *Nannochloropsis oculata* in a two-stage cultivation process. *J. Appl. Phycol.* 23, 903–908. doi:10.1007/s10811-010-9609-4
27. Tseng, C. K. (1981) Commercial Cultivation. IN: Lobban, C. S. & Wynne, M. J. (Eds) The Biology of Seaweeds, Oxford: Blackwell Scientific Publication. 680-725.
28. Wang, H., Zhang, W., Chen, L., Wang, J., and Liu, T. (2013). The contamination and control of biological pollutants in mass cultivation of microalgae. *Bioresour. Technol.* 128, 745–750. doi:10.1016/j.biortech.2012.10.158.
29. Yew, G. Y., Khoo, K. S., Chia, W. Y., Ho, Y., Law, C. L., Leong, H. Y., & Show, P. L. (2020). A novel lipids recovery strategy for biofuels generation on microalgae *Chlorella* cultivation with waste molasses. *Journal of Water Process Engineering*, 38, 101665. <https://doi.org/10.1016/j.jwpe.2020.101665>.

