

ORIGINAL ARTICLE

Comprehensive Cataract Nursing Care Plan

Dinesh K.

HOW TO CITE THIS ARTICLE:

Dinesh K. Comprehensive Cataract Nursing Care Plan. *Ophthalmol Allied Sci.* 2025;11(1): 07–12.

ABSTRACT

Cataracts, a leading cause of visual impairment globally, necessitate surgical intervention as the primary treatment. A robust and patient-centered nursing care plan is paramount to ensuring optimal outcomes for individuals undergoing cataract surgery. This article outlines a comprehensive nursing care plan, encompassing pre-operative, intra-operative, and post-operative phases, grounded in the nursing process. It highlights key elements such as detailed assessment, appropriate nursing diagnoses, goal-setting, evidence-based interventions, and thorough evaluation. The aim is to enhance patient safety, comfort, education, and ultimately, facilitate a successful recovery and improved quality of life.

KEYWORDS

• Cataract • Nursing care • Ophthalmology • Surgery • Phacoemulsification • Lens replacement • Preoperative care • Postoperative care • Patient education • Complications management

INTRODUCTION

Cataracts, characterized by the clouding of the eye's natural lens, progressively impair vision and significantly impact an individual's daily activities and independence. Affecting millions worldwide, surgical removal of the opaque lens and implantation of an intraocular lens (IOL) is the most effective and common treatment. While the surgical procedure itself is often quick and minimally invasive, the patient's

journey extends far beyond the operating room. Comprehensive nursing care plays a pivotal role in every stage of this journey, from initial diagnosis and pre-operative preparation to post-operative recovery and long-term visual rehabilitation.¹

The complexity of managing patients undergoing cataract surgery is multifaceted, involving not only the physical aspects of the eye but also the psychological, social, and

AUTHOR'S AFFILIATION:

Ex-Staff Nurse, Kailash Hospital & Heart Institute, Sector 27, Noida, Uttar Pradesh 201301, India.

CORRESPONDING AUTHOR:

Dinesh K., Ex-Staff Nurse, Kailash Hospital & Heart Institute, Sector 27, Noida, Uttar Pradesh 201301, India..

E-mail: dineshrfp@gmail.com

➤ Received: 10.04.2025 ➤ Accepted: 22.05.2025



Creative commons non-commercial CC BY-NC: This article is distributed under the terms of the creative commons attribution non-commercial 4.0 License (<http://www.creativecommons.org/licenses/by-nc/4.0/>) which permits non-commercial use, reproduction and distribution of the work without further permission provided the original work is attributed as specified on the Red Flower Publication and Open Access pages (<https://rfppl.co.in>)

educational needs of the individual. Nurses, as primary caregivers and patient advocates, are uniquely positioned to address these holistic needs. A well-structured nursing care plan serves as a roadmap, ensuring systematic, individualized, and high-quality care delivery. This article aims to delineate a detailed nursing care plan for patients undergoing cataract surgery, emphasizing the critical components that contribute to successful outcomes and patient satisfaction.²

METHODS AND MATERIALS

Framework for Care Plan Development

The development of a comprehensive nursing care plan for cataract patients is rooted in the systematic application of the nursing process: Assessment, Diagnosis, Planning, Implementation, and Evaluation. This framework ensures a holistic, individualized, and goal-oriented approach to patient care.^{3,4}

1. Assessment

This initial phase involves gathering comprehensive data about the patient's health status, focusing on factors relevant to cataract surgery and recovery.

- **Pre-operative Assessment:**
 - **Medical History:** Review of past medical conditions (e.g., hypertension, diabetes, cardiac issues), previous surgeries, allergies, and current medications (especially anticoagulants).
 - **Ocular History:** History of other eye conditions (e.g., glaucoma, macular degeneration), previous eye surgeries, and use of eye drops.
 - **Visual Acuity:** Baseline visual assessment (Snellen chart, near vision) to quantify the extent of impairment.
 - **Functional Status:** Assessment of ability to perform activities of daily living (ADLs), impact of vision loss on lifestyle, and assistive devices used.
 - **Psychological Status:** Assessment of anxiety levels, fears, coping mechanisms, and expectations regarding the surgery and outcomes.
 - **Social Support:** Identification of

family or caregiver availability for assistance post-operatively.

- **Knowledge Level:** Patient's understanding of the procedure, pre-operative instructions, post-operative care, and potential complications.
- **Intra-operative Assessment:** Brief but crucial monitoring of vital signs, patient comfort, and adherence to sterile technique.
- **Post-operative Assessment:** Immediate and ongoing assessment of vital signs, pain level, nausea, eye dressing, drainage, redness, swelling, and patient's ability to follow instructions.

2. Diagnosis

Based on the comprehensive assessment, specific nursing diagnoses (using NANDA-I terminology) are formulated to identify actual or potential health problems amenable to nursing intervention.

Common diagnoses for cataract patients include:

- **Disturbed Sensory Perception:** Visual related to lens opacity and surgical intervention.
- **Anxiety related to surgical procedure,** potential outcomes, or loss of vision.
- **Deficient Knowledge related to surgical procedure,** pre-operative preparation, and post-operative care.
- **Risk for Injury related to impaired vision,** altered balance, or post-operative restrictions.
- **Acute Pain related to surgical incision or post-operative inflammation.**
- **Risk for Infection related to compromised primary defenses (surgical incision).**

3. Planning

This stage involves establishing patient-centered, measurable, achievable, relevant, and time-bound (SMART) goals and desired outcomes. It also includes developing individualized nursing interventions to achieve these goals.

- **Examples of Goals:**
 - Patient will demonstrate understanding of pre-operative instructions by (date/time).

- Patient will report anxiety level as manageable or reduced by discharge.
- Patient will verbalize understanding of post-operative care instructions before discharge.
- Patient will remain free from signs of infection post-operatively.
- Patient will report pain level as acceptable (e.g., <3/10) within 24 hours post-surgery.

4. Implementation

This phase involves carrying out the planned nursing interventions. This includes direct care, patient education, medication administration, monitoring, and coordination of care.

- **Pre-operative Interventions:** Patient education, medication instructions, NPO status, vital signs, pre-op drops.
- **Intra-operative Interventions:** Maintaining sterile field, monitoring patient, providing comfort.
- **Post-operative Interventions:** Pain management, vital sign monitoring, eye care, discharge teaching.

5. Evaluation

The final stage involves continuously evaluating the effectiveness of the nursing care plan and the patient's progress toward achieving the established goals. This may lead to modifications in the care plan as needed.

- **Examples of Evaluation:**
 - Did the patient accurately describe the post-operative eye drop regimen?
 - Was the patient's pain effectively managed?
 - Were there any signs of infection?
 - Did the patient achieve improved visual acuity as expected?

The "materials" for this framework largely consist of standardized assessment forms, patient education leaflets, visual aids, medication administration records, and electronic health record (EHR) systems that facilitate seamless documentation and communication.⁵⁻⁷

RESULTS

Expected Outcomes of the Care Plan

The diligent application of a comprehensive cataract nursing care plan yields numerous positive outcomes for the patient. These results contribute significantly to the overall success of the surgical intervention and the patient's quality of life.

- **Improved Visual Acuity:** The primary expected outcome, allowing patients to regain functional vision and participate more fully in daily activities.
- **Reduced Anxiety and Fear:** Effective pre-operative education and psychological support help alleviate patient apprehension, leading to a calmer surgical experience.
- **Effective Pain Management:** Appropriate pain assessment and interventions ensure patient comfort post-operatively, facilitating quicker recovery.
- **Prevention of Complications:** Adherence to sterile techniques, proper eye care, and timely identification of warning signs minimize risks such as infection, hemorrhage, or elevated intraocular pressure.
- **Enhanced Patient Understanding and Adherence:** Comprehensive teaching empowers patients to actively participate in their recovery, correctly administer eye drops, and adhere to activity restrictions.
- **Timely and Safe Discharge:** A well-planned discharge ensures patients are ready to return home with proper support and understanding of their continued care needs.
- **Optimal Long-Term Visual Rehabilitation:** By providing ongoing education and facilitating follow-up care, the care plan supports sustained visual health and adaptation to improved vision.
- **Increased Patient Satisfaction:** A holistic and supportive nursing approach significantly enhances the patient's overall experience and satisfaction with their care.

DISCUSSION

The comprehensive cataract nursing care plan is a dynamic and patient centered tool that guides nurses in delivering high quality, individualized care throughout the perioperative period. Each phase pre-operative, intra-operative, and post-operative presents unique challenges and opportunities for nursing intervention.⁹

Pre-operative Phase: Foundations of Success

This phase is critical for laying the groundwork for a successful surgical experience.

- **Education:** Detailed patient education is paramount. Nurses must explain the surgical procedure (phacoemulsification, IOL implantation), expected sensations during and after surgery, the importance of pre-operative eye drops (e.g., antibiotics, anti-inflammatory), and NPO status guidelines. Using visual aids, simple language, and confirming comprehension are essential.
- **Psychological Support:** Many patients experience anxiety. Nurses should provide reassurance, address fears, and clarify misconceptions. Identifying support systems (family, friends) is vital, as these individuals also need to understand their role in post-operative care.
- **Medication Reconciliation:** A thorough review of all medications, especially anticoagulants (which may need temporary discontinuation per physician's orders) and medications for co-morbidities like diabetes or hypertension, is crucial for patient safety.

Intra-operative Phase: Vigilance and Coordination

While the patient is under the direct care of the surgical team, the circulating nurse plays a critical role in safety and support.

- Maintaining a sterile environment is fundamental.
- Monitoring the patient's vital signs and level of consciousness, and reporting any changes immediately to the surgical team.
- Ensuring patient comfort through proper positioning and padding.

- Providing gentle reassurance to the patient if local anesthesia is used.

Post-operative Phase: Recovery and Rehabilitation

This is the most intensive phase for nursing care, focusing on immediate recovery and preparing the patient for home care.

- **Immediate Post-operative Care:** Monitoring vital signs, assessing pain, checking for nausea/vomiting, and observing the operative eye for drainage, excessive redness, or swelling. The application of an eye shield or patch as prescribed is important for protection.
- **Pain Management:** Administering prescribed analgesics and assessing their effectiveness. Non-pharmacological interventions like a cool compress may also be useful.
- **Discharge Teaching:** This is arguably the most vital post-operative nursing intervention. It must be comprehensive and tailored to the patient's learning capacity and home environment. Key teaching points include:
 - **Eye Drop Instillation:** Detailed instructions and demonstration, often requiring a return demonstration from the patient or caregiver. A written schedule is invaluable.
 - **Activity Restrictions:** Avoiding heavy lifting, bending at the waist, strenuous activities, rubbing the eye, and swimming. Explaining the rationale behind these restrictions fosters compliance.
 - **Signs of Complications:** Educating patients on warning signs such as severe pain, sudden vision loss, increasing redness, discharge, or flashes of light, and when to contact the ophthalmologist.
 - **Eye Protection:** Emphasizing the use of eye shields at night and sunglasses during the day.
 - **Follow-up Appointments:** Ensuring the patient understands the importance and schedule of post-operative visits.

SAMPLE NURSING CARE PLAN TABLE

A structured approach using a care plan table enhances consistency and clarity in care delivery

NANDA Nursing Diagnosis	Desired Outcome(s)	Nursing Interventions	Rationale	Evaluation
Disturbed Sensory Perception: Visual related to lens opacity and surgical procedure.	Patient will report improved vision and reduced visual distortions within 24-48 hours post-surgery. Patient will ambulate safely with assistance if needed.	1. Assess baseline visual acuity and peripheral vision. 2. Orient patient to surroundings and available assistive devices (e.g., call bell, handrails). 3. Encourage patient to use prescribed eye drops correctly. 4. Instruct patient to avoid rubbing/ touching the operative eye. 5. Emphasize importance of wearing eye shield/sunglasses as directed. 6. Provide clear, concise instructions on activity restrictions (e.g., no heavy lifting, bending). 7. Assist with ambulation and ADLs as needed, ensuring a clutter-free environment.	1. Establishes a baseline for comparison and identifies degree of impairment. 2. Promotes safety and reduces anxiety in an unfamiliar environment. 3. Reduces inflammation and prevents infection, supporting clear vision. 4. Prevents damage to the surgical site and potential for infection. 5. Protects the eye from accidental trauma, dust, and UV light, which can impede healing. 6. Prevents increased intraocular pressure (IOP) and stress on the surgical incision, which could lead to complications. 7. Minimizes risk of falls and injury, especially in the immediate post-operative period when vision may still be adapting.	Patient reports clearer vision and no significant visual distortions. Patient ambulates without falls and articulates understanding of safety precautions.
Anxiety related to surgical procedure, fear of unknown, and potential loss of vision.	Patient will verbalize reduced anxiety and express confidence in the care plan by discharge. Patient will demonstrate relaxed body language.	1. Assess patient's level of anxiety using a subjective scale (e.g., 0-10). 2. Provide clear, concise, and accurate information about the procedure and expected outcomes. 3. Address patient's specific fears and misconceptions. 4. Encourage questions and active participation in care planning. 5. Teach relaxation techniques (e.g., deep breathing, guided imagery). 6. Ensure consistent communication from the healthcare team. 7. Involve family/support persons as appropriate.	1. Provides a quantifiable measure of anxiety for evaluation and intervention. 2. Knowledge reduces fear of the unknown and fosters a sense of control. 3. Debunking myths and providing factual information can significantly alleviate anxiety. 4. Empowerment and involvement can reduce feelings of helplessness. 5. Promotes a sense of calm and helps manage physiological manifestations of anxiety. 6. Builds trust and confidence in the care team. 7. Support from loved ones provides comfort and reinforces coping.	Patient verbalizes feeling calmer and less anxious. Patient actively participates in discharge planning and asks relevant questions.
Deficient Knowledge related to post-operative care and medication regimen.	Patient will accurately describe post-operative care instructions and demonstrate correct eye drop instillation technique before discharge.	1. Assess current knowledge level regarding post-operative care. 2. Provide written and verbal instructions on eye care, activity restrictions, and signs of complications. 3. Demonstrate correct eye drop instillation, and request a return demonstration from patient/caregiver. 4. Provide a detailed schedule for eye drop administration. 5. Emphasize the importance of follow-up appointments. 6. Use teach-back method to confirm understanding. 7. Provide contact information for questions/concerns.	1. Identifies gaps in knowledge to tailor education appropriately. 2. Reinforces learning and provides a reference for home care. 3. Ensures proper medication delivery and reinforces learning through active participation. 4. Reduces confusion and promotes adherence to the prescribed regimen. 5. Crucial for monitoring healing, detecting complications, and assessing long-term outcomes. 6. Verifies comprehension and identifies areas needing further clarification. 7. Provides a resource for ongoing support and reduces anxiety about unknown post-discharge issues.	Patient correctly explains all post-operative instructions and demonstrates proper eye drop instillation. Patient identifies signs of complications and states when to contact the doctor.

Challenges and Interdisciplinary Collaboration

While the framework is robust, challenges exist. Patients with cognitive impairments, multiple co-morbidities (e.g., severe arthritis limiting eye drop instillation), language barriers, or limited social support require adapted care plans. Nurses often coordinate with social workers, physical therapists, and family members to ensure a safe transition home. Interdisciplinary collaboration with ophthalmologists, anesthesiologists, and pharmacists is continuous, ensuring that the nursing care plan aligns with medical orders and patient needs. The rise of telehealth and digital health tools also presents new avenues for patient education and post-operative monitoring, offering opportunities to enhance the reach and effectiveness of nursing care. Continuous quality improvement initiatives are vital to refine care pathways and incorporate new evidence-based practices.¹⁰⁻¹²

CONCLUSION

The development and implementation of a comprehensive cataract nursing care plan are integral to achieving superior patient outcomes and enhancing the quality of life for individuals undergoing cataract surgery. By systematically applying the nursing process assessment, diagnosis, planning, implementation, and evaluation nurses can deliver individualized, evidence-based care that addresses not only the physical aspects of the surgery but also the psychological, educational, and social needs of the patient. The nurse's role as educator, advocate, and caregiver is indispensable in guiding the patient through the perioperative journey, ensuring safety, managing symptoms, preventing complications, and empowering patients with the knowledge needed for effective self-care. As healthcare evolves, the nursing care plan remains a cornerstone of professional practice, continuously adapting to new technologies and patient demographics to deliver optimal visual rehabilitation and sustained well-being.

REFERENCES

1. Smeltzer, S. C., Bare, B. G., Hinkle, J. L., & Cheever, K. H. (2018). *Brunner & Suddarth's Textbook of Medical-Surgical Nursing* (14th ed.). Wolters Kluwer.
2. Ackley, B. J., Ladwig, G. B., & Makic, M. B. F. (2017). *Nursing Diagnosis Handbook: An Evidence-Based Guide to Planning Care* (11th ed.). Elsevier.
3. American Academy of Ophthalmology. (2023). *Cataract: Patient Information*. Retrieved from www.aao.org.
4. Ignatavicius, D. D., Workman, M. L., & Rebar, C. R. (2021). *Medical-Surgical Nursing: Concepts for Interprofessional Collaborative Care* (10th ed.). Elsevier.
5. World Health Organization. (2023). *Blindness and Vision Impairment*. Retrieved from www.who.int.
6. Aliancy JF, Mamalis N. Crystalline Lens and Cataract. In: Kolb H, Fernandez E, Jones B, Nelson R, editors. *Webvision: The Organization of the Retina and Visual System* [Internet]. University of Utah Health Sciences Center; Salt Lake City (UT): Aug 15, 2017. [PubMed]
7. Ermiş SS, Öztürk F, Inan UU. Comparing the efficacy and safety of phacoemulsification in white mature and other types of senile cataracts. *Br J Ophthalmol*. 2003 Nov;87(11):1356-9. [PMC free article] [PubMed]
8. Guan JY, Ma YC, Zhu YT, Xie LL, Aizezi M, Zhuo YH, Wumaier A. Lens nucleus dislocation in hypermature cataract: Case report and literature review. *Medicine (Baltimore)*. 2022 Sep 02;101(35):e30428. [PMC free article] [PubMed]
9. Ono T, Miyata K. Morgagnian Cataract. *JMA J*. 2022 Jul 15;5(3):379-380. [PMC free article] [PubMed]
10. Rong X, Rao J, Li D, Jing Q, Lu Y, Ji Y. TRIM69 inhibits cataractogenesis by negatively regulating p53. *Redox Biol*. 2019 Apr;22:101157. [PMC free article] [PubMed]
11. Vlastra W, Claessen BE, Beijl MA, Sjaauw KD, Streekstra GJ, Wykrzykowska JJ, Vis MM, Koch KT, de Winter RJ, Piek JJ, Henriques JPS, Delewi R. Cardiology fellows-in-training are exposed to relatively high levels of radiation in the cath lab compared with staff interventional cardiologists-insights from the RECAP trial. *Neth Heart J*. 2019 Jun;27(6):330-333. [PMC free article] [PubMed]
12. Sugawa H, Matsuda S, Shirakawa JI, Kabata K, Nagai R. [Preventive Effects of Aphanothece sacrum on Diabetic Cataracts]. *Yakugaku Zasshi*. 2019;139(3):381-384. [PubMed]