

## Comparative Evaluation of two Different Intravenous Doses of Midazolam to Aid the Insertion of LMA Classic as Adjuvants to Propofol Anaesthesia

Geetanjali Pushkarna<sup>1</sup>, Varun Pushkarna<sup>2</sup>, Preetveen Kaur<sup>3</sup>, Pankaj Sarangal<sup>4</sup>

<sup>1,3,4</sup>Assistant Professor, Department of Anaesthesia, Sri Guru Ram Das Institute of Medical Sciences and Research, Amritsar, Punjab Punjab 143501, India. <sup>2</sup>Consultant Critical Care, Fortis Escorts Hospital, Amritsar, Punjab 143004, India.

### Abstract

**Background:** Propofol is a widely accepted medication for the successful insertion of laryngeal mask airway, but as a sole induction agent, it has a very low success rate. A number of co-induction agents have been used with propofol to increase the success rate of LMA insertion. **Aim and Objectives:** To evaluate the efficacy of two different doses of midazolam i.e. 0.05 mg/kg and 0.08 mg/kg intravenously when used with propofol versus propofol alone for LMA insertion. To find the appropriate dose of midazolam that provides ideal condition and maximum haemodynamic stability. **Material & Methods:** 75 adult patients of ASA I & II grade randomly divided into 3 groups. Group P: Propofol + Saline Group PM1: Midazolam 0.05 mg/kg 3 minutes before propofol + Propofol Group PM2: Midazolam 0.08 mg/kg 3 minutes before propofol + Propofol. LMA insertion attempted 1 minute after the administration of Propofol. Total dose of propofol used, the insertion conditions and haemodynamic changes were noted. **Results:** There was no statistical difference in demographic profile. Haemodynamically patients were more stable in group PM1 & PM2 than in group P. Dose of propofol used in PM2 (2.05+0.21 in mg/kg) & PM1 (2.38+0.21) was less than in Group P (2.84 mg/kg). 100% success rate in LMA insertion was observed in group PM1 & PM2 in first attempt whereas it was 80% in group P. 88% in group PM2, 60% in group PM1 & only 32% patients in group P had excellent insertion conditions. **Conclusion:** Propofol as a sole agent does not provide LMA insertion conditions. Midazolam when used with propofol provides adequate conditions for LMA insertion in a dose dependent manner.

**Keyword:** Laryngeal mask airway; Midazolam; Propofol.

### How to cite this article:

Geetanjali Pushkarna, Varun Pushkarna, Preetveen Kaur et al. Comparative Evaluation of two Different Intravenous Doses of Midazolam to Aid the Insertion of LMA Classic as Adjuvants to Propofol Anaesthesia. Indian J Anesth Analg. 2019;6(3):697-704.

### Introduction

The two fundamental requirements of general anaesthesia are to maintain the patency of the airway and to ensure adequate ventilation. Endotracheal intubation, the gold standard for securing the

airway, being an invasive measure, is associated with greater haemodynamic alterations [1]. Laryngeal Mask Airway (LMA) serves as a bridge between the facemask and endotracheal tube. It is a good alternative to intubation in short duration procedures, being less invasive and less stimulating, thus, Preventing stress response [2].

**Corresponding Author:** Geetanjali Pushkarna, Assistant professor in Department of Anaesthesia, Sri Guru Ram Das Institute of Medical Sciences and Research, Amritsar, Punjab 143501, India.

**E-mail:** gitzgitz@gmail.com

**Received on** 06.03.2018, **Accepted on** 04.04.2019

However, airway reflexes have to be obtunded by general/topical anaesthesia or muscle relaxants before LMA insertion to prevent the patient movements, coughing, gagging and laryngeal spasm [2,3]. Hence, an adequate depth of anaesthesia and muscle relaxation have to be ensured. Propofol is the best known induction agent for LMA insertion, as it effectively suppresses the oro-pharyngeal reflexes, cough reflex and sensitivity of upper airway [4]. When used as a sole induction agent, much larger doses are required to achieve the desired end points, which can precipitate hypotension, respiratory depression and delayed awakening [3].

Many drugs such as- opioids [5,6,7], clonidine [11], midazolam [8,9], dexmedetomidine, lignocaine [12] and succinylcholine [9,10] have been employed as co-induction agents to decrease the induction dose of propofol, so as to get a better success rate, allowing early recovery with minimal side effects. Though, an adjuvant is yet to be found [4]. Opioids are the most commonly used adjuvants with propofol to aid LMA insertion, but respiratory depression, apnoea and chest rigidity are commonly encountered side effects [5]. Benzodiazepines, like midazolam, when given intravenously produce significant depression of the upper airway sensitivity but without haemodynamic instability and other side effects of opioids. Midazolam also reduces the induction dose of propofol, via a synergistic action and improves the LMA insertion conditions than propofol alone [5].

In this study, we evaluated the effect of different doses of Midazolam premedication i.e 0.05 and 0.08 mg/kg I/V with propofol versus propofol alone for LMA insertion and associated Haemodynamic changes before, during and after LMA insertion. The primary objective was to find out the appropriate dose of midazolam that provides the ideal condition for LMA insertion with maximum haemodynamic stability when used along with propofol.

## Material and Methods

After obtaining the approval from institutional ethics committee, the present trial was conducted on 75 patients, aged 20-60 years, of either sex, belonging to ASA grade I and II, undergoing elective surgical procedures of short duration. Informed consent was obtained in written from all the patients after explaining the procedure and were divided into three groups of 25 each by computer generated random number slips. Unwilling uncooperative

patients, patients at risk of aspiration (hiatus hernia, pregnancy, full stomach, intestinal obstruction etc.), patients with low pulmonary compliance- obesity, patients with pre op sore throat and URI, patients with oral pathology and patients allergic to the study drug were excluded from the study.

Patients were kept NPO for at least 6 hours prior to surgery. Tablet pantoprazole 40 mg was given orally to all the patients a night before surgery. The study drug solution was prepared with normal saline upto total volume of 5 ml by an anaesthesiologist not involved in this study and dispensed in unlabelled syringe. Group P was given Propofol and normal saline (control group). Group PM<sub>1</sub> was given Midazolam 0.05 mg/kg with propofol and group PM<sub>2</sub> received midazolam 0.08mg/kg IV with propofol IV. Electrocardiogram, NIBP and pulse oximetry were used for monitoring and mean arterial blood pressure, oxygen saturation and heart rate were recorded pre-induction, after induction of anaesthesia and after laryngeal mask insertion. Injection glycopyrrolate 0.2 mg intravenously was given before induction. After pre-oxygenation with 100% oxygen for 3 minutes, the patients were given their assigned drug solution over 10 seconds by an observer not involved in the study. 3 minutes after giving the study drug, the initial dose of Propofol was injected with IV at a constant rate over 30 seconds. Desired end points chosen to assess the adequacy of anesthesia were the loss of response to verbal commands and loss of eyelash reflex. If required, further boluses of propofol 0.2 mg/kg IV were given every 15 seconds and the total dose of propofol was calculated.

LMA insertion was attempted 30 seconds after the loss of eyelash reflex in all the patients using the technique described in Intravent laryngeal mask instruction manual. Jaw relaxation before LMA insertion was assessed and graded as (according to Young's criteria [13]):

Grade I: Absolutely relaxed jaw with no muscle tone.

Grade II: Moderately relaxed jaw with some degree of muscle tone.

Grade III: Poorly relaxed jaw with full muscle tone.

Incidence of gagging and coughing [14] was also graded on a 4 point scale with Grade I being no gagging / coughing and Grade IV being the most severe.

Overall insertion conditions were assessed according to the modified scheme of Lund and Stovener [15] as- Excellent, Good, Poor or

Unacceptable. If failed on first attempt, additional bolus dose of propofol 0.5 mg/kg was given. Insertion of LMA was attempted to a maximum of three attempts and patients were intubated if third attempt did not succeed. After securely positioning the LMA, patients were kept on spontaneous respiration using isoflurane and 66% N<sub>2</sub>O with O<sub>2</sub>. No further data was collected. Only the patients with first successful attempt of LMA insertion were taken for comparing the haemodynamic changes and LMA insertion conditions.

*Statistical Analysis:* The data from the present study was systematically collected, compiled and statistically analyzed to draw conclusions. Chi square test, student t test and ANOVA test were

applied and p value < 0.05 was taken as significant.

### Results

Demographic data, such as age, weight and duration of surgery were comparable in all the three groups (p>0.05) [Table 1]. Total induction dose of propofol was significantly reduced in group PM<sub>1</sub> and PM<sub>2</sub> as compared group P (p<0.001), with group PM<sub>2</sub> having the least requirement (p< 0.001) [Table 2], [Figure 1]. In group PM1, total dose was 2.38 ± 0.21 mg /kg while in group PM2 it was 2.05 ± 0.21 mg/kg [Table 2], [Figure 2]. Increasing the dose of midazolam further decreased the propofol requirement. There was 16% reduction in propofol

**Table 1:** Demographic Data:

Group	Age (in years)	Weight (in kg)	Duration of surgery (in minutes)
I	31.56 ± 8.47	54.88 ± 6.38	26.40 ± 6.21
II	34.56 ± 11.53	56.20 ± 4.86	27.40 ± 5.02
III	29.88 ± 11.94	56.28 ± 4.78	30.00 ± 6.12

**Table 2:** Dose of Propofol used in mg:


	P	PM1	PM2
Total Dose of propofol (mg)	155.8 ± 17.54	132.20 ± 15.28	109.80 ± 24.56
Dose in Propofol (mg/kg)	2.84 ± 0.14	2.38 ± 0.21	2.05 ± 0.21

**Table 3:** Jaw relaxation:

Group	p value
I & II	p<0.05
I & II	p<0.001
II & III	p<0.01

**Table 4:** Incidence of gagging and coughing

Group	p value
I & II	p<0.05
I & III	p<0.001
II & III	p<0.001



**Fig. 1:**

dose required in group PM1 and 27% reduction in group PM2. LMA was successfully inserted in the first attempt in 100% patients in group PM<sub>1</sub> and PM<sub>2</sub>, whereas More than one attempt were required in group P [Figure 3].

Ten (10)% patients in group PM<sub>2</sub> had absolute grade-I jaw relaxation while 70% patients in group PM<sub>1</sub> and only 40% group P had the absolute jaw relaxation [Figure 4]. This difference in all the three

groups was highly significant ( $p < 0.001$ ) [Table 3].

The number of patients with no gagging, coughing or movements was significantly higher in group PM<sub>2</sub> (88%) and PM<sub>1</sub> (60%) as compared to group P (32%) [Table 4]. 24% patients in group P experienced severe gagging, while it was not seen in other two groups [Figure 5]. The overall insertion conditions were graded excellent in 88% patients in group PM<sub>2</sub>, 60% patients in group PM<sub>1</sub>, with a




Fig. 2:




Fig. 3:

highly significant ( $p < 0.001$ ) difference. Group P had the least optimal conditions for insertion in only 32% patients, with highly significant difference from other two groups [Figure 6].

A significant fall in mean arterial pressure was seen in all the three groups after induction.

However, maximum decrease occurred in group P as compared to group PM<sub>1</sub>, which was more than in group PM<sub>2</sub> [Figure 7]. Post induction increase in heart rate was observed in all the three groups but it increased only minimally in group III [Figure 8]. A significant reduction in SpO<sub>2</sub> trends was seen after LMA insertion in group P but it remained




Fig. 4:




Fig. 5:




Fig. 6:




Fig. 7:




Fig. 8:




Fig. 9:

near pre induction levels in group PM<sub>1</sub> and PM<sub>2</sub> [Figure 9]. Successful LMA insertion in first attempt was seen in 100% patients in group PM<sub>1</sub> & PM<sub>2</sub> whereas in group P, it was successful in 80% patients in first attempt.

## Discussion

Use of LMA has gained wide acceptance for routine airway management, difficult airways and in emergency situations, owing to its ease of insertion. LMA is well tolerated at lower anaesthetic concentrations than the endotracheal tube, which allows earlier emergence from anaesthesia, making it a suitable option for day care procedures [16]. LMA insertion does not require neuromuscular blockade provided the depth of anaesthesia is adequate with obtundation of airway reflexes. Although, propofol is an ideal induction agent for LMA insertion [1], but when used alone larger doses are required adversely causing cardio-respiratory depression [3].

In the present study, the mean dose of propofol required for LMA insertion was found to be  $2.84 \pm 0.14$  mg/kg with the incidence of successful insertion of LMA in first attempt was 80% patients. Excellent insertion conditions were found in 32% patients. Jain Namrata et al. [18] found that 3.1 mg/kg of propofol was required with the incidence of successful insertion in first attempt in only 50% patients with side effects like coughing, gagging, head/limb movements, hypotension and prolonged apnoea. Wafaa TS et al. [9] too found that with the dose of 3.0 mg/kg propofol, successful insertion was seen only in 60% patients with significant fall in MAP and increase in Heart Rate in post induction period with excellent insertion conditions in 20% patients only. To overcome these limitations of propofol, it is being used in combination with drugs like benzodiazepines [8,9], opioids [5,6,7], intravenous or topical lignocaine [12] and muscle relaxants [9,10].

Benzodiazepines are known to reduce upper airway reflexes, besides providing excellent anxiolytic, sedative and amnesic properties. Midazolam is being extensively used in operating room because of its rapid onset of action and brief half life. When used as a premedication, midazolam is known to decrease the dose of propofol and the incidence of adverse haemodynamic effects [19,20,21]. Further addition of midazolam to propofol improves the insertion conditions for LMA [9,18]. In the present study, we compared the

conditions for LMA insertion and haemodynamic changes using propofol alone and in combination with two different doses of midazolam- 0.05 mg/kg I/V and 0.08 mg /kg I/V.

Dose of propofol decreased by 16% in group PM<sub>1</sub> and by 27% in group PM<sub>2</sub>. Several authors have published similar results. Short and Chui [19] suggested that the dose of propofol was reduced to 52% when midazolam was used with propofol. Bhasker, et al. [22] also found that the dose of propofol was decreased with use of midazolam.

Absolute jaw relaxation was observed in 100% patients in group PM<sub>2</sub> while 76% patients in PM<sub>1</sub> and 48% in group P had absolute jaw relaxation. None of the patients in group PM<sub>1</sub> and PM<sub>2</sub> had grade 4 gagging while 24% experienced it in group P. The incidence of gagging and coughing was even lower in PM<sub>2</sub> than PM<sub>1</sub> (40%). The insertion conditions were the best in group PM<sub>2</sub> where midazolam was used in Dose of 0.08mg/kg with 83% patients in Group PM<sub>2</sub>, 60% in group PM<sub>1</sub> and only 32% in group P had excellent insertion conditions. Similar findings were reported by Dhamotharan et al. [22]. They found that LMA insertion was easy in 80% patients where midazolam 0.05 mg/kg was used with in combination with propofol however it was 33.33% in patients where propofol was used alone. Salem [9] too found that LMA was successfully inserted in first attempt in 95% patients and insertion conditions were excellent in 100% patients in propofol-midazolam group.

The patients receiving midazolam propofol combination could have been more deeply anaesthetized or more likely this combination could depress airway reflexes to greater degree.

We observed that increasing the dose of midazolam from 0.05 mg /kg to 0.08 mg /kg further improved the conditions for LMA insertion. This could be attributed to dose dependent depression of upper airway reflexes by midazolam. MAP in all three groups decreased after induction from the pre-induction levels with 7% decrease in group P, 3% in PM<sub>1</sub> and 2.8% in PM<sub>2</sub> respectively. After LMA insertion, MAP was lower in all the groups but this decrease was comparable in group P and PM<sub>1</sub> whereas in group PM<sub>2</sub> the fall was not much. Heart rate increased in post induction and post insertion period in all the groups. This increase was max in group P. Group PM<sub>1</sub> had modest increase in heart rate while in group PM<sub>2</sub> there was least variation. There was a trend towards decrease in saturation after LMA insertion in group P but it remained stable in group PM<sub>1</sub> and PM<sub>2</sub>. Our study shows that

the use of midazolam with propofol results in better hemodynamic stability than when propofol is used alone to insert the LMA. This finding is consistent with previous studies where midazolam has been shown to improve haemodynamic stability when used with propofol [9,18,20].

### Conclusion

Propofol when used alone does not provide ideal conditions for LMA insertion. The use of midazolam as co-induction agent with propofol provides excellent insertion conditions with Stable Haemodynamics in a dose dependent manner. It also decreases the incidence of side effects like coughing, gagging and laryngospasm.

### References

- Dorsch JA, Dorsch SE. Face Mask and Airways In Understanding anesthetic Equipment. 3<sup>rd</sup> ed. Baltimore: William and Wilkins; 1994.
- Jarineshin H, Kashani S, Vatankeh M, AAbdulazade Baghaee AA, Sattari S, Fekrat F. Better hemodynamic profile of laryngeal mask airway insertion compared to laryngoscopy and tracheal intubation. Iran Red Crescent Med J. 2015;17(8):e28615.
- Brimacombe J, Berry A. Insertion of LMA-an indication for propofol? Anaesthesia intensive care. 192 Aug;20(3):394-395.
- Wilkins CJ, Cramp PG, Staples J, Stevens WC. Comparison of the anesthetic requirement for tolerance of laryngeal mask airway and endotracheal tube. Anesth Analg. 1992;75:7947.
- Singh R, Arora M, Vajifdar H. Randomised double blind comparison of ketamine-propofol and fentanyl-propofol for the insertion of laryngeal mask airway in children. J Anaesthesiol Clin Pharmacol. 2011;27:91-6.
- Begee Z, Demirbilek S, Onal D, Erdil F, Ilksen Toprak H, Ozcan Ersoy M. Ketamine or alfentanil administration prior to propofol anaesthesia: The effects of ProSeal laryngeal mask airway conditions and haemodynamic changes in children. Anaesthesia. 2009;64:282-6.
- Gupta A, Kaur S, Attri JP, Saini N. Comparative evaluation of ketamine- Propofol, fentanyl-propofol and butarphanol-propofol on haemodynamics and laryngeal mask airway insertion conditions. J Anaesthesiol Clin Pharmacol. 2011;27:74-8.
- Goel S, BBhardwaj N, Jain K. Efficacy of ketamine and midazolam as co-induction agents with propofol for laryngeal mask insertion in children. Pediatr Anaesth. 2008;18:628-34.
- Salem WT. A comparison of midazolam and mini-dose succinylcholine to aid laryngeal masks airway insertion during propofol anaesthesia. J Egypt Natl Cancer Inst. 2000;12:65-9.
- Aghamohammadi D, Eydi M, Hosseinzadeh H, Amiri Rahimi M, Golzari SE. Assessment of mini-dose succinylcholine effect on facilitating laryngeal mask airway insertion. J Cardiovasc Thorac Res. 2013;5:17-21.
- Goyagi T, Tanaka M, Nishikawa T. Oral clonidine pre-medication reduces propofol requirement for LMA insertion. Can J Anaesth. 2000;47:627-630.
- Cook TM, Seavell CR, Cox CM. Lignocaine to aid the insertion of the laryngeal mask airway with thiopentone. Anaesthesia. 1996;51:787-790.
- Young MSA, Clarke RSJ, Dundee JW. Intubating conditions with AH8165 and Suxamethonium. Anaesthesia. 1975;30:30-33.
- Nimmo SM, Mc Cann N, Broome IJ, Rob HM. Effectiveness and sequelae of very low dose suxamethonium for nasal intubation. Br J Anaesth 1995;74:31-34.
- Lund I, Stovener J. Dose response curves for tubocurarine, alcuronium and pancuronium. Acta Anaesthesiol Scand. 1970;37(5):238-242.
- Joshi GP. Rapid Recovery from Ambulatory Surgery: The New Paradigm in Ambulatory Anesthesia. Available from: <http://www.iars.org/assets/1/7/RCL-01.pdf>.
- Scanlon P, Carey M, Power M, Kirby F. Patient response to laryngeal mask insertion after induction of anaesthesia with propofol or thiopentone. Can J Anaesth. 1993;50:464-66.
- JG Namrata, P Parikh Geeta. Midazolam or mini-dose Succinylcholine as a co-induction agent to aid Laryngeal mask airway insertion during propofol Anaesthesia. In J Res Med. 2015;4(1):62-68.
- Short TG, Chui PT. Propofol and midazolam act synergistically in combination. British journal of anaesthesia. 1991 Nov;67(5):539-45.
- Oxorn D, Ferris LE, Harrington E, Orser BA. The effect of midazolam on propofol anaesthesia: propofol dose requirements, mood profiles and peri-operative dreams. Anaesth Analg. 1997;85:553-59.
- Murphy PJ, Erskine R, Langton JA. Effect of intravenous administration of diazepam, midazolam and flumazenil on sensitivity of upper airway reflexes. Anaesthesia 1994;49:105-110.
- Dhamotharan S, Singh NR, Singh SS, Singh MB. Comparative evaluation of fentanyl and midazolam with propofol induction on laryngeal mask airway insertion conditions: A study. J Med Soc. 2014;28: 185-9.