

# International Journal of Food, Nutrition and Dietetics

**Editor-in-Chief**

**Balwinder Sadana**

College of Home Science, Punjab Agricultural University  
Ludhiana

## **National Editorial Board**

**E. Lakshmi**, Kattankulathur

**Jamuna Prakash**, Mysore

**Ruma Bhattacharyya**, Jorhat

**S.S. Thorat**, Rahuri

**Vandana Bharti**, Indore

**Vijaya M. Nalwade**, Prabhani

## **International Editorial Board**

**Aruna Amarasinghe**, Sri Lanka

**Alaa Jabbar Al-Manhel**, Iraq

**Samrita Dogra**, USA

**Managing Editor:** A. Lal

**Publication Editor:** Manoj Kumar Singh

**International Journal of Food, Nutrition & Dietetics (IJFND)** (pISSN 2322-0775; eISSN: 2455-569X), a broad-based peer-reviewed journal publishes the most exciting researches with respect to the subjects of nutrition and food sciences. The journal covers current thinking on food and nutrition emphasizing the practical and social application of ideas. Special editions focusing on topics including micronutrients, special diets for management of health problems and cost sector catering provide readable content that is an invaluable resource for practitioners and academics wishing to inform themselves, their colleagues, or the public on modern thinking, research, and attitudes to food and nutrition.

**Readership:** Academics and researchers in the field, Dietitians, Food company managers, Food research institutes, Health care professionals, Nutritionists.

**Indexing Information:** Index Copernicus, Poland; ProQuest, USA; Genamics JournalSeek; CiteFactor, USA; Cosmos Impact Factor, Germany etc.

---

**For all other queries** Red Flower Publication Pvt. Ltd., 48/41-42, DSIDC, Pocket-II, Mayur Vihar Phase-I, Delhi - 110 091 (India), Phone: 91-11-22754205, Fax: 91-11-22754205, E-mail: info@rfppl.co.in, Web: www.rfppl.co.in

---

**Disclaimer** The opinion in this publication is those of the authors and is not necessarily those of the International Physiology the Editor-in-Chief and Editorial Board. Appearance of an advertisement does not indicate International Physiology approval of the product or service.

© Red Flower Publication Pvt. Ltd. 2019. All rights reserved. No part of the journal may be reproduced, stored in a retrieval system or transmitted in any form or by any means, electronic, mechanical, photocopying, recording or otherwise, without prior permission of the New Indian Journal of Surgery.

**Printed at** Saujanya Printing Press, D-47, Okhla Industrial Area, Phase-1, New Delhi - 110 020.

**International Journal of Food, Nutrition & Dietetics (IJFND) (pISSN: 2322-0775, eISSN: 2455-569X)**, a broad-based peer-reviewed journal publish the most exciting researches with respect to the subjects of nutrition and food sciences. The journal covers current thinking on food and nutrition emphasizing the practical and social application of ideas. Special editions focusing on topics including micronutrients, special diets for management of health problems and cost sector catering provide readable content that is an invaluable resource for practitioners and academics wishing to inform themselves, their colleagues, or the public on modern thinking, research, and attitudes to food and nutrition.

**Readership:** Academics and researchers in the field, Dietitians, Food company managers, Food research institutes, Health care professionals, Nutritionists

**Indexing Information:** Index Copernicus, Poland; ProQuest, USA; Genamics JournalSeek; CiteFactor, USA; Cosmos Impact Factor, Germany etc.

#### Subscription Information

##### India

**Institutional** (1 year) (Print+Online): INR 5500

##### Rest of the World

**Institutional** (1 year) (Print+Online): USD430

#### Payment instructions

##### *Online payment link:*

<http://rfppl.co.in/payment.php?mid=15>

##### *Cheque/DD:*

Please send the US dollar check from outside India and INR check from India made. Payable to 'Red Flower Publication Private Limited'. Drawn on Delhi branch

##### *Wire transfer/NEFT/RTGS:*

Complete Bank Account No. 604320110000467

Beneficiary Name: Red Flower Publication Pvt. Ltd.

Bank & Branch Name: Bank of India; Mayur Vihar

MICR Code: 110013045

Branch Code: 6043

IFSC Code: BKID0006043 (used for RTGS and NEFT transactions)

Swift Code: BKIDINBBDOS

**Send all Orders to:** Subscription and Marketing Manager, Red Flower Publication Pvt.

Ltd., 48/41-42, DSIDC, Pocket-II, Mayur Vihar Phase-I, Delhi - 110 091(India),

Phone: 91-11-45796900, 22754205, 22756995, E-mail: [sales@rfppl.co.in](mailto:sales@rfppl.co.in),

Website: [www.rfppl.co.in](http://www.rfppl.co.in)

---

## Contents

---

### *Original Articles*

- Organic Farming for Sustainable Food and Nutrition with Rural Empowerment** 5  
Sangeeta Ahuja
- Nutritional and Ethnomedicinal Potential Plants of the Qur'an: An Overview - I** 11  
T.P. Mall
- Consumer's Knowledge, Attitude and Practices (KAP) Regarding Organic Foods in Relation to Demographic Factors of Punjab** 23  
Seema Rani, Neerja Singla, Poonam Bakheta
- Impact of Nutrition Counseling on Dietary Intake of Obese Women from Low, Middle and High Income Group of Hoshiarpur City of Punjab** 33  
Shagun Aggarwal, Neerja Singla, Poonam Bakheta

### *Review Articles*

- Vegetable Pigments: Effect of Ph and Heat: A Review** 41  
E. Lakshmi, E. Saraswathy
- Guidelines for Authors** 45

## SUBSCRIPTION FORM

I want to renew/subscribe international class journal "International Journal of Food, Nutrition and Dietetics" of Red Flower Publication Pvt. Ltd.

**Subscription Rates:**

- Institutional: INR5500/USD430

Name and complete address (in capitals): \_\_\_\_\_

**Payment detail:**

**Online payment link:** <http://rfppl.co.in/payment.php?mid=15>

Cheque/DD: Please send the US dollar check from outside India and INR check from India made payable to 'Red Flower Publication Private Limited'. Drawn on Delhi branch.

**Wire transfer/NEFT/RTGS:**

Complete Bank Account No. 604320110000467

Beneficiary Name: Red Flower Publication Pvt. Ltd.

Bank & Branch Name: Bank of India; Mayur Vihar

MICR Code: 110013045

Branch Code: 6043

IFSC Code: BKID0006043 (used for RTGS and NEFT transactions)

Swift Code: BKIDINBBDOS

**Term and condition for supply of journals**

1. Advance payment required by Demand Draft payable to **Red Flower Publicaion Pvt. Ltd.** payable at **Delhi**
2. Cancellation not allowed except for duplicate payment.
3. Agents allowed 10% discount.
4. Claim must be made within six months from issue date.

**Mail all orders to**

Subscription and Marketing Manager

Red Flower Publication Pvt. Ltd.

48/41-42, DSIDC, Pocket-II

Mayur Vihar Phase-I

Delhi - 110 091(India)

Phone: 91-11-45796900, 22754205, 22756995, Cell: +91-9821671871

E-mail: [sales@rfppl.co.in](mailto:sales@rfppl.co.in)

## Organic Farming for Sustainable Food and Nutrition with Rural Empowerment

Sangeeta Ahuja

---

### How to cite this article:

Sangeeta Ahuja. Organic Farming for Sustainable Food and Nutrition with Rural Empowerment. *Int J Food Nutr Diet.* 2019;7(1): 5-10.

### Abstract

The organic food is grown or raised can have a major impact on the mental and emotional health as well as the environment. Organic foods often have more beneficial nutrients, such as antioxidants, than their conventionally-grown counterparts and people with allergies to foods, chemicals, or preservatives often find their symptoms lessen or go away when they eat only organic foods. There is a direct mapping of nutrition management and Agriculture. Agriculture plays a very important role for the growth and development of any country. Agriculture is the sedentary human civilization whereas farming created food surpluses that enabled to live in cities. The Organic farming provides an opportunity to achieve sustainable agriculture that has grown out of the conscious efforts contributing towards rural development and livelihoods to farmers without destroying the natural resource basis. This has also benefited in the way as it reduces the environmental impacts of conventional agriculture and increase productivity in small farmers' land holdings and the costly external inputs, and guarantees price premiums for organic products. The scientific and technological inputs have been major drivers of growth and development in agriculture and allied sectors that have enabled us to achieve self-reliant food security with a reasonable degree of resilience even in times of natural calamities, in recent years. In the present times, agricultural development is faced with several challenges relating to state of natural resources, climate change, fragmentation and diversion of agricultural land to non-agricultural uses, factor productivity and global trade. There is an urgent need to identify the issues of research and strategies for organic research in agri-horticultural sector, integrated farming systems, resource bases and agro-ecological zones. Decision support systems, expert systems and statistical methodologies/ techniques have to be developed in the field of organic farming in order to increase the agricultural production of the country.

**Keywords:** Small and marginal farmers; Sustainable livelihoods; Decision support system; Nutrition.

---

### Introduction

India has tremendous potential to become a major exporter of agricultural commodities in the international market. During the next decade, the population in developing countries like India, will

face changes in climate patterns and variability that will contribute to severe water shortages or flooding, and also will cause shifts in crop growing seasons due to the rising temperatures which in turn will increase shortage in food and distribution of disease vectors, putting populations at greater environmental health risks. Organic agriculture thus, as an adaptation strategy to climate variability, is a concrete and promising option for rural communities and has additional potential as a mitigation strategy. In India, organic farming was started first by the agribusiness entrepreneurs around 2000, supported by the Ministry of Commerce. Initially, organs of the Ministry of Commerce and APEDA to promote organic production of export commodities due to

---

**Author Affiliation:** Scientist (S.S.), IASRI (ICAR), Ministry of Agriculture and Farmers Welfare, New Delhi 110012, India.

**Corresponding Author:** Sangeeta Ahuja, Scientist (S.S.), IASRI (ICAR), Ministry of Agriculture and Farmers Welfare, New Delhi 110012, India.

**E-mail:** reach2san@yahoo.com

new trade regime of Non-tariff barriers linked to residue contaminations and unsafe agri-products and because of premium prices organic enterprises could enjoy. Indian organic initiative therefore started with setting up of regulatory mechanism, necessary for exports, rather than on principles of organic. Traditionally, Indian agriculture is look like an organic culture in many ways. About 75 percent of the farmers are small and marginal, being practicing organic farming by default due to the unaffordable modern agriculture technologies and rain-fed ecosystems. Presently, the farming and farmers are passing through a transitional phase, comprising of several factors and processes, which include both constraints and opportunities. Integrated farming system is the strength of hill farming.

### Current scenario of organic farming

At present, organic farming is practiced in 162 countries and 37 million hectares of agricultural land are managed organically by 1.8 million farmers. The current market for organic foods in India is pegged at Rs. 2500 crores, which according to ASSOCHAM, is expected to reach Rs. 6,000 crores by 2015. Thus, a huge potential is seen in the nascent Indian organic sector. Organic products, which until now were mainly being exported, are now finding consumers in the domestic market also [4] (Table 1). In order to feed this larger population, food production must increase by 48.5 percent. India ranks second worldwide in farm output but

**Table 1:** Status of organic products in India (2013-14)

<b>Total quantity exported</b>	<b>135</b>
Value of total export	US\$ 374 million
Total certified area (including under 5.21 million hectare cultivation, forest and wild harvest)	5.21 million hectare
Organic crops/ commodities/ products produced	Sugarcane, Cotton, Basmati rice, Pulses, Tea, Spices, Coffee, Oil Seeds, Fruits and their value added products, organic cotton fiber, functional food products etc.
Countries importing Indian organic products	EU, US, Switzerland, Canada, South East Asian countries and South Africa.
Share of Indian organic products in export	Oil seeds - Soybean (41%) lead among the products exported followed by Cane Sugar (26%), Processed food products (14%), Basmati Rice (5%), Other cereals & millets (4%), Tea (2%), Spices (1%), Dry fruits (1%).

the economic contribution of agriculture to India's GDP is steadily declining with the country's broad-based economic growth. The contribution of this important sector to the national GDP is declining (14.6%). Still, agriculture is demographically the broadest economic sector and plays a significant role in the overall socio-economic fabric of India.

### Experimentation

Rigorous experimentation with survey analysis have been conducted in the various branches of Agriculture and the data have been statistically computed.


### Land Holdings

In India, the average holding size is estimated to be 0.32 and 0.24 ha in 2030 and 2050 respectively. 63 percent holdings are below 1 ha accounting for 19 percent of the operated area while over 86 percent of holdings are less than 2 ha account for nearly 40 percent of the area (APCAS, 2010). As per estimates, more than 95% of the holdings will be under the category of small and marginal holders in 2050 (Figs. 1-3)[5,6].

### Export Markets: Buying from India

The scenario of the export market shown in Fig. 4 [7]

The Organic produce have lot of beneficial effects. It contains fewer pesticides are often fresher, is better for the environment. Organically raised animals are not given antibiotics, growth hormones, or fed animal by-products. Organic meat and milk are richer in certain nutrients.



**Fig. 1:** Decreasing farm land holdings size.




Fig. 2: Food demand projections for 2050




Fig. 3: Changing food consumption pattern




Fig. 4: Export Market

**Solution**

The solution of good health and nutrition is organic food and vice versa and by implementing the Organic farming in rural places the rural empowerment can be sustained (Fig. 5).

The research and development i.e. the issues & challenges are tackled by departments and institutes of Agriculture viz., SAUs, ICAR Institutes, and KVKs. The comparison of production costs, yields




Fig. 5: Solutions of organic agriculture

and income of organic and conventional farming systems is needed. The improvement of organic production technologies. More research on inputs to organic farming and make results available to producers and policy makers. The private sector can potentially collaborate with SAUs and ICAR units for developing quality inputs. It is shown in fig. 6.

These are fusion of Technologies for Sustainable Rural Development viz., Database Technology, Internet/Intranet Technology [1-3], GIS and Remote Sensing, Image Processing, GPS, Artificial Intelligence, Decision Support System and Statistical Modeling.

Organic Certification & Accreditation is also needed as shown in Fig. 7.

By 2050, the world’s population will reach 9.1 billion, 34 percent higher than today and India will be the most populous country (1.6 billion) on the earth. Urbanization will continue at an accelerated pace (2.4%) and about 50 percent of the India’s population will be urban as compared to present 29.5 percent. Currently, India ranks 33<sup>rd</sup> in terms of total land under organic cultivation and 88<sup>th</sup> position for agriculture land under organic crops to total farming area. The cultivated land under certification is around 4.43 million ha. At present, India produced around 3.88 million MT of certified organic products including all varieties of




Fig. 6: Implementation




Fig. 7: Decision Making Process



Fig. 8: Decision Support Systems




Fig. 9: Spatial decision support for policy decisions for crop productivity management



Fig. 10: Certification

food products. Global consumers are increasingly looking forward to organic food that is considered safe and hazard-free. Global demand for organic products remains robust, with sales increasing by over 5 billion US Dollars a year.

### Conclusion

The nutrition with sound health is directly related to organic farming which further leads to rural empowerment. The livelihood of many people are dependent on this field. The survival of mankind without food is not possible on earth so organic farming not only sustain the rural empowerment but very much beneficial for health and nutrition. The application of ICT and their involvement in every stage of production from soil identification to till production. In present scenario, every part of the world is conducting the survey to gather empirical information on crop, livestock and other agriculture resource. Planning and development process of agriculture DSS constitute a multi-complex

problem which is very difficult to be solved, if it is not faced thoroughly. The decision making process is demand from producer (information)/ decision maker a qualitative evaluation. The model will lead to make agriculture DSS more appropriate and lead to expected agriculture solution model.

### References

1. Kruglinski, David. Inside Visual C++, I Edition. Microsoft Press, Washington, 1996.
2. Jeff, Prosize. Programming Windows with MFC, II Edition. Microsoft Press, Washington, 1999.
3. Richter - Programming Applications for Microsoft Windows, IV Edition. Microsoft Press, Washington, 1999.
4. <http://www.apeda.gov.in>.
5. Singh, et al. Food demand projections for 2050. 2013.
6. Amarasinghe et al. Changing food consumption pattern. 2007.
7. <http://www.FiBL.org>

## Nutritional and Ethnomedicinal Potential Plants of the Qur'an: An Overview - I

T.P. Mall

---

### How to cite this article:

T.P. Mall. Nutritional and Ethnomedicinal Potential Plants of the Qur'an: An Overview - I. Int J Food Nutr Diet. 2019;7(1):11-21.

### Abstract

Our ancient literatures of Hindus', Muslims as well as of Cristian's are full of plants described about their nutritional and medicinal potential. A significant number of plants described in their texts are still unidentified because of least or no interest in plant taxonomy among neither graduate scientists in well-developed different organisations nor respected teachers of Universities or Colleges of repute. We can say with confidence that there are no teachers who can teach Plant Taxonomy in significant numbers of academic institutions. A man of Biotechnology can do nothing if he/she can not identify the plant on which the work has to be worked out. As a result, due to non-identification of plants we cannot use our wealth of knowledge which has been provided by our ancestors present in form of scripts. We were presented a book written by Dr. M.I.H. Farooqi entitled "*Quruani Poudhe-Vagayanic Dhrsti Se*" while in Seminar organised by UP Biodiversity Board, Lucknow. We found that there are seventy-one plants being reported in Holy Qur'an and Bible. We have consulted the literatures available as well as the tribal of Bahraich about the uses of the plants available. The perusal of the alphabetical list of plants of Holy Qur'an reveals that there are 71 plant species representing 48 genera of 30 families. Brassicaceae, Cucurbitaceae and Moraceae family were found to be the biggest family represented by 6 plant species each whereas Caesalpiniaceae, Papilionoideae and Poaceae with 5 plant species each; Rosaceae and Rhamnaceae with 4 plant species; Mimosaceae, Liliaceae, Pinaceae, Euphorbiaceae, Oleaceae, Lytharaceae, Lamiaceae and Arecaceae with 2 plant species and rest fourteen species viz., Malvaceae, Asclepidiaceae, Lauraceae, Bixaceae, Dipterocarpaceae, Juglandaceae, Cupressaceae, Lecnoraceae, Loranthaceae, Anacardiaceae, Sterculiaceae, Ericaceae, Salvadoraceae, and Vitaceae is being represented by single plant species each.

**Keywords:** Qur'an; Nutritional; Ethnomedicinal; Potential Plants.

---

### Introduction

Herbal Medicine is the oldest form of medicine known to mankind. It was the mainstay of many early civilizations and still the most widely practiced form of medicine in the world today.

**Author Affiliation:** Ex-Associate Professor & Head, Postgraduate Department of Botany, Kisan PG College, Bahraich, Uttar Pradesh 271 801, India.

**Corresponding Author:** T.P. Mall, Ex-Associate Professor & Head, Postgraduate Department of Botany, Kisan PG College, Bahraich, Uttar Pradesh 271 801, India.

**E-mail:** [drtejpratapmall@gmail.com](mailto:drtejpratapmall@gmail.com)

Ethnobotany is one of the most interesting themes of economic botany which might have first of all came into the existence probably when earliest man of "stone age" observed the animals mostly the apes and monkeys eating certain plants or plant parts ex. Fruits, leaves and even inflorescences to satisfy their hunger. Therefore, on the basis of plants usage first of all by animals and later on by the human beings the concepts of Ethnobotany and Ethnozoology were evolved, which merged into a common term known as Ethnobiology. However, the term Ethnobotany was first of all used in the last of 19<sup>th</sup> century by J.W. Harsh Berger (1895) to indicate the interrelationship of plants with aboriginal people or tribal societies [Trivedi and

Sharma, 2011]. In many parts of the world, wild plants are obtained from forests or wild areas designated for extractive resources and managed by local communities [Jadhav *et al.*, 2011]. Wild edible plants provide food quantity as well as medicines [Patale *et al.*, 2015].

India is one of the twelve mega-biodiversity countries of the World having rich vegetation with a wide variety of plants. As per the 2001 census, the tribal population of India is 8.43 crore, constituting 8.2% of total population of the country [Annual Report, 2005-2006]. With enormously diversified ethnic groups and rich biological resources, India represents one of the great emporia of ethnobotanical wealth [Pal, 2000]. Even today, tribal's and certain local communities in India still collecting and preserving locally available wild and cultivated plant species for their day today life [Mahishi, *et al.*, 2005 and Ayyanar, *et al.*, 2010].

We were presented a book written by Dr. M.I.H. Farooqi entitled "*Quruani Poudhe-Vagayanic Dhrusti Se*" while in Seminar organised by UP Biodiversity Board, Lucknow. We found that there are seventy-one plants being reported in Holy Qur'an and Bible. We have consulted the literatures available as well as the tribal of Bahraich about the uses of the plants available. The perusal of the alphabetical list of plants of Holy Qur'an reveals that there are 71 plant species representing 48 genera of 30 families. Brassicaceae, Cucurbitaceae and Moraceae family were found to be the biggest family represented by 6 plant species each whereas Caesalpiniaceae, Papilionoideae and Poaceae with 5 plant species each; Rosaceae and Rhamnaceae with 4 plant species; Mimosaceae, Liliaceae, Pinaceae, Euphorbiaceae, Oleaceae, Lytharaceae, Lamiaceae and Arecaceae with 2 plant species and rest fourteen species viz., Malvaceae, Asclepidiaceae, Lauraceae, Bixaceae, Dipterocarpaceae, Juglandaceae, Cupressaceae, Lecnoraceae, Loranthaceae, Anacardiaceae, Sterculiaceae, Ericaceae, Salvadoraceae, and Vitaceae is being represented by single plant species each. We are enumerating only five plants viz., *Acacianilotica*, *Acacia senegal*, *Acacia seyal*, *Alhigimaurorum* and *Allium cepa* in detail.

### Enumeration

1. *Acacia nilotica* (L.) Delile., Gum arabic tree, Babla; Baboul; Babul; Dauria; Godi; Godi babul; Kabuli kikar; Kaora; KauliaKauria; Kavadi; Kikar; Kikkar; Ram babul; Ramkanta; Ramkati babul; Teli babul; Telia; Telia babul; Vedi; Arabic gum tree; Babul acacia; Blackthorn; Egyptian mimosa; Egyptian thorn; Prickly acacia; Prickly mimosa;

Scented thorn; Cented-pod acacia; *Acacia gomifera*; *Acacia a gomme*; *Acacia d'Arabie*; *Acacia de cayenne*; Gommier rouge garad; Sunt; Babul (*Mimosaceae*):

Sub species: *Acacia nilotica* subsp. *adstringens*; *Acacia nilotica* subsp. *cupressiformis*, *Acacianilotica* subsp. *hemispherica*; *Acacia nilotica* subsp. *indica*; *Acacia nilotica* subsp. *kraussiana*; *Acacia nilotica* subsp. *leiocarpa*; *Acacia nilotica* subsp. *nilotica*, *Acacia nilotica* subsp. *subalata*; *Acacia nilotica* subsp. *tomentosa*.

Synonymized variety: *Acacia arabica* var. *indica*; *Acacia arabica* var. *nilotica*.

*Acacia nilotica* is a pioneer species that is relatively fast growing on arid sites. It is an important riverine tree in India, Sudan and Senegal, where it is planted for timber. Le Houerou (1988) describes native range habitat types for several subspecies, *tomentosa*, *adstringens* and *nilotica*. These are all described in the context of flood plain or water course habitats with varying degrees of flooding. According to Weber (2003), *Acacia nilotica* invades grass and savanna habitats. In Queensland, Australia it has invaded plains of Mitchell grass (*Astrelba* spp.), occurring on soils with a high clay content and sandy loams providing there is sufficient moisture, waterways and seasonally flooded plains including saline areas (Carter, 1998).

### Ethnobotanical Potential

*Acacia nilotica* is popular as an agroforestry tree, either sown in lines 5 m apart in agricultural fields, or on field crop boundaries. As a fodder tree, it is utilized in many different silvi-pastoral systems, and its sweet-smelling pods are particularly sought out by animals. It is extensively used in land rehabilitation, being planted on saline and alkaline soils. It will also grow when irrigated with tannery effluent, or saline water, and effectively colonizes waste heaps from coal mines. The tree is popular as a shelterbelt, and there is interest in *Acacia nilotica* subsp. *cupressiformis* as a windbreak surrounding fields because its narrow crown form produces less shade than other taxa.

It is also a popular ornamental tree, and is frequently planted in India as an avenue tree.

Since the time of the Pharaoh's, large timber trees have been exploited from the riverine forests of the Nile in Sudan and Egypt. At present, forests in the Sudan are managed on a 20-30-year rotation, producing termite resistant timber especially suitable for railway sleepers. In India and Pakistan, the riverine plantations are managed on 15-20-year rotations for fuelwood and timber such as mine props. The strong and durable wood is nearly twice

as hard as teak, and is very shock resistant. It is used for a wide range of construction uses, and in tool handles and carts. Wood properties are reviewed in Rao and Purkayastha (1972), Goldsmith and Carter (1981), Tewari and Rajput (1987) and Troup and Joshi (1983), and its rayon pulp and paper pulp properties in FAO (1980) and Guha *et al.* (1974). The dark brown hardwood has a high calorific value of 4950 kcal/kg, making excellent fuelwood and high-quality charcoal (NAS, 1980).

The pods and leaves have high levels of crude (12.4%) and digestible protein (8%) and energy (7.2 MJ), and are rich in minerals (Le Houerou, 1980). Pods are used as a supplement to poultry rations in India. Dried pods are relished and particularly sought out by animals grazing on rangelands as the pods mature towards the end of the dry season. In India, branches are commonly lopped for fodder. Pods are best fed as a supplement.

A by-product from felling is the bark, which has high levels of tannin (12-20%) used for tanning leathers in India, and the pods of *A. nilotica* subsp. *nilotica* have been used for over 6000 years in Egypt for tanning. *A. nilotica* subsp. *adstringens* is used both for tanning and as a dye in Nigeria.

The gum was originally called gum arabic and has been collected from the Nile forests since the time of the Pharaoh's, for use in paints and medicines. It has some properties similar to those of true gum arabic (from *Acacia senegal*) and is frequently used in calico printing, dyeing, sizing material for silk and cotton and in paper manufacture in India. In Mumbai, India, it is marketed as Amravati gum.

It also has been found to be a powerful molluscicide and algicide, and the fruits, when added to ponds in Sudan, killed snail species which carry schistosomiasis without affecting fish (Ayoub, 1982). The tree is a good host plant for growing lac (shellac) in Sind, Pakistan. An extract of the root is a potential inhibitor of Tobacco mosaic virus. In eastern Java, sprouted seeds are eaten as vegetables, and well-roasted seeds are mixed with coffee (Lemmens and Wulijarni-Soetjpto, 1991). There are many other reported uses (Fagg and Greaves, 1990).

#### Uses List

*Animal feed, fodder, forage:* Fodder/animal feed, Invertebrate food for lac/wax insects.

*Environmental:* Agroforestry, Boundary, barrier or support, Revegetation, Shade and shelter.

Windbreak, Fuels, Charcoal, Fuelwood.

*Materials:* Carved material, Dye/tanning, Fibre, Gum/resin.

*Miscellaneous materials:* Poisonous to mammals, Wood/timber, Wood Products, Charcoal, Railway sleepers, Round wood, Building poles, Carpentry/joinery (exterior/interior) Engineering structures, Exterior fittings, Fences, Flooring, For heavy construction, For light construction.

*Industrial and domestic woodware:* Tool handles, Wood carvings.

#### Ethnomedicinal Potential

The tannin content contributes to the many medicinal uses of *A. nilotica*, acting as a powerful astringent. Medicinal, pharmaceutical. Source of medicine/pharmaceutical uses.

**2. Acacia senegal** (L.) Willd., Gum Arabic Tree; Gum Acacia; Gum Senegal Tree; Kumththa; Kumatiyo; Kumat; Kumatio; Khair; Babul; Shvetakhadira (Mimosaceae):

Synonyms: *Acacia verek*, *Mimosa senegal* L., *Senegaliasenegal* (L.) Britton, *Acacia circummarginata* Chiov., *Acacia cufodontii* Chiov., *Acacia glaucophylla* sensu Brenan, *Acacia kiniongesensu* Brenan, *Acacia oxyosprion* Chiov., *Acacia rupestris* Boiss., *Acacia senegal* (L.) Willd. subsp. *modesta* (Wall.) Roberty, *Acacia senegal* (L.) Willd. subsp. *senegalensis* Roberty, *Acacia somalensis* sensu Brenan, *Acacia spinosa* Marloth & Engl., *Acacia thomasi* sensu Brenan, *Acacia volkii* Suess., *Mimosa senegal* L. (L.) Britton.

Gum Arabic tree is a small deciduous acacia tree, native to semi-desert regions of Sub-Saharan Africa, Pakistan, and north-western India. It grows to a height of 5-12m, with a trunk up to 30 cm in diameter. Thorns are placed just below the nodes, either in threes up to 7 mm long, with the middle one hooked downwards and the lateral ones curved upwards, or solitary with the laterals absent. Leaves are double-compound, up to 2.5 cm long. Leaf-axis is finely downy with 2 glands; pinnae are 6-20 pairs; leaflets are small, 7-25 pairs, rigid, leathery, smooth, linear to elliptic-oblong, ciliate on margins, pale glaucous-green, tip blunt to somewhat pointed. Flowers are borne in not very dense spikes 5-10 cm long, carried on stalks 0.7-2 cm long. Flowers are normally produced with the leaves. Sepal cup is bell-shaped, glabrous, deeply toothed. Flowers are white to yellowish, fragrant, stalkless. Pod is straight or slightly curved, ret rap-shaped, 7.5-18 cm long, 2.5 cm wide, thin, light brown or gray, papery or woody, firm, smooth. The tree produces gum arabic, which is

used as a food additive, in crafts, and as a cosmetic. The gum is drained from cuts in the bark, and an individual tree will yield 200 to 300 grams. Seventy percent of the world's gum arabic is produced in Sudan. Flowering: January–March.

#### *Ethnobotanical Potential*

Gum arabic is the exudate formed in the bark of *Acacia senegal*, *Acacia seyal* Delile and a few related species. In Sudan and sometimes in international trade, a distinction is made between 'gum hashab' from *Acacia senegal* and 'gum talha' from *Acacia seyal*.

Gum arabic is easily soluble in water and forms solutions over a wide range of concentrations. It has highly valued emulsifying, stabilising, thickening and suspending properties and does not become highly viscous.

The food industry uses 60–75% of the world production. In confectionery, gum arabic is used to prevent crystallisation of sugar, as an emulsifier, and as a glaze or topping in bakery products; in soft drinks and alcoholic drinks it is used either as a vehicle for flavouring or as a stabiliser or clouding agent; in frozen dairy products gum arabic is used for encapsulating flavours such as citrus oils.

Its pharmaceutical use has decreased, but it is still used as a suspending or emulsifying agent and in tablet manufacture, where it functions as a binding agent or as a coating prior to sugar coating.

Gum arabic is used in the printing industry for coating offset lithographic plates to prevent oxidation, to increase their hydrophilic properties and to make them repellent to ink.

It is also a base for photosensitive chemicals. In ceramics, gum arabic helps to strengthen the clay. Other technical applications include pyrotechnics and ink manufacturing.

In textiles, paints, paper size and adhesives (including the traditional office glue and postage stamps) its use has decreased to very low levels in recent years.

Gum arabic is used locally in special dishes and as chewing gum; it has medicinal applications for both humans and livestock, e.g. to treat skin diseases and inflammation.

*Acacia senegal* is a multipurpose tree. The foliage and pods are an important fodder source for camels and goats.

Seed may be dried and conserved for human consumption mainly as an emergency food.

The wood is used for small-scale construction purposes and agricultural implements; it yields a fuelwood of good quality, that can be made into good charcoal.

The thorny branches are often used to make 'dead fences' to enclose livestock or protect agricultural fields.

Being a very drought-resistant tree, it is planted for sand dune fixation, windbreaks and shelter belts in arid regions.

The flowers are a source of honey.

Cordage is made from the roots, either directly or after beating to extract the fibres; its strength makes it suitable for well ropes and fishing nets.

The seed contains a fat which is used both in medicine and for soap making.

#### *Ethnomedicinal Potential*

The gum is used for soothing mucous membranes of the intestine and to treat inflamed skin.

It is also reportedly used as for its astringent properties, to treat bleeding, bronchitis, diarrhoea, gonorrhoea, leprosy, typhoid fever and upper respiratory tract infections.

Bark, leaves and gum are used as an astringent to treat colds, ophthalmia, diarrhoea and haemorrhages.

#### *Environmental impact*

Soil binder and soil improver.

Gum arabic trees can help prevent desertification through dune stabilisation and by acting as a wind break. It is valued in agroforestry systems where it is combined with crops such as millet, sorghum, sesame, and groundnut, and where it is reported to improve soil fertility, though its N-fixing status is debated (Orwa *et al.*, 2009).

Flowers are a good bee forage; honey amber coloured, very mild aroma, granulates rapidly, within 2 weeks.

#### *Nutritional attributes*

Gum arabic tree leaves are of good nutritional value, with a protein content ranging from 15 to 33% of DM when the leaves are fresh. The leaves are also relatively low in fibre (crude fibre 14-25% of DM). The pods are also rich in protein but with a higher fibre content.

### Ruminants

Gum arabic tree leaves are a valuable forage for all ruminants except cattle. Though forage yield remains low, leaves are relished by livestock, and provide good quality forage with high protein content and digestibility (in vitro OM digestibility ranges from 66 to 86%), and a low tannin content. Gum arabic tree forage is available at the end of the dry season, when other forages are still scarce. At that time the leaves may account for up to 15-20% of the feed intake of sheep and camelids, and 45% and above for goats. In terms of its relative palatability, *Acacia senegal* ranks highest among local fodder species (Ickowicz *et al.*, 2005).

Pods are palatable to livestock but their consumption destroys seed stocks and may have deleterious effects on stand management (Ickowicz *et al.*, 2005). It was shown that only 33% of gum arabic tree seeds could be retrieved from cattle faeces, and only 1% from sheep and goats. Their viability was moreover significantly reduced: 1% germination for seeds found in cattle faeces and no germination for those retrieved from sheep and goats' faeces (Danthu *et al.*, 1996).

The most important use of the gum arabic tree is the harvest of gum arabic, an exudate from the bark that is tapped for this purpose during the dry season (Orwa *et al.*, 2009). Several thousand tons of gum arabic are internationally traded every year, mainly in Europe and the USA.

### Commercial uses

Food, (flavour fixative, emulsifier, stabilizer of dairy products), pharmaceuticals (these two sectors representing 60-75% of the use of gum arabic), and industrial products (inks, pigments, polishes).

Gum arabic was reported to have antidotal effects as it can destroy many alkaloids (Duke, 1983).

*Acacia senegal* seeds are traditionally used for human nutrition in Rajasthan (Ram *et al.*, 2014).

The wood is valued as firewood, and can be used to produce charcoal. The wood is also used to make utensils, poles and fence-posts.

The bark and the roots provide fibre and make strong ropes and fishing nets (Orwa *et al.*, 2009; Duke, 1983). Gum arabic trees provide valuable fodder to sheep, goats and camels. Leaves and pods are browsed by domestic and wild ruminants. Flowers provide valuable nectar to bees for honey production (Orwa *et al.*, 2009).

**3. *Acacia seyal* Del. (*Vachellia seyal* (Delile) Hurter)** White galled acacia; Shittim Wood; White

### Whistling wood (*Caesalpinaceae*):

*Varieties: Vachellia seyal* var. *fistula* (Schweinf.) Kyal. & Boatwr.; *Vachellia seyal* var. *seyal* (Delile) Hurter.

**Synonyms:** *Acacia fistula* Schweinf., *Acacia flava* (Forssk.) Schweinf. var. *seyal* (Delile) Roberty, *Acacia seyal* Delile, *Acacia stenocarpa* Rich.

Tree up to 9 m high with usually flattened crown; bark on trunk powdery, white to greenish yellow or orange red; young branchlets almost glabrous, epidermis reddish, conspicuously flaking off to expose a greyish or reddish powdery under-surface. Stipular spines up to 8 cm long, some sometimes basally inflated and fused into bilobed "ant-galls". Leaves: pinnae 3-7 pairs; leaflets 11-20 pairs, 3-8 x 0.7-1.5 mm. Flowers bright yellow, in heads; involucre in lower half of peduncle, often subtending a few additional flowers. Calyx 2-2.5 mm long. Corolla 3.5-4 mm long. Pods ± falcate and constricted between the seeds, dehiscent, 7-20 x 0.5-0.9 cm, finely longitudinally veined, glabrous. Seeds elliptic, compressed, 7-9 x 4.5-5 mm; areole 5-6 x 2.5-3.5 mm [Thulin *et al.*, 1993]

### Ethnobotanical Potential

According to some Biblical scholars, the Shittah tree is mentioned in the Bible only once (I will plant in the wilderness, The Shittah tree. Isaiah 41), but its wood is referred to many times as shittium, which is the plural of shittah in Hebrew. Some even speculate that it was only natural that Moses should turn to shittium when he came to build the Ark of the Covenant and the Tabernacle and needed beams and timber. No one can really be sure which species of *Acacia* was meant.

Wood is white to yellow-brown, finely-striated with dark lines, coarse-grained, soft, easy to work, polishes well, but discolours easily with mold and is susceptible to insect attack.

Ancient Egyptians made coffins, some still intact, from the wood.

Nigerians used sapling stems, or also the roots for spear shafts.

Tree also yields a gum of good quality, inferior to that of *A. senegal*. Systematic tapping has produced a product of better colour and taste.

Bark contains tannin and yields a red liquid extract.

The gum is said to be edible. Eating the gum is supposed to afford some protection against bronchitis and rheumatism (Duke, 1983a).

The leaves are important for forage and the wood for fuel where the trees are abundant.

In parts of Africa the tree is important for livestock, natives driving their animals to where it is common and lopping off branches for them, both leaves and young pods being eaten.

The pods are sold, especially for fattening sheep. The tree is believed to provide the best firewood in Chad, and the best fodder in Sahelian savannas (NAS, 1980a; Duke, 1983a).

It is regarded by the Mbeere as a major bee forage; barrel hives hung in the tree and afforded protection by the very thorny branches from predation by the nratel or honey badger.

The *Acacia seyal* var. *seyal* asan important honey source; honey-white, very mild aroma.

#### *Ethnomedicinal Potential*

The bark is used to treat dysentery and bacterial infections of the skin, such as leprosy. The bark is also used as a stimulant [Purdue University].

The gum is used as an aphrodisiac, to treat diarrhoea, as an emollient, to treat haemorrhaging, inflammation of the eye, intestinal ailments and rhinitis. The gum is used to ward off arthritis and bronchitis [Purdue University].

Incense from the wood is used to treat pain from rheumatism and to keep expectant mothers from contracting rhinitis and fevers [Purdue University].

#### *Folk Medicine*

The gum is believed to be aphrodisiac.

The bark decoction is used for dysentery and leprosy.

Tanganyikans use the bark as a stimulant in tropical Africa.

The gum is used as emollient and astringent for colds, diarrhoea, haemorrhage and ophthalmia.

Mixed with *Acacia sieberana* DC, it is used for intestinal ailments on the Ivory Coast.

Wood used as a fumigant for rheumatic pains, and to protect puerperal mothers from colds and fevers. Eating the gum is supposed to afford some protection against bronchitis and rheumatism (Duke, 1983a).

*Vachellia seyal* is, along with other *Vachellias*, an important source for gum arabic, a natural polysaccharide, that exudes from damaged stems and solidifies [Purdue University]. The gum of

*Vachellia seyal* is called gum talha, from the Arabic name of the tree: (Talh).

Parts of the tree have a tannin content of up to 18-20%. The bark and seed pods of *Vachellia seyal* var. *seyal* have a tannin content of about 20% [www.fao.org].

Wood from the tree is said to have been used in Ancient Egypt to make coffins and also the Ark of the Covenant [*Vachellia seyal*].

**4. Alhagi maurorum** Shook, Aqool, ShookEl Jamal, Shprim, Lehlah; Camel thorn bush, Caspian manna, Persian manna; Alhagi des Maures, Kameldorn, Manna-, Mannastrauch; Bharbhara, Jawasa; Lupinellaalhagi, Manna di Persia; Kameeldoringbos, Volstruisdoring [AG Medicinal plants].

**Synonyms:** *Alhagigraecorum*, *Alhagicamelorum* Fisch., *Alhagipersarum* Boiss. & Buhse, *Alhagipseudalhagi*, *Alhagipseudalhagi* (M. Bieb.) Fisch., *Hedysarumalhagi* L., *Hedysarum pseudalhagi* M. Bieb., *Alhagimaurorum* Medik. subsp. *maurorum*, *Alhagipseudalhagi* subsp. *persarum* (Boiss. & Buhse) Takht., *Alhagicamelorum* var. *spinis-elongatis* Boiss [EOL, 2014; USDA, NRCS, 2010; TF Pakista, 2011; Euromed Plant data base].

It is a deep rooted, rhizomatous, perennial shrub, with roots that can extend six to seven feet into the ground. The spiny, intricately-branched shrub reaches 1.5 to 4 feet in height. The plant, which is grayish green and hairless, has simple, entire leaves that are alternately arranged. The leaf shape is oval to lance-shaped. The small pea-like flowers are pinkish purple to maroon and are borne on short, spine-tipped branches that arise from the leaf axils. The reddish-brown to tan fruits are found between the seeds, with a short narrow beak at the end [A G Medicinal plants; *Alhagi maurorum*, 2012].

#### *Ethnobotanical Potential*

All plant parts including the roots are being used in medicines [Ali Esmail Al-Snafi, 2015].

*Alhagi maurorum* plant is grazed by livestock.

It is cut in late spring and used for making hay for small livestock and camels [Al-Snafi, 2013 and Bhandari, 2013].

Manna, a sugar exudate, is formed on stems and leaves and shaken from the bushes at flowering.

In Indian markets it is sold under the name (torajabin) and is imported from Afghanistan and Iran.

Today, manna is used for extracting mannitol,

made into tablets and used in the cosmetic and pharmaceutical industries to produce laxatives, diuretics and sweeteners [Brandis, 1972; Bamber, 1916; Maheshwari, 1963 and Thalen, 1979]. It composed from monomeric units mainly consisting of galactose and uronic acids [Goncharov *et al.*, 2001].

#### *Ethnomedicinal Potential*

The previous studies showed that *Alhagimaurorum* contained many secondary metabolites including flavonoids, fatty acids, coumarins, glycosides, sterols, steroids, resins, vitamins, alkaloids, carbohydrates, tannins, unsaturated sterols and triterpenes. It exerted anti-bacterial, anti-inflammatory, antipyretic, analgesic, antioxidant, gastrointestinal, cardiovascular, diuretic, and dermatological and many other effects [Ali Esmail Al-Snafi, 2015].

*Alhagi maurorum* is customarily used in folk medicine as a remedy for rheumatic pains, bilharziasis, liver disorders, various types of gastrointestinal discomfort, general tonic, anthelmintic, to treat constipation, jaundice and arthritis.

It also used as diuretic, blood purifier, antimicrobial, for treatment of dysentery, upper respiratory system problems, wounds, haemorrhoids and uterine problems. The roots were used as aphrodisiac [EMP, 2005 and Chakravarti, 1976].

The plant is used as laxative, diuretic and expectorant in India. The oil extracted from leaves is used for curing rheumatism [Singh *et al.*, 1990]. A decoction made from seeds of *Alhagi maurorum* is used for curing kidney stones [Fahmy, 1963].

#### **5. *Allium cepa* Linn. Common onion, Bulb onion, (Liliaceae, Amaryllidaceae):**

*Synonyms: Allium cepa var. aggregatum* G. Don, *Allium cepa var. bulbiferum* Regel, *Allium cepa var. cepa* Linn., *Allium cepa var. multiplicans* Bailey, *Allium cepa var. proliferum* (Moench) Regel, *Allium cepa var. solaninum* Alef, *Allium cepa var. viviparum* (Metz) Mansf. [Linnaeus, 1753 and USDA Plant list].

The onion (*Allium cepa* L., from Latin *cepa* "onion"), is a vegetable that is the most widely cultivated species of the genus *Allium*. Its close relatives include the garlic, leek, chive, [The Plant List and Chinese onion Block, 2013].

This genus also contains several other species variously referred to as onions and cultivated for food, such as the Japanese bunching onion (*Allium*

*fistulosum*), the tree onion (*A. ×proliferum*), and the Canada onion (*Allium canadense*). The name "wild onion" is applied to a number of *Allium* species, but *A. cepa* is exclusively known from cultivation. Its ancestral wild original form is not known, although escapes from cultivation have become established in some regions [Allergy Net, 2010]. The onion is most frequently a biennial or a perennial plant, but is usually treated as an annual and harvested in its first growing season. The onion plant has been grown and selectively bred in cultivation for at least 7,000 years. It is a biennial plant, but is usually grown as an annual. Modern varieties typically grow to a height of 15 to 45 cm (6 to 18 in). The leaves are yellowish- to bluish green and grow alternately in a flattened, fan-shaped swathe. They are fleshy, hollow, and cylindrical, with one flattened side. They are at their broadest about a quarter of the way up, beyond which they taper towards a blunt tip. The base of each leaf is a flattened, usually white sheath that grows out of a basal disc. From the underside of the disc, a bundle of fibrous roots extends for a short way into the soil. As the onion matures, food reserves begin to accumulate in the leaf bases and the bulb of the onion swells [Fritsch and Friesen, 2002].

In the autumn, the leaves die back and the outer scales of the bulb become dry and brittle, so the crop is then normally harvested. If left in the soil over winter, the growing point in the middle of the bulb begins to develop in the spring. New leaves appear and a long, stout, hollow stem expands, topped by a bract protecting a developing inflorescence. The inflorescence takes the form of a globular umbel of white flowers with parts in sixes. The seeds are glossy black and triangular in cross section [Fritsch and Friesen, 2002]. The average pH of an onion is around 5.5 [Brickell, 1996].

The onion plant has a fan of hollow, bluish-green leaves and its bulb at the base of the plant begins to swell when a certain day-length is reached. The bulbs are composed of shortened, compressed, underground stems surrounded by fleshy modified scale (leaves) that envelop a central bud at the tip of the stem. In the autumn (or in spring, in the case of overwintering onions), the foliage dies down and the outer layers of the bulb become dry and brittle. The crop is harvested and dried and the onions are ready for use or storage. The crop is prone to attack by a number of pests and diseases, particularly the onion fly, the onion eelworm, and various fungi cause rotting. Some varieties of *A. cepa*, such as shallots and potato onions, produce multiple bulbs.

Onions are cultivated and used around the world.

As a food item, they are usually served cooked, as a vegetable or part of a prepared savoury dish, but can also be eaten raw or used to make pickles or chutneys. They are pungent when chopped and contain certain chemical substances which irritate the eyes.

#### *Ethnobotanical Potential*

Onions are commonly chopped and used as an ingredient in various hearty warm dishes, and may also be used as a main ingredient in their own right, for example in French onion soup, creamed onions, and onion chutney. They are versatile and can be baked, boiled, braised, grilled, fried, roasted, sautéed, or eaten raw in salads [Smith, 2013]. Their layered nature makes them easy to hollow out once cooked, facilitating stuffing them, as in Turkish sogan-dolma.

Onions pickled in vinegar are eaten as a snack around the world, and as a side serving in pubs and fish and chip shops throughout the United Kingdom and the Commonwealth. They are part of a traditional British pub's ploughman's lunch, usually served with crusty bread, English cheese, and ale.

Similar to garlic [Onion Good food, 2013] onions can show an additional colour – pink-red – after cutting, an effect caused by reactions of amino acids with sulphur compounds [Lukes, 1986].

Homeopathic remedies are made from plants, but they also come from some of the foods we eat. Perhaps the most well-known, food-based homeopathic remedy is made from the common red onion and is called *Allium cepa*.

The onion, also called the common onion or bulb onion, is the most cultivated species of the *Allium* genus, and it is related to the leek, garlic, and shallot. Like its other *Allium* family members, onions contain important health-promoting sulphur phytonutrients, such as allyl propyl sulfoxides. As a result, eating onions help boost immunity, treat infections, lower cholesterol, reduce blood pressure, reduce the risk of heart disease, and maintain blood sugar. Onions have particularly large cells that are readily observed under low magnification. Forming a single layer of cells, the bulb epidermis is easy to separate for educational, experimental, and breeding purposes [Lee *et al.*, 2012 and Suslov *et al.*, 2009].

Onions are, therefore, commonly employed in science education to teach the use of a microscope for observing cell structure [Xu *et al.*, 2014].

Onions are toxic to dogs, cats, guinea pigs, and many other animals [Mc Cabe, 2007 and Cope, 2005] evils.

Most onion cultivars are about 89% water, 9% carbohydrates (including 4% sugar and 2% dietary fibre), 1% protein, and negligible fat. Onions contain low amounts of essential nutrients and have an energy value of 166 kJ (40 Calories) in a 100 g (3.5 oz) amount. Onions contribute savoury flavour to dishes without contributing significant caloric content [Ansari, 2007].

#### *Ethnomedicinal Potential*

*Allium cepa*, is used for a wide variety of infections, cold symptoms, allergies, eye irritations, and neuralgic pains.

*Allium cepa* is often recommended for allergies or allergic rhinitis – also called hay fever. Common symptoms include watery eyes, a stuffy nose, sneezing, and a sore throat.

*Allium cepa* is useful for acute allergy symptoms, especially hay fever with profuse watering of the eyes, and catarrh that irritates and inflames the upper lip and nostrils. Symptoms improve from fresh air and a cool room but worsen from cold or damp weather, and warm rooms.

Let's look a little deeper on how *Allium cepa* helps a stuffy nose and catarrh. The homeopathic remedy is when profuse and watery catarrh also burns the skin of the nose and upper lip. The nose and lip will become raw, red, and very painful. The nose also drips like a faucet, and congestion will alternate between one nostril or both. The person will also experience constant violent sneezing.

Catarrh, a stuffy nose, and violent sneezing are also common symptoms of colds and the flu. As a result, *Allium cepa* is great for infections of all kinds, especially colds, the flu, and chest and throat infections. *Allium cepa* is also an appropriate remedy for sore throats, laryngitis, bronchitis, and a hacking cough that settles in the chest area.

People will have a sensation where the larynx is torn and split due to pain when they speak. Coughing worsens while inside in warm rooms, and during the night. Symptoms can also be brought on by cold air exposure.

Eye irritation is common for people that require *Allium cepa*. The eyes burn and will worsen from coughing. The eyes are also sensitive to light and are itchy and red. The person will also often want to rub their eyes, which will lead to further irritation. Vision will be impaired where near objects appear to be far away.

Neuralgia is a burning, stabbing, and quite often a severe pain that occurs due to a damaged nerve. The damaged nerve commonly occurs in the neck and face. *Allium cepa* is an excellent remedy for neuralgic pains that are accompanied by a burning and piercing pain that will shift from one side of the body to another. The pain may originate with a toothache in the molars, an earache, or a headache at the back of the brow.

The remedy is also given to people that endure phantom limb pains after an amputation. The pain will mainly occur in the face, head, teeth, sinuses, chest, or neck.

The list of possible side-effects that may occur in medicines that contain *Allium cepa* are viz.,

Dermatologic disease, Skin rash, Chronic blistering disease, Gastrointestinal disorder, Heartburn, Dyspepsia, Gastric acidity, Gastroesophageal reflux disease, Swelling, Eye ulceration.

*Allium cepa*, a spice plant, is commonly known as onion and belongs to the family Liliaceae. Since ancient times, it has been used traditionally for the treatment of different diseases. Among various activities of *Allium cepa*, regulation of hypoglycaemic activity is considered one of its important effects in DM. Sulphur compounds including S-methyl cysteine and flavonoids such as quercetin are mainly responsible for the hypoglycaemic activity of *Allium cepa*. S-methyl cysteine and flavonoids help to decrease the levels of blood glucose, serum lipids, oxidative stress, and lipid peroxidation, as well as increasing antioxidant enzyme activity and insulin secretion. Extracts of onion also have been shown to have hypoglycemic and hypolipidemic effects by normalizing the activities of liver hexokinase, glucose 6-phosphatase and HMG coenzyme-A reductase. In preliminarily clinical trials, patients with diabetes safely consumed slices of *Allium cepa*, exhibiting sufficient hypoglycemic activity [Akash *et al.*, 2014].

## References

1. A Guide to Medicinal Plants in North Africa. *Asphodelus tenuifolius* Cavan, 21-22.
2. Akash MS, Rehman K, Chen S. 2014. Spice plant *Allium cepa*: dietary supplement for treatment of type 2 diabetes mellitus. *Nutrition*. 2014; 30(10):1128-37.
3. *Alhagimaurorum*, <http://www.cabi.org/isc/datasheet/4215> [21 September 2012].
4. Ali Esmail Al-Snafi. *Alhagimaurorum* as a potential medicinal herb: An Overview *International Journal of Pharmacy Review & Research*. 2015;5(2): 130-136.
5. Allergy Net – Allergy Advisor Find. Allallergy.net. Archived from the original on 15 June 2010. Retrieved 14 April 2010.
6. *Allium cepa* L. World Checklist of Selected Plant Families (WCSP). Royal Botanic Gardens, Kew – via The Plant List.
7. *Allium cepa* Natural Resources Conservation Service PLANTS Database. USDA. Retrieved 2013-03-31.
8. Al-Snafi AE. Encyclopedia of the constituents and pharmacological effects of Iraqi medicinal plants. Thiqr University, 2013.
9. Ansari NA. Onion Cultivation and Production in Iran (PDF). *Middle Eastern and Russian Journal of Plant Science and Biotechnology*. 2007;1(2):26-38.
10. Ayoub SMH. TAN. A new molluscicide and algicide from the fruits of *Acacia nilotica*. *Journal of Chemical Technology and Biotechnology*, 1982;32(7):728-34.
11. Ayyanar, M., Sankarasivaraman, K., Ignacimuthu, S and Sekar, T. 2010. Plant species with ethno botanical importance other than medicinal in Theni district of Tamil Nadu, Southern India. *Asian J. Exp. Biol. Sci.* 2010;1(4):765-771.
12. Bamber CJ. *Plants of Punjab*. Punjab, India, Superintendent Government Printing, 1916.
13. Bhandari MM. *Flora of the Indian Desert*. Jodhpur, India: Scientific Publishers, 1978, 51.
14. Block E. *Garlic and Other Alliums: The Lore and the Science*. Royal Society of Chemistry. ISBN 0-85404-190-7.
15. Bolus L. *Medicinal Plants of North Africa*. Cairo, Egypt, Reference Publications Inc. 1983:368.
16. Brandis D. *The Forest-Flora of North-West and Central India*. Dehradun, India, Bishen Singh Mahendra Pal Singh, 1972.
17. Brickell, Christopher. *The Royal Horticultural Society Encyclopedia of Gardening*. Dorling Kindersley. 1992.p.345. ISBN 978-0-86318-979-1.
18. Carter JO. *Acacia nilotica*: a tree legume out of control. *Forage tree legumes in tropical agriculture.*, 1994;27:338-351.
19. Chakravarty A. *The Plant Wealth of Iraq (A Dictionary of Economic Plants)*. Botany Directorate, Ministry of Agriculture & Agrarian Reform, England, 1976.
20. Cope R.B. *Allium* species poisoning in dogs and cats. *Veterinary Medicine*. 2005 Aug;100(8):562-66. ISSN 8750-7943.
21. Duke J.A. *Medicinal plants of the Bible*. Trado-Medic Books, Owerri, NY. 1983a

22. Encyclopedia of medicinal plants in UAE. Health Authority Abu Dhabi. Zaied Center for Traditional Medicine and Herbs Researches 2005.pp.15-20.
23. EOL, Encyclopedia of the life, Alhagimaurorum 2014, <http://eol.org/pages/703028/names/synonyms>.
24. EuroMed plant database, Alhagimaurorum, ww2.bgbm.org/EuroPlusMed/PTaxonDetail.asp?NameId=63693&PTRefFk=8500000 [January 2011].
25. Fagg CW; Greaves A, 1990. *Acacia nilotica* 1869-1988. Annotated Bibliography - CAB International, No. F42:xiv + 81 pp.; [Published in collaboration with Oxford Forestry Institute (OFI)]; 632 ref.
26. Fahmy IR. Drug plants of Egypt. *Planta Medica*. 1963,11:202-224.
27. Fritsch R.M., N. Friesen. Chapter 1: Evolution, Domestication, and Taxonomy. In H.D. Rabinowitch; L. Currah. *Allium Crop Science: Recent Advances*. Wallingford, UK: CABI Publishing. 2002.pp.20-21. ISBN 0-85199-510-1.
28. Goldsmith B, Carter DT. The indigenous timbers of Zimbabwe. *Zimbabwe Bulletin of Forestry Research* 1981 No.9 pp.x + 406 pp. ref.22.
29. Goncharov MY, Yakovlev GP and Vitovskaya GA. Composition of polysaccharides from above-ground part of *Alhagimaurorum* Medic. *Rastitel'nye Resursy*. 2001;37:60-63.
30. Gupta RK. Multipurpose trees for agroforestry and wasteland utilisation. *Multipurpose trees for agroforestry and wasteland utilisation*. 1993 xv + 562 pp.
31. Harshberger J.W. The purpose of Ethnobotany. *Bot. Gaz.* 1896;21:146-158.
32. Jadhav V.D., Mahadkar S.D., Valvi, S.R. Documentation and ethnobotanical survey of wild edible plants from Kolhapur district. *Recent Research in Science and technology*, 2011;3(12): 58-63.
33. Le Houerou HN, 1988. *Acacia nilotica* (L.) Willd. Ex Del.
34. Lee Eunjin, Rezenom, Yohannes H., Russell David H., Patil Bhimanagouda S., Yoo Kil Sun. Elucidation of chemical structures of pink-red pigments responsible for 'pinking' in macerated onion (*Allium cepa* L.) using HPLC-DAD and tandem mass spectrometry. *Food Chemistry*. 2012;131(3):852-861.
35. Lemmens RHMJ; Wulijarni-Soetjipto N (Editors), *Plant resources of South East Asia*. No. 3. Dye and tannin-producing plants. 1991.pp.195.
36. Linnaeus, Carolus. *Species Plantarum* (in Latin). 1. Stockholm: Laurentii Sal vii. 1753.p.262.
37. Lukes, T.M. Factors Governing the Greening of Garlic Puree. *Journal of Food Science*. 1986;51(6):1577-77.
38. Maheshwari JK. *The Flora of Delhi*. New Delhi, India, CSIR, 1963.
39. Mahishi P., Srinivasa B.H. & Shivanna M.B. Medicinal plant wealth of local communities in some villages in Shimoga District of Karnataka, India. *J. Ethnopharmacol.* 2005;98:307-12.
40. McCabe A., Mick O'Donnell, Rachel Whittaker *Advances in Language and Education*. Bloomsbury Publishing. 2007.p.35. ISBN 978-1-4411-0458-8.
41. Ministry of Tribal Affairs, (2005-2006). Annual Report.
42. N.A.S. *Firewood crops. Shrub and tree species for energy production*. National Academy of Sciences, Washington, DC. 1980a.
43. National Academy of Sciences. *Firewood Crops: Shrub and Tree Species for Energy Production*. Washington DC, USA; National Academy of Sciences. 1980.
44. Onion. *Good Food*. BBC. Retrieved 2013-04-02.
45. Orwa C., Mutua A., Kindt R., Jamnadass R., Anthony S. *Agroforest tree Database: a tree reference and selection guide version 4.0*. World Agroforestry Centre, Kenya. 2009.
46. Pal D.C. (Eds. Singh, N.P., Singh, D.K., Hajra, P.K. and Sharma, B.D.M.) *Flora of India. Introductory volume - part II*. Botanical Survey of India, Calcutta, India, 2000.pp.303-20.
47. Patale C., Nasare K., Praveenkumar N, Narkhede S.D. Ethnobotanical studies on wild edible plants of Gond, Halba and Kawar tribes of Salekasa taluka, Gondia district, Maharashtra state, India. *Int. Res. J. Pharm.* 2015;6(8):512-18. <http://dx.doi.org/10.7897/2230-8407.06810>.
48. Rao RK, Purkayastha SK. *A. arabica* Willd. In: *Indian Woods, their identification, properties and uses*. Vol III. Leguminosae to Combretaceae. Delhi, India: Manager of Publications. 1972.pp.20-22.
49. Singh Y, Wadhvani AM and Johri BM. *Dictionary of Economic Plants of India*. New Delhi, India: Indian Council of Agricultural Research, 1990.
50. Smith S.E. What is onion powder. *Wise Geek. Conjecture Corporation*. Retrieved 2013-04-02.
51. Suslov D, Verbelen J.P., Vissenberg K. Onion epidermis as a new model to study the control of growth anisotropy in higher plants. *Journal of Experimental Botany*. 2009;60(14):4175-87.
52. Tewari MC. Rajput SS. *Acacia nilotica* - utilisation aspects. *Journal of the Timber Development Association of India*, 1987;33(2):28-32.
53. Thalen DCP. *Ecology and Utilization of Desert Shrub Rangelands in Iraq*. The Hague, The Netherlands: Dr. W. Junk, BV Publishers, 1979.
54. *The flora of Pakistan, Alhagimaurorum, Freshbiology*, 2011.
55. Thulin M., A.S. Hassan & B.T. Styles. *Acacia in Thulin, M. Flora Somalia, Vol 1, (1993) [updated by M. Thulin, 2008]*.
56. Trivedi P.C. and Sharma N. *Text book of Ethnobotany*, Pointer Publishers, Jaipur, India,

- 2011.
57. Troup RS, Joshi HB. The Silviculture of Indian Trees. Vol IV. Leguminosae. Delhi, India; Controller of Publications. 1983.
58. USDA, NRCS. The Plants Database. National Plant Data Center, Baton Rouge, LA, USA 2010.
59. Vachelliaseyal in BoDD - Botanical Dermatology Database.
60. Weber E. Invasive plant species of the world: A reference guide to environmental weeds. Wallingford, UK: CAB International, 2003.p.548. www.fao.org. Retrieved 2017-08-03.
61. Xu K, Huang X, Wu M, Wang Y, Chang Y, Liu K, Zhang J, Zhang Y, Zhang F, Yi L, Li T, Wang R, Tan G, Li C. A rapid, highly efficient and economical method of Agrobacterium-mediated in planta transient transformation in living onion epidermis. PLoS ONE. 2014;9(1):e83556.
-

## Revised Rates for 2019 (Institutional)

Title of the Journal	Frequency	India(INR)		Outside India(USD)	
		Print Only	Online Only	Print Only	Online Only
Dermatology International	Semiannual	5500	5000	430	391
Gastroenterology International	Semiannual	6000	5500	469	430
Indian Journal of Anatomy	Quarterly	8500	8000	664	625
Indian Journal of Anesthesia and Analgesia	Bi-monthly	7500	7000	586	547
Indian Journal of Cancer Education and Research	Semiannual	9000	8500	703	664
Indian Journal of Communicable Diseases	Semiannual	8500	8000	664	625
Indian Journal of Dental Education	Quarterly	5500	5000	430	391
Indian Journal of Diabetes and Endocrinology	Semiannual	8000	7500	597	560
Indian Journal of Genetics and Molecular Research	Semiannual	7000	6500	547	508
Indian Journal of Hospital Administration	Semiannual	7000	6500	547	508
Indian Journal of Hospital Infection	Semiannual	12500	12000	938	901
Indian Journal of Medical & Health Sciences	Semiannual	7000	6500	547	508
Indian Journal of Pathology: Research and Practice	Bi-monthly	12000	11500	938	898
Indian Journal of Preventive Medicine	Semiannual	7000	6500	547	508
International Journal of Neurology and Neurosurgery	Quarterly	10500	10000	820	781
International Physiology	Triannual	7500	7000	586	547
Journal of Cardiovascular Medicine and Surgery	Quarterly	10000	9500	781	742
Journal of Global Medical Education and Research	Semiannual	5900	5500	440	410
Journal of Global Public Health	Semiannual	12000	11500	896	858
Journal of Microbiology and Related Research	Semiannual	8500	8000	664	625
Journal of Organ Transplantation	Semiannual	26400	25900	2063	2023
Journal of Orthopedic Education	Triannual	5500	5000	430	391
Journal of Pharmaceutical and Medicinal Chemistry	Semiannual	16500	16000	1289	1250
Journal of Practical Biochemistry and Biophysics	Semiannual	7000	6500	547	508
Journal of Radiology	Semiannual	8000	7500	625	586
New Indian Journal of Surgery	Bi-monthly	8000	7500	625	586
Ophthalmology and Allied Sciences	Triannual	6000	5500	469	430
Otolaryngology International	Semiannual	5500	5000	430	391
Pediatric Education and Research	Quarterly	7500	7000	586	547
Physiotherapy and Occupational Therapy Journal	Quarterly	9000	8500	703	664
Urology, Nephrology and Andrology International	Semiannual	7500	7000	586	547
Indian Journal of Maternal-Fetal & Neonatal Medicine	Semiannual	9500	9000	742	703
Indian Journal of Obstetrics and Gynecology	Quarterly	9500	9000	742	703
Indian Journal of Emergency Medicine	Quarterly	12500	12000	977	938
Indian Journal of Trauma and Emergency Pediatrics	Quarterly	9500	9000	742	703
Journal of Emergency and Trauma Nursing	Semiannual	5500	5000	430	391
Indian Journal of Forensic Medicine and Pathology	Quarterly	16000	15500	1250	1211
Indian Journal of Forensic Odontology	Semiannual	5500	5000	430	391
Indian Journal of Legal Medicine	Semiannual	8500	8000	664	625
International Journal of Forensic Sciences	Semiannual	10000	9500	781	742
Journal of Forensic Chemistry and Toxicology	Semiannual	9500	9000	742	703
Community and Public Health Nursing	Triannual	5500	5000	430	391
Indian Journal of Surgical Nursing	Triannual	5500	5000	430	391
International Journal of Pediatric Nursing	Triannual	5500	5000	430	391
International Journal of Practical Nursing	Triannual	5500	5000	430	391
Journal of Gerontology and Geriatric Nursing	Semiannual	5500	5000	430	391
Journal of Nurse Midwifery and Maternal Health	Triannual	5500	5000	430	391
Journal of Psychiatric Nursing	Triannual	5500	5000	430	391
Indian Journal of Ancient Medicine and Yoga	Quarterly	8000	7500	625	586
Indian Journal of Law and Human Behavior	Semiannual	6000	5500	469	430
Indian Journal of Medical Psychiatry	Semiannual	8000	7500	625	586
Indian Journal of Biology	Semiannual	5500	5000	430	391
Indian Journal of Library and Information Science	Triannual	9500	9000	742	703
Indian Journal of Research in Anthropology	Semiannual	12500	12000	977	938
Indian Journal of Waste Management	Semiannual	9500	8500	742	664
International Journal of Political Science	Semiannual	6000	5500	450	413
Journal of Social Welfare and Management	Triannual	7500	7000	586	547
International Journal of Food, Nutrition & Dietetics	Triannual	5500	5000	430	391
Journal of Animal Feed Science and Technology	Semiannual	7800	7300	609	570
Journal of Food Additives and Contaminants	Semiannual	5000	4500	391	352
Journal of Food Technology and Engineering	Semiannual	5000	4500	391	352
Indian Journal of Agriculture Business	Semiannual	5500	5000	413	375
Indian Journal of Plant and Soil	Semiannual	6500	6000	508	469

**Terms of Supply:**

- Agency discount 12.5%. Issues will be sent directly to the end user, otherwise foreign rates will be charged.
- All back volumes of all journals are available at current rates.
- All Journals are available free online with print order within the subscription period.
- All legal disputes subject to Delhi jurisdiction.
- Cancellations are not accepted orders once processed.
- Demand draft / cheque should be issued in favour of "Red Flower Publication Pvt. Ltd." payable at Delhi
- Full pre-payment is required. It can be done through online (<http://rfppl.co.in/subscribe.php?mid=7>).
- No claims will be entertained if not reported within 6 months of the publishing date.
- Orders and payments are to be sent to our office address as given above.
- Postage & Handling is included in the subscription rates.
- Subscription period is accepted on calendar year basis (i.e. Jan to Dec). However orders may be placed any time throughout the year.

**Order from**

Red Flower Publication Pvt. Ltd., 48/41-42, DSIDC, Pocket-II, Mayur Vihar Phase-I, Delhi - 110 091 (India),

Mobile: 8130750089, Phone: 91-11-45796900, 22754205, 22756995 E-mail: [sales@rfppl.co.in](mailto:sales@rfppl.co.in), Website: [www.rfppl.co.in](http://www.rfppl.co.in)

## Consumer's Knowledge, Attitude and Practices (KAP) Regarding Organic Foods in Relation to Demographic Factors of Punjab

Seema Rani<sup>1</sup>, Neerja Singla<sup>2</sup>, Poonam Bakheta<sup>3</sup>

### How to cite this article:

Seema Rani, Neerja Singla, and Poonam Bakheta. Consumer's Knowledge, Attitude and Practices (KAP) Regarding Organic Foods in Relation to Demographic Factors of Punjab. *Int J Food Nutr Diet.* 2019;7(1):23-31.

### Abstract

The present study was carried out with an aim to study the knowledge, attitude and practices regarding organic foods on a total of 120 selected subjects (aged between 35 and 45 years) belonging to middle income group (MIG) and high income group (HIG) from two cities of Punjab namely Ludhiana and Patiala. The questionnaire consisted of various statements pertaining to knowledge, attitude and practice regarding organic foods. The major sources of information regarding organic foods were found to be television and radio, followed by newspaper, friends, organic shops, farmers and labels, values being 28.3, 18.3, 15.8, 15.8, 12.5 and 9.1%, respectively. Majority of the respondents (88%) strongly agreed that organic foods were healthier than conventional foods. Organic foods were found to be consumed by 96.7 and 93.3% percent of the MIG and HIG subjects respectively of Ludhiana city where as in Patiala city corresponding values were 83.3 and 90% respectively. There was no significant difference in the overall knowledge and attitude score of both MIG and HIG respondents. However, a statistically significant ( $p \leq 0.05$ ) difference was observed in the overall practice score of respondents belonging to MIG and HIG. Hence, it can be concluded that as the income increases the practice of using organic foods also increases.

**Keywords:** Attitude; Knowledge; Practice; Organic foods; Middle Income group; High Income group.

### Introduction

Healthy life style has become a global trend and has been underlined by a slogan 'back to nature'. The environment a lists in 1960s and 1970s concentrated on the effects of chemical fertilizers and pesticides on the biophysical and human health leading to

take an alternative food system i.e. organic farming which also hold into account consumer safety. The issues of soil quality, food contamination and conservation of biodiversity was highlighted by the organic food industry [6]. Though the organic products originated in the developed countries, but due to increasing awareness about environmental issues and their alarming effects on health, consumers from developing countries also started accepting the notion of "Go organic". Presently the consumers prefer healthy and high-quality food mainly for two reasons, which are food safety and sustainability. Therefore, the green foods with less chemical residues have become more popular worldwide [3].

Modernization of society has reportedly affected the food choices and preferences of consumers. The consumers very intelligently choose healthy foods because of their improved standard of living and

---

**Author Affiliation:** <sup>1</sup>Post-graduate student, <sup>2</sup>Scientist, <sup>3</sup>Nutritionist, Department of Food and Nutrition, Punjab Agricultural University, Ludhiana, Punjab 141004, India.

**Corresponding Author:** Neerja Singla, Scientist, Department of Food and Nutrition, Punjab Agricultural University, Ludhiana, Punjab 141004, India.

**E-mail:** [singlaneerja@pau.edu](mailto:singlaneerja@pau.edu)

education level. This habit of consumers emerged because there has been some apprehensions about the safety of conventional foods which ultimately influenced lifestyle and food habits worldwide [4]. The organic products are environment-friendly so far as the production methods and the final products are concerned.

The consumers demand bundles of characteristics in a product and their choice is influenced by their perception about desirable characteristics of organic foods. In organic market awareness and knowledge are important in two ways. Firstly, a large segment of the population is not yet informed about organic foods. Secondly, the consumers do not have enough detailed information to clearly differentiate the unique attributes of organic from conventionally grown alternatives [5]. The important drive for the growth of organic food market are consumer's knowledge and attitude as well as their practices. So, the present investigation was conducted with an objective to assess and compare the acquired knowledge, attitude and practices of the consumers regarding organic foods belonging to two different income groups i.e. MIG and HIG from two cities of Punjab.

### Material and Methods

For the present study, two cities namely Ludhiana and Patiala were selected which represent Central and South-eastern Zones of Punjab, respectively (in the year 2016-17). In each city two localities were selected on the basis of socio-economic status of the families. A total of one hundred and twenty subjects

including 60 from Ludhiana and 60 from Patiala city were selected, which comprised of 30 from Middle income group (MIG) and 30 from high income group (HIG) from both the cities (Fig. 1). The selection of subjects was independent of gender. The respondents selected from HIG of both the cities had monthly family income >Rs.1,00,000, while respondents from MIG had monthly family income between Rs.50,000-1,00,000.

The data regarding demographic profile of respondents pertaining to their gender (M/F), age, marital status, education, total family income, religion, type of family, number of family members and locality was recorded.

#### *Assessment of knowledge, attitude and practices (KAP) score*

A questionnaire consisting of two parts was developed. First part of the questionnaire contained all the general information of the subjects i.e. demographic profile, socio-economic status, education, marital status etc. Second part of the questionnaire contained a total of 29 questions out of which 10 were regarding the knowledge, 12 regarding attitude and 7 regarding the practice towards organic foods. The questionnaire was developed after thoroughly reviewing the available literature.

Pre-testing of the questionnaire was done on 20 subjects. Thereafter, necessary modifications were incorporated. These 20 respondents were excluded from the final respondents. The modified questionnaire was then used for data collection by personally interviewing the subjects.




Fig. 1: Selection of subjects

*Scoring of Knowledge, Attitude and Practices regarding organic foods*

Various statements (as mentioned in Table 3) to check the knowledge of the consumers regarding organic foods were asked and the responses were quantified by a score system from 3 to 1 as given in the following table:

Scoring of Knowledge regarding organic foods

Agreement	Scores
True	3
Do not know	2
False	1

Attitude of the selected respondents towards organic foods was assessed through various statements (as mentioned in Table 4) the results were calculated by a score system from 5 to 1 as mentioned below:

Scoring of Attitude towards organic foods

Agreement	Scores
Strongly agree	5
Agree	4
Neutral/No idea	3
Disagree	2
Strongly disagree	1

Statements regarding practices (as mentioned in Table 5) of the selected subjects towards organic foods were given scores as given below:

Table 1: General profile of the selected respondents (N=120)

Particulars		MIG			HIG		
		Ludhiana (n=30)	Patiala (n=30)	Total (n=60)	Ludhiana (n=30)	Patiala (n=30)	Total (n=60)
Age (years)	35-45	18 (60.0)	23 (76.7)	41 (68.3)	16 (53.3)	19 (63.3)	35 (58.3)
	>45	12 (40.0)	7 (23.3)	19 (31.7)	14 (46.7)	11 (18.3)	25 (41.7)
Gender	Female	26 (86.7)	28 (93.3)	54 (90.0)	26 (86.7)	26 (86.7)	52 (86.7)
	Male	4 (13.3)	2 (6.7)	6 (10.0)	4 (13.3)	4 (13.3)	8 (13.3)
Marital Status	Married	28 (93.3)	29 (96.7)	57 (95.0)	27 (90.0)	28 (93.3)	55 (91.7)
	Unmarried	2 (6.7)	1 (3.3)	3 (5.0)	3 (10.0)	2 (6.7)	5 (8.3)
Education	Primary	1 (3.3)	0 (0.0)	1 (1.7)	0 (0.0)	0 (0.0)	0 (0.0)
	High school	5 (16.7)	2 (6.7)	7 (11.7)	2 (6.7)	0 (0.0)	2 (3.3)
	Graduate	15 (50.0)	15 (50.0)	30 (50.0)	18 (60.0)	12 (40.0)	30 (50.0)
	Post graduate	9 (30.0)	13 (43.3)	22 (36.7)	10 (33.3)	17 (56.7)	27 (45.0)
Religion	Hindu	18 (60.0)	13 (43.3)	31 (51.7)	14 (46.7)	13 (43.3)	27 (45.0)
	Sikh	11 (36.7)	17 (56.7)	28 (46.7)	16 (53.3)	16 (53.3)	32 (53.3)
	Others	1 (3.3)	0 (0.0)	1 (1.67)	0 (0.0)	1 (3.3)	1 (1.7)
Type of family	Joint	7 (23.3)	6 (20.0)	13 (21.7)	4 (20.0)	10 (33.3)	14 (23.3)
	Nuclear	23 (76.7)	24 (80.0)	47 (78.3)	26 (80.0)	20 (66.7)	46 (76.7)
Type of residence	Owned	18 (60.0)	27 (90.0)	45 (75.0)	30 (100.0)	30 (100.0)	60 (100.0)
	On rent	12 (40.0)	3 (10.0)	15 (25.0)	0 (0.0)	0 (0.0)	0 (0.0)
Occupation	Working	22 (73.3)	21 (70.0)	43 (71.7)	28 (93.3)	26 (86.7)	54 (90.0)
	Non- working	8 (26.7)	9 (30.0)	17 (28.3)	2 (6.7)	4 (13.3)	6 (10.0)
Total monthly income	50,000-1,00,000	30 (100)	30 (100)	60 (100)	0 (0.0)	0 (0.0)	0 (0.0)
	>1,00,000	0 (0.0)	0 (0.0)	0 (0.0)	30 (100)	30 (100)	60 (100)

Figures in parenthesis indicate percentage

Scoring of Practices towards organic foods

Agreement	Scores
Yes	1
No	0
Organic food consumption frequency scores	
Daily	4
Weekly	3
Monthly	2
Rarely	1

*Statistical analysis*

The collected data was analyzed using various statistical tools such as frequency, percentages and the comparisons between categories of respondents were done using z-test and correlation coefficient of various demographic and socio-economic factors were also calculated.

**Results and Discussion**

*General information*

The general information of the respondents i.e. age, gender, marital status, education, religion, type of family, type of residence, occupation and income of the respondents has been presented in Table 1. As evident from Table 1 the age of respondents of both the cities belonging to MIG as well as HIG ranged between 35-45 years. Further, gender distribution showed that 90% of the respondents

in MIG and 86.7% in HIG were females. Majority of the respondents were married in MIG and HIG i.e. 95 and 91.7% respectively. In both the income group from both the cities 50% of the selected respondents were found to be graduates while 45% of the HIG subjects and 36.7% of the MIG subjects were educated up to postgraduate level indicating good educational level of the subjects. The majority of the selected respondents in both the cities were Hindu/Sikh. The majority of the selected respondents i.e. 78.3% in MIG and 71.7% in HIG belonged to nuclear families. All the subjects in the HIG had their own houses while 75% MIG had their own houses, while others had rented houses. It was also found that 71.7 and 90% selected respondents from MIG and HIG, respectively were working. All the subjects in MIG were having monthly income between Rs. 50,000-1 lakh and in HIG all the subjects were having monthly income more than Rs. 1,00,000.

#### *Source of information regarding organic foods*

As evident from Table 2, 28.3% of the subjects from both the cities found TV/Radio as the major source of information followed by newspaper/magazines (18.3%), friends/relatives (15.8%), organic shops (15.8%), farmers (12.5%) and labels (9.1%), respectively. It was reported [2] that 28.5% of the respondents used television and radio for obtaining information about organic food products where as newspapers, friends, farmers and other type of information delivery was 14.9, 19.7, 14.3 and 22.6%, respectively. Quality information, labeling and advertisement play important role in enhancement of knowledge [9].

#### *Knowledge, attitude and practices towards organic foods*

##### *Knowledge about organic foods*

Knowledge can be defined as familiarity or understanding of something such as facts, information and description etc. In the present

study knowledge of the respondents was assessed through a series of questions about organic foods and they were asked to state whether the statement were true/false or they do not know. The statement "organic production does not involve use of chemical pesticides and fertilizers" was found to be true by 44% of the total subjects in middle income group of both the cities where as 57% of the higher income group subjects of both the cities found this statement to be true. The organic foods have higher nutrient content and healthier than conventional foods, this statement was found to be true by majority of the respondents of both the income group in both cities (Table 3). The values being 70 and 71.7% respectively in middle and high income group of both the cities. Among the middle income group 51.7% subjects of both the cities found the statement that organic foods can be judged by their appearance as false, where as 41.7% of the high income group subjects found this statement to be true.

Statement regarding no preservatives in organic foods was found to be true by 43.3 and 48.3% from MIG and HIG respectively. However, 35% of the respondents from MIG did not know about presence or absence of preservatives in organic foods. Fifty percent of the respondents from MIG knew that bio fertilizers are used in organic foods where as the corresponding values in HIG was 47.7% for the same statement. Regarding the availability of organic foods majority of the subjects from both the income groups i.e. 46.7 and 58.3% from MIG and HIG respectively stated it false that real organic products can be purchased only from supermarkets, departmental stores or organic food stores. Among MIG subjects 43.3% of the respondents reported it true that only certified and labeled foods are real organic foods. However, 51.7% of the HIG subjects found the same statement to be false. The statement that organic farming can serve as an effective measure for the safety and health of the people was found to be true by a great majority of the subjects in both the income groups (Table 3).

**Table 2:** Source of information regarding organic foods among middle and high income group subjects of both the cities.

Source of information	MIG (n=60)	HIG(n=60)	Total (N=120)	z-value
Television/Radio	18(30.0)	16(26.7)	34(28.3)	0.41
Newspaper/magazines	10(16.7)	12(20.0)	22(18.3)	0.47
Friends/Relatives	10(16.7)	9(15.0)	19(15.8)	0.25
Organic shops	9(15.0)	10(16.7)	19(15.8)	0.25
Farmers	7(11.7)	8(13.3)	15(12.5)	0.28
Labels	6(10.0)	5(8.3)	11(9.1)	0.32

Figures in parenthesis indicate percentage

**Table 3:** Knowledge regarding organic foods among middle and high income group subjects of selected cities (N=120)

Statements		MIG (n=60)			HIG (n=60)		
		Do not know	False	True	Do not know	False	True
Organic production does not apply chemical pesticide and fertilizers.	Ludhiana	3 (10.0)	2 (6.67)	25 (83.3)	1 (3.3)	1 (3.3)	28 (93.3)
	Patiala	5 (16.7)	6 (20.0)	19 (63.3)	0 (0.0)	1 (3.3)	29 (96.7)
	z-value	0.76	1.52	1.75	1.01	0	0.59
	Total	8 (13.3)	8 (13.3)	44 (73.3)	1 (1.7)	2 (3.3)	57 (95.0)
Organic food are grown under natural conditions.	Ludhiana	2 (6.67)	26 (86.7)	2 (6.67)	2 (6.7)	27 (90.0)	1 (3.3)
	Patiala	5 (16.7)	20 (66.7)	5 (16.7)	1 (3.3)	23 (76.7)	6 (20.0)
	z-value	1.21	1.83	1.21	0.59	1.39	2.01*
	Total	7 (11.7)	46 (76.7)	7 (11.7)	3 (5.0)	50 (83.3)	7 (11.7)
Organic food has higher nutrition content than conventional food.	Ludhiana	3 (10.0)	0 (0.00)	27 (90.0)	12 (40.0)	3 (10.0)	15 (50.0)
	Patiala	10 (33.3)	5 (16.7)	15 (50.0)	1 (3.3)	1 (3.3)	28 (93.3)
	z-value	2.19*	2.34*	3.38**	3.45**	1.04	3.72**
	Total	13 (21.7)	5 (8.3)	42 (70.0)	13 (21.7)	4 (6.7)	43 (71.67)
We can judge if a product is organic or not by its outlook appearance.	Ludhiana	7 (23.3)	15 (50.0)	8 (26.7)	6 (20.0)	9 (30.0)	15 (50.0)
	Patiala	7 (23.3)	16 (53.3)	7 (23.3)	6 (20.0)	14 (46.7)	10 (33.3)
	z-value	0	0.26	0.3	0	1.33	1.31
	Total	14 (23.3)	31 (51.7)	15 (25.0)	12 (20.0)	23 (38.3)	25 (41.7)
There is no preservative in organic food.	Ludhiana	12 (40)	5 (16.7)	13 (43.3)	6 (20)	11 (36.7)	13 (43.3)
	Patiala	9 (30)	8 (26.7)	13 (43.3)	8 (26.7)	6 (20)	16 (53.3)
	z-value	0.81	0.94	0	0.61	1.43	0.78
	Total	21 (35.0)	13 (21.7)	26 (43.3)	14 (23.3)	17 (28.3)	29 (48.3)
Organic farming use bio-fertilizers.	Ludhiana	19 (63.3)	2 (6.7)	9 (30.0)	8 (26.7)	9 (30.0)	13 (43.3)
	Patiala	5 (16.7)	4 (13.3)	21 (70)	9 (30)	6 (20)	15 (50)
	z-value	3.69**	0.86	3.10**	0.29	0.89	0.52
	Total	24 (40.0)	6 (10.0)	30 (50.0)	17 (28.3)	15 (25.0)	28 (46.7)
Real organic products can only be bought in large supermarkets, department stores or organic food stores.	Ludhiana	1 (3.3)	13 (43.3)	16 (53.3)	4 (13.3)	21 (70.0)	5 (16.7)
	Patiala	6 (20)	15 (50)	9 (30)	3 (10)	14 (46.7)	13 (43.3)
	z-value	2.01*	0.52	1.83	0.4	1.83	2.25*
	Total	7 (11.67)	28 (46.7)	25 (41.7)	7 (11.7)	35 (58.3)	18 (30.0)
Only those products with organic certified labels are real organic products.	Ludhiana	2 (6.7)	11 (36.7)	17 (56.7)	3 (10)	16 (53.3)	11 (36.7)
	Patiala	9 (30)	12 (40)	9 (30.0)	5 (16.7)	15 (50)	10 (33.3)
	z-value	2.34*	0.27	2.08*	0.76	0.26	0.27
	Total	11 (18.3)	23 (38.3)	26 (43.3)	8 (13.3)	31 (51.7)	21 (35)
Organic farming can serve as an effective measure for safety and health of the people.	Ludhiana	1 (3.3)	2 (6.7)	27 (90)	3 (10)	1 (3.3)	26 (86.7)
	Patiala	4 (13.3)	2 (6.7)	24 (80)	2 (6.7)	2 (6.7)	26 (86.7)
	z-value	1.4	0	1.08	0.47	0.59	0
	Total	5 (8.3)	4 (6.7)	51 (85)	5 (8.3)	3 (5.00)	52 (86.7)
Several organic foods are available in the market.	Ludhiana	1 (3.3)	7 (23.3)	22 (73.3)	4 (13.3)	0 (0.0)	26 (86.7)
	Patiala	12 (40.0)	2 (6.7)	16 (53.3)	3 (10)	2 (6.7)	25 (83.3)
	z-value	3.45**	1.81	1.61	0.4	1.44	0.36
	Total	13 (21.7)	9 (15)	38 (63.3)	7 (11.7)	2 (3.3)	51 (85)

Figures in parenthesis indicate percentage

\* Significant at 5% level

\*\* Significant at 1% level

The values being 85 and 86.7% among MIG and HIG, respectively. A higher number of the subjects from HIG i.e. 85% stated that several organic foods are available in the market as true.

#### *Attitude towards organic foods*

An attitude is a tendency to respond in a positive or in a negative manner towards a certain idea, object, person or situation. It influences one's choice of action and response to challenges and rewards. In the present study attitude of the selected subjects towards organic foods was assessed by a well structured questionnaire which consisted of 12 statements and the subjects were asked to respond whether they agreed, disagreed or neutral to the mentioned statement. It was found that 46.7% of the MIG subjects and 50% of HIG subjects from Ludhiana city strongly disagreed to the statement that "I am not interested in organics" (Table 4). However, 50% of HIG subjects from Patiala city were neutral to the above mentioned statement. The statement regarding preference of buying organic foods was agreed by 36.7% of the MIG subjects in Ludhiana city where as in Patiala city half of the subjects disagreed to the same statement, where as 26.7% of the higher income group subjects in Ludhiana city and 30% in Patiala city agreed to the same statement. Among the MIG subjects of Ludhiana and Patiala city who agreed to the statement regarding safety and reliability of organic foods was 56.7 and 60% respectively and 50% HIG subjects in Ludhiana city and 30% subjects in Patiala city strongly agreed to the same statement. Similarly, the statement regarding better quality of organic foods was agreed upon by 53.3 and 56.7% of the MIG subjects and 50 and 33.3% of HIG subjects in Ludhiana and Patiala city, respectively. No subjects in both income groups of both the cities disagreed to the statement that "organic products are healthier and more nutritious i.e. all the subjects were aware that organic foods are healthier. It has been reported that majority of the subjects about 88% t strongly agreed that organic foods were healthier than conventional foods and only 2% of consumers disagreed to the statement (Table 4). It was further reported that 10% of the respondents were neutral in this respect.

All the MIG subjects from Ludhiana city and 3.3% in Patiala city strongly agreed to the statement and fifty percent HIG subjects remained neutral to the statement that organic foods are supreme products and consumed only by rich people. Nooripoor *et al.*, (2008) indicated that higher price of organic foods is one of the major barrier to their consumption. All the subjects in both income groups of both the

cities strongly agreed to the statement that organic products are available in limited variety in the market. Majority of the respondents i.e. 43.3% in MIG and 40% in HIG in Ludhiana city agreed to the statement that there are a lot of sale points for organic products, where as 50% in MIG and 46.7 in HIG of Patiala city were neutral to the same statement. The statement regarding trust on organic products was agreed upon by 56.7 and 43.3% of the selected MIG subjects from Ludhiana and Patiala city, respectively. However, 46.7% t subjects of Ludhiana city and 40% subjects of Patiala city remained neutral to the statement.

Further, it was also found that all the subjects agreed that organic foods are more nutritious but at the same time all of them agreed that these foods are costlier. Majority of the subjects agreed that organic foods are available in limited variety. On contrary, it was found that a higher number of respondents in Ludhiana city were interested in organic foods and preferred to purchase organic foods as compared to those in Patiala city. The reason for this difference might be that though the respondents in both the cities were well aware about the better quality, safety and nutrition of organic foods but the respondents from Ludhiana were more interested in purchasing organic foods, which might be due to their higher purchasing power and more availability of organic foods in Ludhiana city.

#### *Practices towards organic foods*

The practice is the actual application or use of a method/idea/belief as opposed to theories relating to it. Along with the knowledge and attitude, practices towards organic foods adopted by the respondents were also assessed in middle and high income group subjects of both the selected cities. Organic foods were found to be consumed by 96.7 and 93.3% of the MIG and HIG subjects of Ludhiana city respectively, where as in Patiala city values were 83.3 and 90% respectively, indicating that in Ludhiana city income level of selected subjects did not influence their purchase of organic foods, but in Patiala city the subjects from HIG were purchasing more of organic foods (Table 5). Among MIG subjects of both the cities 90% of the respondents stated that they were consuming organic foods while HIG subjects 93.3% of the subjects responded the same. This finding can be correlated well with the attitude of the respondents from both the cities which indicated that more number of subjects in Ludhiana city were interested in purchasing organic foods and all the subjects in both the cities responded that organic foods were costlier.

**Table 4:** Attitude regarding organic foods among middle and high Income group subjects of selected cities (N=120)

Statements		MIG (n=60)					HIG (n=60)				
		Strongly Agree	Agree	Neutral	Disagree	Strongly Disagree	Strongly Agree	Agree	Neutral	Disagree	Strongly Disagree
I am not interested in organic foods	Ludhiana	0 (0.0)	2 (6.7)	9 (30.0)	8 (26.7)	14 (46.7)	0 (0.0)	2 (6.7)	10 (33.3)	3 (10)	15 (50)
	Patiala	1 (3.3)	4 (13.3)	15 (50.0)	7 (23.3)	3 (10)	0 (0.0)	3 (10.0)	11 (36.7)	2 (6.7)	14 (46.7)
	z-value	1.01	0.86	1.58	0.51	3.83**	NA	0.47	0.31	0.52	0.31
	Total	1 (1.7)	6 (10)	24 (40)	15 (25)	17 (28.3)	NA	5 (8.3)	21 (35)	5 (8.3)	29 (48.3)
I prefer buying organic foods.	Ludhiana	0 (0.0)	11 (36.7)	9 (30)	7 (23.3)	3 (10)	11 (36.7)	8 (26.7)	8 (26.7)	3 (10)	0 (0.0)
	Patiala	3 (10)	7 (23.3)	1 (3.3)	15 (50)	4 (13.3)	8 (26.7)	9 (30)	12 (40)	1 (3.3)	0 (0.0)
	z-value	1.78	1.13	2.77**	2.14*	0.40	0.83	0.29	1.10	1.04	NA
	Total	3 (5.0)	18 (30.0)	10 (16.7)	22 (36.7)	7 (11.7)	19 (31.7)	17 (28.3)	20 (33.3)	4 (6.7)	0 (0.0)
Organic products are safer and more reliable.	Ludhiana	0 (0.0)	17 (56.7)	4 (13.3)	1 (3.3)	8 (26.7)	15 (50.0)	11 (36.7)	4 (13.3)	0 (0.0)	0 (0.0)
	Patiala	2 (6.7)	18 (60)	8 (26.7)	2 (6.7)	0 (0.0)	9 (30.0)	10 (33.3)	11 (36.7)	0 (0.0)	0 (0.0)
	z-value	1.44	0.26	1.29	0.59	3.04**	1.58	0.27	2.09*	NA	NA
	Total	2 (3.3)	35 (58.3)	12 (20.0)	3 (5.0)	8 (13.3)	24 (40.0)	21 (35)	15 (25.0)	0 (0.0)	0 (0.0)
Organic products are of better quality.	Ludhiana	0 (0.0)	16 (53.3)	4 (13.3)	3 (10.0)	7 (23.3)	7 (23.3)	15 (50)	8 (26.7)	0 (0.0)	0 (0.0)
	Patiala	1 (3.3)	17 (56.7)	7 (23.3)	5 (16.7)	0 (0.0)	10 (33.3)	10 (33.3)	9 (30.0)	1 (3.3)	0 (0.0)
	z-value	0.21	0.26	1.00	0.76	2.82**	0.86	1.31	0.29	1.01	NA
	Total	1 (1.7)	33 (55)	11 (18.3)	8 (13.3)	7 (11.7)	17 (28.3)	25 (41.7)	17 (28.3)	1 (1.7)	0 (0.0)
Organic products are healthier and more nutritious	Ludhiana	7 (23.3)	15 (50.0)	6 (20.0)	2 (6.7)	0 (0.0)	6 (20.0)	4 (13.3)	15 (50.0)	5 (6.7)	0 (0.0)
	Patiala	4 (13.3)	12 (40.0)	9 (30.0)	5 (16.7)	0 (0.0)	7 (23.3)	6 (20.0)	17 (56.7)	0 (0.0)	0 (0.0)
	z-value	1.07	0.78	0.89	1.21	NA	0.31	0.69	0.52	2.34*	NA
	Total	11 (6.7)	27 (45.0)	15 (25.0)	7 (11.6)	0 (0.0)	13 (21.7)	10 (16.7)	32 (23.3)	5 (8.3)	0 (0.0)
Organic products are very expensive.	Ludhiana	0 (0.0)	11 (36.7)	6 (20.0)	6 (20.0)	7 (23.3)	13 (43.3)	7 (23.3)	9 (30.0)	1 (3.3)	0 (0.0)
	Patiala	0 (0.0)	12 (40.0)	13 (43.3)	5 (16.7)	0 (0.0)	6 (20.0)	7 (23.3)	13 (43.3)	4 (13.3)	0 (0.0)
	z-value	NA	0.27	1.94	0.33	2.82**	1.94	0.0	1.07	1.40	NA
	Total	0 (0.0)	23 (38.3)	19 (31.7)	11 (18.3)	7 (11.7)	19 (31.7)	14 (23.3)	22 (36.7)	5 (8.3)	0 (0.0)
The packing of organic products looks less pleasing to the eye.	Ludhiana	4 (13.3)	6 (20.0)	7 (23.3)	13 (43.3)	0 (0.0)	5 (16.7)	2 (6.7)	15 (50.0)	1 (3.3)	7 (23.3)
	Patiala	2 (6.7)	9 (30.0)	9 (30.0)	6 (20.0)	4 (13.3)	2 (6.7)	2 (6.7)	20 (66.7)	6 (20.0)	0 (0.0)
	z-value	0.44	0.89	0.58	1.94	0.21	1.21	0.0	1.31	2.01*	2.82**
	Total	6 (3.3)	15 (25.0)	16 (26.7)	19 (31.7)	4 (13.3)	7 (11.7)	4 (6.7)	35 (58.3)	7 (11.7)	7 (11.7)
Organics are supreme products consumed only by rich people.	Ludhiana	0 (0.0)	9 (30.0)	10 (33.3)	5 (16.7)	6 (20.0)	0 (0.0)	4 (13.3)	15 (50.0)	1 (3.3)	10 (33.3)
	Patiala	1 (3.3)	9 (30.0)	12 (40.0)	5 (16.7)	3 (10.0)	2 (6.7)	1 (3.3)	15 (50.0)	12 (40.0)	0 (0.0)
	z-value	0.22	0	0.54	0.0	1.08	1.44	1.40	0.0	3.45**	3.46**
	Total	1 (1.7)	18 (30.0)	22 (36.7)	10 (16.7)	9 (15.0)	2 (3.3)	5 (8.3)	30 (50.0)	13 (21.7)	10 (16.7)
There is small variety of organic products.	Ludhiana	0 (0.0)	12 (40.0)	8 (26.7)	3 (10.0)	7 (23.3)	6 (20.0)	9 (30.0)	5 (16.7)	7 (23.3)	3 (10.0)
	Patiala	0 (0.0)	13 (43.3)	8 (26.7)	7 (23.3)	2 (6.7)	4 (13.3)	10 (33.3)	15 (50.0)	1 (3.3)	0 (0.0)
	z-value	NA	0.26	0	1.39	1.81	0.69	0.28	2.74**	2.28*	1.78
	Total	0 (0.0)	25 (41.7)	16 (26.7)	10 (16.7)	9 (15.0)	10 (16.7)	19 (31.7)	20 (33.3)	8 (13.3)	3 (5.0)
There are a lot of sales points for organic products.	Ludhiana	0 (0.0)	13 (43.3)	6 (20.0)	6 (20.0)	5 (16.7)	2 (6.7)	2 (6.7)	12 (40.0)	11 (36.7)	3 (10.0)
	Patiala	1 (3.3)	10 (33.3)	15 (50.0)	3 (10.0)	1 (3.3)	0 (0.0)	9 (30.0)	14 (46.7)	7 (23.3)	0 (0.0)
	z-value	1.01	0.80	2.44*	1.08	1.72	1.44	2.34*	0.52	1.13	1.78
	Total	1 (1.7)	23 (38.3)	21 (35.0)	9 (15.0)	6 (10.0)	2 (3.3)	11 (18.3)	26 (43.3)	18 (30.0)	3 (5.0)
It is difficult to identify real organic products.	Ludhiana	0 (0.0)	7 (23.3)	11 (36.7)	5 (16.7)	7 (23.3)	14 (46.7)	8 (26.7)	6 (20.0)	2 (6.7)	0 (0.0)
	Patiala	1 (3.3)	14 (46.7)	8 (26.7)	6 (20.0)	1 (3.3)	5 (16.7)	7 (23.3)	14 (46.7)	4 (13.3)	0 (0.0)
	z-value	1.01	1.89	0.83	0.33	2.28*	2.50*	0.30	2.19*	0.86	NA
	Total	1 (1.7)	21 (35.0)	19 (31.7)	11 (18.3)	8 (13.3)	19 (31.7)	15 (25.0)	20 (33.3)	6 (10.0)	0 (0.0)
I trust organic products.	Ludhiana	0 (0.0)	17 (56.7)	5 (16.7)	3 (10.0)	5 (16.7)	5 (16.7)	10 (33.3)	14 (46.7)	1 (3.3)	0 (0.0)
	Patiala	2 (6.7)	13 (43.3)	11 (37)	4 (13.3)	0 (0.0)	7 (23.3)	10 (33.3)	12 (40.0)	1 (3.3)	0 (0.0)
	z-value	1.44	1.55	2.25*	0.42	2.34*	0.65	0.0	0.52	0.0	NA
	Total	2 (3.3)	28 (46.7)	18 (30.0)	7 (11.7)	5 (8.3)	12 (20.0)	20 (33.3)	26 (43.3)	2 (3.3)	0 (0.0)

Figures in parenthesis indicate percentage,

\*Significant at 5% level,

\*\* Significant at 1% level

It was found that subjects of MIG of both cities had no practice to use organic foods daily. However, among HIG subjects 26.7 and 16.7% of the respondents from Ludhiana and Patiala city respectively were using organic foods daily (Table 5). It can be concluded that with the increase in income, consumption of organic foods increases on daily basis which might be due to higher affordability of these products with the increase in income. It was found that purchase intention towards organic foods and the level of income had a positive but insignificant relationship [8]. On weekly basis consumption of organic foods was found to be higher (50%) among HIG subjects as compared to that among MIG (30%) subjects. Overall, rare consumption of organic foods was found to be higher among MIG subjects as compared to HIG subjects. When the practices regarding the methods of washing, cutting, and cooking of organic foods

were compared with those of unconventional foods, they were found to be similar in all the subjects of both the cities. However, chopping of organic vegetables was found to be more convenient as compared to conventional vegetables by subjects of both the income groups from both the cities.

#### Knowledge, Attitude and Practice (KAP) Score

As shown in Table 6, there was no significant difference in the overall knowledge and attitude score of both MIG and HIG respondents. However, there was a statistically significant ( $p \leq 0.05$ ) difference in the overall practice score of respondents belonging to MIG and HIG. Hence, it can be concluded that although the level of income does not affect the knowledge and attitude regarding organic foods, but as the income increases the practices regarding using organic foods do increase.

**Table 5:** Practices towards organic foods among middle and high income group-subjects of selected cities (N=120)

Practices Ludhiana	MIG				HIG				
	Patiala	z-value	Total	Ludhiana	Patiala	z-value	Total		
Consumption of organic products	29 (96.7)	25 (83.3)	1.72	54 (90)	28 (93.3)	27 (90)	0.22	56 (93.3)	
Frequency of using organic foods	Rarely	9 (30)	9 (30)	0.58	18 (30)	2 (6.7)	3 (10)	0.94	5 (8.3)
	Monthly	14 (50)	8 (26.7)	1.61	22 (36.7)	5 (16.7)	7 (23.3)	0.65	12 (20)
	Weekly	5 (16.7)	13 (43.3)	0.00	18 (30)	15 (50)	15 (50)	0.24	30 (50)
	Daily	0 (0.0)	0 (0.0)	NA	2 (3.3)	8 (26.7)	5 (16.7)	0.94	13 (21.7)
Methods of washing vegetables	19 (63.3)	20 (66.7)	0.27	39 (65.0)	25 (83.3)	26 (86.7)	0.36	51 (85.0)	
	Before cutting								
After cutting	19 (63.3)	17 (56.7)	0.53	36 (60.0)	14 (46.7)	16 (53.3)	0.52	30 (50.0)	
Dipping	2 (6.7)	3 (10.0)	0.47	5 (8.3)	8 (26.7)	14 (46.7)	1.61	22 (36.7)	
Rinsing	2 (6.7)	3 (10.0)	0.47	5 (8.3)	10 (33.3)	10 (33.3)	0.00	20 (33.3)	
Rubbing while washing	1 (3.3)	4 (13.3)	1.40	5 (8.3)	2 (6.7)	1 (3.3)	0.59	3 (5.0)	
Chopping is easier	22 (73.3)	23 (76.7)	0.30	45 (75.0)	21 (70.0)	17 (56.7)	2.36*	45 (75.0)	
Cooking practice vary for organic food products	6 (20.0)	0 (0.0)	2.58*	6 (10.0)	0 (0.0)	4 (13.3)	2.07*	4 (6.7)	
Cooking takes same time	15 (50.0)	8 (26.7)	1.86	23 (38.30)	12 (40.0)	18 (60.0)	1.55	30 (50.0)	
Cooking Methods									
	Open Pan	14 (46.7)	24 (80.0)	2.68*	38 (63.3)	23 (76.7)	12 (40.0)	2.88**	35 (58.3)
	Pan with lid	12 (40.0)	2 (6.7)	3.05**	14 (23.3)	17 (56.7)	14 (46.7)	0.78	31 (51.7)
	Pressure Cooking	21 (70.0)	12 (40.0)	2.34*	33 (55.0)	15 (50.0)	19 (63.3)	1.04	34 (56.7)

Figures in Parenthesis indicate percentage

\* Significant at 5% level,

\*\* Significant at 1% level

**Table 6:** Mean Knowledge, Attitude and Practice score in middle and high income group subjects of selected cities (N=120)

Parameters	MIG(n=60)	HIG(n=60)	z-value	Total (N=120)
Knowledge	4.6 ± 0.7	4.8 ± 0.6	0.55	4.7 ± 0.6
Attitude	5.0 ± 0.5	5.2 ± 0.4	0.48	5.1 ± 0.4
Practice	3.0 ± 0.5	3.8 ± 0.4	2.40*	3.4 ± 0.4
KAP score	12.6 ± 1.7	13.8 ± 1.3	1.23	13.2 ± 1.4

Values are Mean ± SD,

\* Significant at 5% level, Ns-non significant

**Table 7:** Correlation coefficients (r) of KAP score with demographic factors

Correlation of with KAP score	Age	Education	Income
Knowledge	0.19*	0.23**	0.11NS
Attitude	0.07NS	0.08NS	0.12NS
Practice	0.09NS	0.09NS	0.18*
KAP score	0.12NS	0.11NS	0.10NS

\*- Correlation is significant at the 0.05 level (2-tailed).

\*- Correlation is significant at the 0.01 level (1-tailed).

NS- Non-significant

### *Correlations between various factors and purchase of organic foods*

In different states of India food consumption varies and the factors which influence consumption are regional, cultural, ethnic, income and difference in agricultural production. In the present study, correlation coefficient values of KAP scores with demographic factors like age, education and income were calculated and are represented in the Table 7.

The knowledge of the respondents had a significantly ( $p \leq 0.05$ ) positive correlation with age and education, indicating that the knowledge increases with age and education, whereas it had a positive but non-significant correlation with income. The attitude of the respondents had a positive but non-significant correlation with age, education and income. Similarly, the practice score had a positive but non-significant correlation with age and education. However, it had a significantly ( $p \leq 0.05$ ) positive correlation with the income.

### **Conclusion**

Consumption of organic foods is increasing because of environmental and health issues. From the present study, it can be concluded that the knowledge and attitude of the consumers regarding organic foods is influenced by their education level and age as well. Furthermore, the level of income does not affect their knowledge and attitude regarding organic foods, but it does affect practice of using the same. The data can help both marketers and policy makers in formulating strategies and policies regarding the supply and availability of organic foods.

### **References**

1. Aertsens J. The influence of subjective and objective knowledge on attitude, motivations and consumption of organic food. *British Food Journal* 2011;113:1353-78.
2. Attanasio S. Carelli A., Cappelli L., and Papetti. Organic food. *International Journal of Latest Research in Science and Technology*. 2013;2:128-32.
3. Baltas G. Nutrition labeling: issues and policies. *European Journal of Marketing* 2001;35: 708-21.
4. Chinnici G.D., Amico M. and Pecorino B. A multivariate statistical analysis on the consumer of organic products. *British Food Journal*. 2002;104: 730-65.
5. Demeritt L. All things organic. A look at the organic consumer. The Hartman group, Bellevue, WA. 2002.
6. Heckman J. A history of organic farming: Transitions from Sir Albert Howard's War in the soil to USDA National Organic Program. *Renewable Agriculture and Food Systems*. 2005;21:143-50.
7. Nooripoor, M. Shahvali, M. Zarafshani, K. Integration of communication media for horticultural sustainability: The application of multiple criteria decision making (MCDM). *American-Eurasian Journal of Agricultural and Environmental Science*. 2008;3:137-47.
8. Omar N.A., Nazri M.A., Osman L.H. and Ahmad M.S. The effect of demographic factors on consumer intention to purchase organic products in Klang Valley. *Malaysian Journal of Society and Space*. 2016;12:68-82.
9. Suh B.W. A study on the relationships between TV programs related to health and diet and consumers' diet life: Focus on the age group in 20s and 30s. *Journal of Korean Food Marketing Association*. 2008;25:29-48.

**STATEMENT ABOUT OWNERSHIP AND OTHER PARTICULARS**  
**“International Journal of Food, Nutrition and Dietetics” (See Rule 8)**

- |   |   |  |
|---|---|--|
| 1. Place of Publication   | : | Delhi  |
| 2. Periodicity of Publication   | : | Quarterly  |
| 3. Printer's Name   | : | <b>Dinesh Kumar Kashyap</b>  |
| Nationality   | : | Indian   |
| Address   | : | 395-A, Pocket-II, Mayur Vihar,<br>Phase-1, Delhi-91  |
| 4. Publisher's Name   | : | <b>Dinesh Kumar Kashyap</b>  |
| Nationality   | : | Indian   |
| Address   | : | 395-A, Pocket-II, Mayur Vihar,<br>Phase-1, Delhi-91  |
| 5. Editor's Name  | : | <b>Dinesh Kumar Kashyap</b>  |
| Nationality   | : | Indian   |
| Address   | : | 395-A, Pocket-II, Mayur Vihar,<br>Phase-1, Delhi-91  |
| 6. Name & Address of Individuals<br>who own the newspaper and particulars of<br>shareholders holding more than one per cent<br>of the total capital | : | <b>Red Flower Publication Pvt. Ltd.</b><br>41/48, DSIDC, Pocket-II<br>Mayur Vihar, Phase-1, Delhi-91 |

I **Dinesh Kumar Kashyap**, hereby declare that the particulars given above are true to the best of my knowledge and belief.

Sd/-

**(Dinesh Kumar Kashyap)**

## Impact of Nutrition Counseling on Dietary Intake of Obese Women from Low, Middle and High Income Group of Hoshiarpur City of Punjab

Shagun Aggarwal<sup>1</sup>, Neerja Singla<sup>2</sup>, Poonam Bakheta<sup>3</sup>

### How to cite this article:

ShagunAggarwal, Neerja Singla, Poonam Bakheta. Impact of Nutrition Counseling on Dietary Intake of Obese Women from Low, Middle and High Income Group of Hoshiarpur City of Punjab. Int J Food Nutr Diet. 2019;7(1): 33-40.

### Abstract

A study was conducted to assess the impact of nutrition counseling on food consumption pattern of obese women from low, middle and high income group. To study the prevalence of obesity, 150 women in the age group of 40-50 years, belonging to low (LIG), middle (MIG) and high (HIG) income group (50 from each income group) were randomly selected from sub-mountainous region of Hoshiarpur city. Twenty five normal (BMI 20-22.9 kg/m<sup>2</sup>) and 25 obese (BMI ≥ 27.5 kg/m<sup>2</sup>) women from each income group were selected. Nutrition counseling was imparted to the subjects for 3 months. Its impact was assessed in terms of their dietary intake and gain in knowledge. Prevalence of overweight and obesity was higher in women belonging to HIG (70.7%) followed by MIG (67.3%) and LIG (56.6%). Percent adequacy of food groups like fats & oils, cereals, milk & milk products, roots and tubers and fruits was significantly ( $p < 0.01$ ) higher in obese subjects as compared to normal subjects. Nutrition counseling of the obese subjects resulted in a gain in their knowledge and quantum of improvement, which further led to a decrease in the intake of fats & oils, cereals, milk & milk products, and roots & tubers by the obese subjects, compared to their consumption before the intervention and compared to women who only received routine counseling.

**Keywords:** Obesity; BMI; Income groups; Nutrition counseling.

### Introduction

Obesity is a medical condition in which excess body fat has accumulated to the extent that it may have a negative effect on health, leading to reduced life expectancy and/or increased health problems. It can be seen as the first wave of a defined cluster

of non-communicable diseases called "New World Syndrome," creating an enormous socio-economic and public health burden in poorer countries. It has become serious epidemic health problem, estimated to be the fifth leading cause of mortality at global level [1]. The World Health Organization has described obesity as one of the most neglected public health problem, affecting every region of the globe.

Increasing evidence of overweight/obesity indicates that India is experiencing a double burden of communicable and non-communicable diseases. In 2005, nearly 14% of women aged 18 to 49 were overweight or obese, with higher rates among urban women (25%) than rural women (8%) [2]. The rate of overweight and obesity in women, overall increased by 3.5% a year from 1998 to 2005. Comparison of two major studies conducted by National Family Health Survey (NFHS- 2 in 1998-1999 and NFHS-

**Author Affiliation:** <sup>1</sup>Student, <sup>2</sup>Assistant Scientist, <sup>3</sup>Assistant Nutritionist, Department of Food and Nutrition, College of Home Science, Punjab Agricultural University, Ludhiana, Punjab 141004, India.

**Corresponding Author: Poonam Bakheta**, Assistant Nutritionist, Department of Food and Nutrition, College of Home Science, Punjab Agricultural University, Ludhiana, Punjab 141004, India.

**E-mail:** [poonambakheta@pau.edu](mailto:poonambakheta@pau.edu)

3 in 2005-2006 showed that prevalence of obesity among Indian women has elevated from 10.6 to 12.6 per cent (increased by 24.52%). The prevalence is found more in women of age between 40-49 yrs (23.7%), residing in cities (23.5%) and households in the highest wealth quantile (30.5%). Highest percentage of obese women (29.9%) was found in Punjab followed by Kerala (28%) and Delhi (26%), all of which are relatively richer states [3]. Therefore, the increasing trend of obesity among Indians is becoming a matter of concern.

Worldwide debates are continuing about using ethnic-specific or standard methods of assessment of obesity. The most common measure of overweight and obesity is the Body Mass Index (BMI), which is determined by the individual's weight and height. When a BMI value is over 25 kg/m<sup>2</sup>, the value indicates an overweight. People are considered obese when their Body Mass Index (BMI) exceeds 30 kg/m<sup>2</sup> [4].

One of the major causes of obesity is the changes in the diet in terms of quantity and quality which has become more "westernized" and sedentary life styles [5]. To effectively cure obesity with diet management, controlling the excessive nutrient intake but also concretely instructing the know how of choosing right foods and desirable eating pattern are necessary [6]. Nutrition education/counseling is therefore needed to motivate and create public awareness to preserve healthy life styles and adopt healthy dietary habits to reduce risk factors of obesity.

Nutrition counseling is one of the most effective tools of changing the food habits of the people without affecting their sentiments. It is a process by which the beliefs, attitude, environmental influents and knowledge about food and health are canalized into actual practices [7]. To maintain the reduced body weight after weight control, it is necessary to correct the way or pattern of eating, exercise and overall lifestyle [8]. Also it is important to acquire the ability to control one's own dietary habit by continuous nutrition education and counseling. Keeping these facts in view, the present community based study was carried out in Hoshiarpur city, Punjab to assess the impact of nutrition counseling on food and nutrient intake of obese women of low, middle and high income groups.

## Materials and Methods

### *Selection of subjects*

A total of 150 women in the age group

40-50 years (50 belonging to each income group *i.e.* low income group, middle income group and high income group) were selected randomly among the localities of Hoshiarpur city to study the prevalence of obesity. The Body Mass Index (BMI) was used as a criterion for the assessment of prevalence of obesity. According to the WHO classification for Asian population the women having BMI more than 27.5 are considered as obese [4]. Based on the BMI, further 50 subjects from each income group were selected into two categories: normal and obese (25 subjects in each category), thus making a total sample size of 150 subjects.

An interview schedule was drafted to obtain the desired information on various aspects such as general information, family history, dietary intake, etc. of the subjects. Pretesting was carried out on 15 subjects to test the reliability and validity of questionnaire and the necessary modifications were made. Pretested subjects were not included in the study.

### *Background Information*

Information about the subjects pertaining to age, educational qualification, monthly income, marital status, family type, family size, nature of work, lifestyle related information, medical history and usage of drugs was recorded through a pre-structured tested questionnaire was collected.

### *Dietary assessment*

Information regarding the intake of food for three consecutive days was collected from the subjects using 24-hour recall method. Mean daily intake of different food groups including cereals, pulses, fruits, green leafy vegetables, roots and tubers, other vegetables, milk and milk products, sugar and jaggery and fats and oils was calculated by taking mean intake of three consecutive days. Cooked food consumed was converted into their raw equivalents. Different food items were converted into their raw equivalents and the average daily intake of energy, proximate principles, important vitamins and minerals were calculated by Dietcal a computer software [9].

### *Assessment of knowledge*

A multiple choice questionnaire was also designed to test the nutrition knowledge of the subjects. After pre-testing the knowledge of the respondent's nutrition education in the form of lectures, trainings, talks, group discussions etc. was imparted to 150 subjects for a period of three months. Full package of knowledge through pamphlet and

booklet on Obesity were also given to the subjects. For evaluating questionnaire 1 score was awarded for each correct and zero for wrong answer. Post nutrition knowledge test was conducted after three months to note the gain in score and quantum of improvements in the diet. Gain in knowledge was calculated using the following equation.

$$\text{Gain in knowledge} = \text{Score of post test} - \text{Score of pre test}$$

$$\text{Quantum of Improvement} = \frac{\text{Post test scores}}{\text{Pre test scores}}$$

*Statistical analysis*

The data on all the parameters viz. food intake, nutrient intake and gain in knowledge were analyzed statistically with the help of mean, standard error, percentage and paired-*t* test.

**Results and Discussion**

*Dietary Intake*

It was observed (Table 1) that majority of the obese subjects *i.e.* 24, 68 and 52 percent were vegetarians whereas 48, 20 and 16 percent were non-vegetarians in LIG, MIG and HIG, respectively. However, corresponding values for ovatarians in all the three income groups were 28, 12 and 32 percent, respectively.

The daily consumption pattern of fruits and salads in the normal category showed that among the LIG (60%) and MIG (40%) consumed twice a week whereas HIG (40%) consumed daily. In the obese category LIG (72%), MIG (52%) and HIG (68%) consumed thrice a week. As evident from the Table 1 majority of the obese subjects *i.e.* 92, 100 and 88 percent in LIG, MIG and HIG, respectively preferred to have food from outside. The data clearly indicated that the frequency of

**Table 1:** Dietary Habits of the selected subjects.

Parameters	LIG (n=50)		MIG (n=50)		HIG (n=50)	
	Normal (n=25)	Obese (n=25)	Normal (n=25)	Obese (n=25)	Normal (n=25)	Obese (n=25)
<b>Dietary Pattern</b>						
Vegetarian	15 (60)	6 (24)	11 (44)	17 (68)	8 (32)	13 (52)
Non-vegetarian	7 (28)	12 (48)	9 (36)	5 (20)	8 (32)	4 (16)
Ovatarian	3 (12)	7 (28)	5 (20)	3 (12)	9 (36)	8 (32)
<b>No. of meals per day</b>						
Twice	1 (4)	-	-	-	-	-
Thrice	24 (96)	22 (88)	16 (64)	1 (4)	7 (28)	1 (4)
Four times	-	3 (12)	9 (36)	9 (36)	17 (68)	9 (36)
More than four times	-	-	-	15 (60)	1 (4)	15 (60)
<b>Fruits and salad consumption</b>						
Daily	7 (28)	4 (16)	7 (28)	12 (48)	10 (40)	3 (12)
Twice a week	15 (60)	3 (12)	10 (40)	-	9 (36)	5 (20)
Thrice a week	3 (12)	18 (72)	8 (32)	13 (52)	6 (24)	17 (68)
<b>Outside food consumption</b>						
Yes	9 (36)	23 (92)	17 (68)	25 (100)	20 (80)	22 (88)
No	16 (64)	2 (8)	8 (32)	-	5 (20)	3 (12)
<b>Outside food consumption pattern</b>						
Once a week	4 (16)	8 (32)	4 (16)	7 (28)	5 (20)	3 (12)
Twice a week	-	4 (16)	2 (8)	13 (52)	3 (12)	14 (56)
Thrice a week	1 (4)	-	2 (8)	4 (16)	-	3 (12)
Fortnightly	4 (16)	9 (36)	9 (36)	1 (4)	12 (48)	2 (8)
Daily	-	3 (12)	-	-	-	-
Rarely	16 (64)	1 (4)	8 (32)	-	5 (20)	3 (12)

LIG-Low income group, MIG- Middle income group and HIG- High income group

Figures in parenthesis indicate percentage.

food consumption from outside was more in obese subjects belonging to all the three income groups, which might be a major contributor to obesity.

#### *Consumption pattern of various food items*

The data regarding food frequency (Table 2) indicated that consumption of fried foods and snacks was significantly higher in obese subjects of all the three income groups, whereas a non-significant difference was observed in the frequency score of carbonated beverages, sweet dishes and bakery products.

#### *Food intake*

Mean daily food intake by normal and obese subjects of all the income groups and their comparison with suggested dietary intake [10] are given in Table 3. Mean daily intakes of energy yielding foods such as cereals, roots and tubers, fats and oils and milk and milk products were significantly higher by obese subjects of all the three income groups as compared to normal subjects. The percent adequacy of all the food groups by LIG group was below the suggested intakes. Among MIG, the intake of fruits (115.8 and 115.0%), milk and milk products (130.1 and 137.8%) and fats and oils (100.6 and 212.9%) was found to be adequate by normal and obese subjects respectively. The percent adequacy of pulses was 144.2% and 166.0% by normal and obese subjects of HIG. The consumption of pulses increased with increase in income level. It was also reported that obese women consumed significantly higher ( $p \leq 0.05$ ) quantities of pulses than normal women [11]. Consumption of roots and tubers was higher in the obese subjects in all the three income groups *i.e.*  $199.8 \pm 58.0$ ,  $185.4 \pm 66.6$  and  $156.2 \pm 30.8$ g as compared to the normal subjects *i.e.*  $145.9 \pm 53.8$ ,  $146.2 \pm 41.1$  and  $120.4 \pm 40.1$ g in

MIG, HIG and LIG, respectively.

It was observed that fruit intake was significantly higher in normal as well as obese subjects belonging to MIG and HIG as compared to that in the LIG. The consumption of milk and milk products was significantly different among the normal and the obese subjects in all the income groups. The LIG usually consume milk in the form of tea and very few were habitual of drinking milk. However, MIG and HIG subjects consumed milk in the form of paneer, cheese, flavoured milk, tea, sweets etc. Kaur (1995) [12] also reported less milk consumption among LIG (325g) as compared to MIG (658g) and HIG (863g). High intake of milk and its products in obese were also reported in literature [13]. The results (Table 3) showed that average daily intake of fats and oils was significantly ( $p \leq 0.05$ ) higher among the obese subjects as compared to the normal subjects with high intake of  $64.4 \pm 13.1$  g by HIG obese subjects in comparison to  $49.4 \pm 16.2$  g by the normal subjects. The percent adequacy for fats and oils in normal subjects belonging to LIG, MIG and HIG was 28.8, 100.6 and 247 percent whereas in the obese subjects it was 41.2, 212.9 and 322 percent, respectively. The higher intake of fats and oils was mainly because of the more consumption of fried food among the three groups. It was also reported in the literature that more than 100 percent adequacy of fats and oil among LIG, MIG and HIG and consumption of higher amount of fat among the HIG (67g) as compared to MIG (58g) and LIG (56g) [14].

#### *Nutrient intake*

The mean daily intake of nutrients by the selected subjects is shown in Table 4. The percent adequacy among normal subjects was found out to be 42.5%, 73.6% and 103.3% whereas in obese subjects it

**Table 2:** Frequency food scores of various food items.

Parameters	LIG (n=50)			MIG (n=50)			HIG (n=50)		
	Normal (n=25)	Obese (n=25)	t value	Normal (n=25)	Obese (n=25)	t value	Normal (n=25)	Obese (n=25)	t value
Fried foods	2.3 ± 1.4	3.8 ± 0.7	4.63*	3.4 ± 1.4	3.2 ± 0.8	2.63**	3.5 ± 0.8	3.9 ± 1.2	2.80**
Snacks	2.8 ± 1.5	3.6 ± 0.8	2.24**	3.6 ± 1.1	4.6 ± 0.5	3.9*	3.5 ± 0.7	4.7 ± 0.5	6.65*
Carbonated beverages	1.2 ± 0.8	1.4 ± 1.0	0.428 <sup>NS</sup>	1.9 ± 1.1	2.2 ± 1.0	0.76 <sup>NS</sup>	2.3 ± 1.3	2.7 ± 1.4	0.92 <sup>NS</sup>
Sweet dishes	2.6 ± 1.3	3.1 ± 1.6	1.12 <sup>NS</sup>	3.2 ± 1.6	3.6 ± 1.8	0.82 <sup>NS</sup>	3.7 ± 1.0	3.9 ± 1.5	0.426 <sup>NS</sup>
Bakery products	0.2 ± 0.4	0.7 ± 0.7	2.65**	1.3 ± 1.1	1.5 ± 1.1	0.73 <sup>NS</sup>	2.2 ± 0.9	2.4 ± 1.1	0.66 <sup>NS</sup>

LIG-Low income group, MIG- Middle income group, HIG- High income group

Values are Mean ± SD Mean scores (0, 1,2,3,4,5,6)

\* Significant at 1%

\*\* Significant at 5% <sup>NS</sup>- Non significant

was 58.5%, 106.3% and 120.7% belonging to LIG, MIG and HIG respectively when compared with RDA [10]. A higher intake of energy in MIG and HIG subjects might be due to higher consumption of fats and oils and also sugar among them. A similar view has also been reported that a higher intake of energy by the obese respondents than their respective RDA [15,16]. Low intake of energy

by LIG as compared to MIG and HIG was also reported in the literature [12].

Except fats, Dietary fiber, calcium and vitamin A the consumption of all other nutrients was significantly different among the normal and the obese subjects in all the income groups. In LIG, the percent adequacy for fat was 100.5 and 114.6 percent

**Table 3:** Daily food intake of the selected subjects

Food groups (g/day)	# Suggested Dietary intake	LIG (n=50)			MIG (n=50)			HIG (n=50)		
		Normal (25)	Obese (25)	t value	Normal (25)	Obese (25)	t value	Normal (25)	Obese (25)	t value
Cereals	270	122.0 ± 30.6	207.2 ± 48.4	7.43*	172.0 ± 32.8	230.8 ± 53.8	4.65*	185.4 ± 39.6	241.1 ± 66.2	3.60*
Pulses and Legumes	55	19.8 ± 8.4	19.6 ± 8.6	0.08 <sup>NS</sup>	46.6 ± 17.5	48.6 ± 33.9	0.48 <sup>NS</sup>	79.3 ± 24.0	82.2 ± 31.3	0.36 <sup>NS</sup>
Green leafy Vegetable	100	2.4 ± 6.2	3.3 ± 6.7	0.50*	4.0 ± 7.0	6.9 ± 19.9	0.69 <sup>NS</sup>	8.6 ± 13.6	8.4 ± 20.7	0.053 <sup>NS</sup>
Roots and Tubers	200	120.4 ± 40.1	156.2 ± 30.8	3.53*	145.9 ± 53.8	199.8 ± 58.0	3.40*	146.2 ± 41.1	185.4 ± 66.6	2.50*
Other vegetable	200	24.5 ± 14.0	28.0 ± 11.6	0.96 <sup>NS</sup>	38.4 ± 26.0	39.7 ± 47.8	1.34 <sup>NS</sup>	41.8 ± 18.5	43.3 ± 38.4	1.75 <sup>NS</sup>
Fruits	100	18.5 ± 21.9	17.2 ± 18.2	0.23 <sup>NS</sup>	115.6 ± 98.4	115.0 ± 94.6	1.53 <sup>NS</sup>	192.2 ± 87.2	193.6 ± 109.8	0.049 <sup>NS</sup>
Milk and milk Products	300	149.6 ± 50.0	175.4 ± 65.0	3.56*	390.2 ± 136.2	413.4 ± 128.4	3.62*	490.9 ± 133.8	542.5 ± 154.7	4.26*
Sugar	20	9.5 ± 6.1	10.0 ± 7.0	0.22 <sup>NS</sup>	14.3 ± 7.9	12.5 ± 12.1	0.63 <sup>NS</sup>	24.1 ± 11.2	18.2 ± 20.2	1.28 <sup>NS</sup>
Fats and Oils	20	5.7 ± 2.5	8.2 ± 3.8	2.68**	20.1 ± 10.0	42.5 ± 15.2	6.12*	49.4 ± 16.2	64.4 ± 13.1	2.29**
Meat and Poultry		19.0 ± 36.2	23.4 ± 28.9	0.47 <sup>NS</sup>	15.0 ± 31.5	21.7 ± 46.5	0.59 <sup>NS</sup>	34.0 ± 72.7	37.6 ± 61.6	0.188 <sup>NS</sup>

LIG-Low income group, MIG-Middle income group, HIG- High income group

\* Significant at 1%

\*\* Significant at 5% <sup>NS</sup>-Non significant

# ICMR (2010)

**Table 4:** Daily nutrient intakes of selected subjects

Nutrients	#RDA	LIG (n=50)			MIG (n=50)			HIG (n=50)		
		Normal (n=25)	Obese (n=25)	t-value	Normal (n=25)	Obese (n=25)	t-value	Normal (n=25)	Obese (n=25)	t-value
Energy (Kcal)	1900	807 ± 94.4	1111 ± 210.0	6.61*	1399 ± 235.6	2020 ± 267.2	8.71*	1963 ± 405.3	2295 ± 323.8	3.19*
Protein (g)	55	28.9 ± 4.8	39.7 ± 7.7	5.93*	45.5 ± 9.0	64.4 ± 10.9	6.67*	58.9 ± 13.4	71.9 ± 11.4	3.69*
Fat (g)	20	20.1 ± 7.6	22.9 ± 5.2	1.52 <sup>NS</sup>	46.1 ± 4.6	56.1 ± 19.6	1.05 <sup>NS</sup>	85.5 ± 21.3	95.0 ± 20.8	1.60 <sup>NS</sup>
Carbohydrates (g)	-	120.8 ± 23.7	185.4 ± 41.3	6.76*	184.3 ± 33.4	250.2 ± 36.8	6.62*	217.9 ± 46.2	264.5 ± 53.6	3.29*
Total dietary fiber (g)	-	7.4 ± 2.5	7.4 ± 2.7	0.06 <sup>NS</sup>	10.3 ± 3.7	13.4 ± 4.6	1.94 <sup>NS</sup>	15.5 ± 3.3	16.6 ± 6.3	0.72 <sup>NS</sup>
Calcium (mg)	600	437.4 ± 112.7	441.4 ± 94.3	0.136 <sup>NS</sup>	892.6 ± 263.6	982.1 ± 292.7	1.67 <sup>NS</sup>	1198.9 ± 344.3	1344.2 ± 310.4	1.56 <sup>NS</sup>
Iron (mg)	21	7.9 ± 1.4	12.0 ± 2.8	6.46*	11.7 ± 2.4	17.8 ± 4.2	6.14*	14.4 ± 2.4	18.8 ± 4.9	4.03*
Vitamin A (µg)	600	182.0 ± 164.6	171.6 ± 126.4	0.24 <sup>NS</sup>	259.2 ± 172.7	351.7 ± 229.5	1.60 <sup>NS</sup>	470.6 ± 330.1	587.9 ± 580.7	0.87 <sup>NS</sup>
Thiamin (mg)	1.0	0.8 ± 0.1	1.2 ± 0.2	7.32*	1.1 ± 0.2	1.6 ± 0.3	6.36*	1.4 ± 0.2	1.7 ± 0.4	3.72*
Riboflavin (mg)	1.1	0.4 ± 0.1	0.7 ± 0.2	5.96*	0.7 ± 0.1	1.0 ± 0.3	4.17*	1.0 ± 0.3	1.2 ± 0.3	2.19**
Niacin (mg)	12	6.1 ± 1.3	10.2 ± 2.5	6.99*	8.0 ± 2.2	11.6 ± 2.8	4.94*	8.8 ± 2.0	12.0 ± 3.5	3.91*
Folic acid (µg)	200	102.2 ± 26.6	140.4 ± 32.3	4.56*	142.0 ± 40.3	204.0 ± 63.7	4.11*	198.7 ± 56.4	235.2 ± 66.2	3.04*
Vitamin C (mg)	40	30.2 ± 9.8	36.4 ± 9.7	2.22**	69.8 ± 35.6	98.0 ± 38.0	2.70**	103.6 ± 34.5	119.6 ± 90.3	2.42**

LIG-Low income group, MIG-Middle income group, HIG- High income group,

# ICMR (2010)

\* Significant at 1%

\*\* Significant at 5% <sup>NS</sup>-Non significant

for normal and obese subjects, all other nutrients were inadequately consumed except thiamin in obese subjects. The obese subjects of MIG and normal subjects of HIG were consuming all the nutrients adequately except iron, riboflavin, niacin and folic acid whereas the obese subjects of HIG were having all the nutrients in adequate amounts as compared to RDA. The percent adequacy was found out to be 100.5%, 230.9% and 427.5% by the normal subjects of LIG, MIG and HIG whereas for the obese subjects it was reported as 114.6, 280.85 and 475.45 percent, respectively. This excessive consumption of fat among Punjabi women has also been reported in the literature [17,18]. The high intake of fat in all the three groups might be due to strong liking and a higher consumption of fried and fast foods. In a previous study, it was that dietary fat promotes more obesity than carbohydrate or protein of same energy value [19]. Similar results have been reported in the literature [20,21,16].

The consumption of vitamins namely thiamine, riboflavin, niacin, folic acid and vitamin C differ significantly among the normal and the obese subjects while vitamin A did not differ significantly among the normal and the obese subjects of the three income groups. The results clearly indicated that

the obese subjects consumed large meals resulting in high intakes of energy, protein, carbohydrates and fat but on the other hand, the large meals also provided them with more vitamins which have positive effect on their overall health.

### Nutrition Counseling

Nutrition counseling regarding obesity, its causes, consequences, control and prevention, balanced diet, dietary recommendations, cooking practices and ill effects of junk foods was imparted to the subjects for a period of three months. The evaluation of scores was done before (pre-test) and after (post-test) nutrition counseling to assess the difference in nutrition knowledge among normal and obese subjects. The average gain in knowledge during nutrition counseling is presented in Table 5. Quantum of improvement in normal and obese subjects of HIG was ( $1.2 \pm 2.1$  and  $1.1 \pm 1.3$  respectively), whereas in MIG it was ( $1.2 \pm 1.2$  and  $1.4 \pm 2.0$  respectively) followed by LIG normal and obese subjects ( $1.3 \pm 2.1$  and  $1.5 \pm 2.0$  respectively) depicting a significant difference among the quantum of improvement of the subjects in all the three income groups.

**Table 5:** Gain in scores before and after nutrition counseling

Groups	LIG (n=50)			MIG (n=50)			HIG (n=50)		
	Normal (n=25)	Obese (n=25)	t value	Normal (n=25)	Obese (n=25)	t value	Normal (n=25)	Obese (n=25)	t value
Pre Test	19.1 ± 2.4	17.6 ± 2.7	1.99 <sup>NS</sup>	21.3 ± 2.3	22.1 ± 2.6	1.14 <sup>NS</sup>	21.9 ± 2.4	23.4 ± 2.1	2.28 <sup>**</sup>
Post Test	25.2 ± 1.9	25.6 ± 1.1	0.98 <sup>NS</sup>	26.1 ± 2.1	26.8 ± 1.3	1.34 <sup>NS</sup>	25.7 ± 2.0	26.1 ± 1.4	0.78 <sup>NS</sup>
t- Value	9.80*	13.22*		7.67*	8.01*		5.96*	5.21*	
Gain	6.0 ± 2.0	8.0 ± 2.0	3.21*	4.8 ± 2.2	4.9 ± 2.0	2.19 <sup>**</sup>	3.8 ± 2.1	2.7 ± 1.3	2.12 <sup>**</sup>
Quantum of improvement	1.3 ± 0.1	1.4 ± 0.2	2.95*	1.2 ± 0.1	1.4 ± 0.1	2.19 <sup>**</sup>	1.1 ± 0.1	1.1 ± 0.0	2.15 <sup>**</sup>

LIG-Low income group, MIG-Middle income group, HIG- High income group <sup>NS</sup>-Non significant \* Significant at 1% \*\* Significant at 5%

**Table 6:** Food intake of the subjects before and after nutrition counseling.

Food groups (g/d)	LIG (n=25)			MIG (n=25)			HIG (n=25)		
	Before	After	t value	Before	After	t value	Before	After	t value
Cereals	207.2 ± 48.4	200.8 ± 44.4	2.24 <sup>**</sup>	230.8 ± 53.8	227.2 ± 51.8	2.45 <sup>**</sup>	241.1 ± 66.2	236.4 ± 65.2	2.41 <sup>**</sup>
Pulses and Legumes	19.6 ± 8.6	17.2 ± 6.6	1.56 <sup>NS</sup>	48.6 ± 33.9	42.9 ± 30.9	1.89 <sup>NS</sup>	82.2 ± 31.3	78.2 ± 30.3	1.78 <sup>NS</sup>
Green leafy vegetable	3.3 ± 6.7	3.8 ± 6.7	1.67 <sup>NS</sup>	6.0 ± 19.9	6.9 ± 18.5	1.76 <sup>NS</sup>	8.4 ± 20.7	7.9 ± 19.7	1.43 <sup>NS</sup>
Roots and Tubers	156.2 ± 30.8	152.5 ± 28.8	1.2 <sup>NS</sup>	199.8 ± 58.0	194.8 ± 54.0	1.87 <sup>NS</sup>	185.4 ± 66.6	180.4 ± 66.6	1.14 <sup>NS</sup>
Other vegetable	28.0 ± 11.6	26.0 ± 10.3	1.48 <sup>NS</sup>	39.7 ± 47.8	36.7 ± 47.6	1.64 <sup>NS</sup>	43.3 ± 38.4	40.5 ± 36.2	1.52 <sup>NS</sup>
Fruits	17.2 ± 18.2	19.6 ± 15.4	1.89 <sup>NS</sup>	115.0 ± 94.6	118.0 ± 93.6	1.56 <sup>NS</sup>	193.6 ± 109.8	195.9 ± 108.8	1.84 <sup>NS</sup>
Milk and Milk Products	175.4 ± 65.0	170.3 ± 62.0	2.33 <sup>**</sup>	413.4 ± 128.4	409.6 ± 125.4	2.76 <sup>**</sup>	542.5 ± 154.7	539.5 ± 153.7	2.84 <sup>**</sup>
Sugar	10.0 ± 7.0	9.0 ± 6.5	1.91 <sup>NS</sup>	12.5 ± 12.1	11.5 ± 10.1	1.76 <sup>NS</sup>	18.2 ± 20.0	16.2 ± 18.0	1.54 <sup>NS</sup>
Fats and Oils	8.2 ± 3.8	7.0 ± 2.5	2.87 <sup>**</sup>	42.5 ± 15.2	40.5 ± 12.2	2.75 <sup>**</sup>	64.4 ± 13.1	62.3 ± 13.1	2.69 <sup>**</sup>
Meat and Poultry	23.4 ± 28.9	21.2 ± 26.9	2.22 <sup>**</sup>	21.7 ± 46.5	19.6 ± 44.5	2.40 <sup>**</sup>	37.6 ± 61.6	36.8 ± 61.6	2.37 <sup>**</sup>

LIG-Low income group, MIG-Middle income group, HIG- High income group <sup>NS</sup>-Non significant \* Significant at 1% \*\* Significant at 5%

**Table 7:** Nutrient intake of the subjects before and after nutrition counseling.

Nutrients	LIG (n=25)			MIG (n=25)			HIG (n=25)		
	Before	After	t value	Before	After	t value	Before	After	t value
Energy (Kcal)	1111 ± 210.0	1100 ± 208.0	2.87**	2020 ± 267.2	2017 ± 267.2	2.76**	2295 ± 323.8	2289 ± 323.8	2.54**
Protein (g)	39.7 ± 7.7	38.7 ± 7.7	1.43 <sup>NS</sup>	64.4 ± 10.9	63.7 ± 10.9	1.11 <sup>NS</sup>	71.9 ± 11.4	70.9 ± 11.4	0.68 <sup>NS</sup>
Fat (g)	22.9 ± 5.2	20.3 ± 5.2	2.88**	56.1 ± 19.6	54.6 ± 19.6	2.75**	95.0 ± 20.8	89.0 ± 20.8	2.45**
Carbohydrates (g)	185.4 ± 41.3	180.2 ± 41.3	2.44**	250.2 ± 36.8	248.5 ± 36.8	2.31**	264.5 ± 53.6	260.5 ± 53.6	2.45**
Total dietary fibre (g)	7.4 ± 2.7	7.4 ± 2.7	1.10 <sup>NS</sup>	13.4 ± 4.6	13.4 ± 4.6	1.45 <sup>NS</sup>	16.6 ± 6.3	16.6 ± 6.3	1.87 <sup>NS</sup>
Calcium (mg)	441.4 ± 94.3	436.4 ± 94.3	2.30**	982.1 ± 292.7	980.4 ± 292.7	2.67**	1344.2 ± 310.4	1342.5 ± 310.4	2.73**
Iron (mg)	12.0 ± 2.8	11.7 ± 2.8	1.89 <sup>NS</sup>	17.8 ± 4.2	16.4 ± 4.2	1.54 <sup>NS</sup>	18.8 ± 4.9	16.8 ± 4.9	1.73 <sup>NS</sup>
Vitamin A (µg)	171.6 ± 126.4	167.8 ± 126.4	1.93 <sup>NS</sup>	351.7 ± 229.5	348.5 ± 229.5	1.59 <sup>NS</sup>	587.9 ± 580.7	580.3 ± 580.7	1.54 <sup>NS</sup>
Thiamin (mg)	1.2 ± 0.2	1.1 ± 0.2	0.98 <sup>NS</sup>	1.6 ± 0.3	1.5 ± 0.3	0.69 <sup>NS</sup>	1.7 ± 0.4	1.6 ± 0.4	0.56 <sup>NS</sup>
Niacin (mg)	10.2 ± 2.5	9.8 ± 2.5	1.12 <sup>NS</sup>	11.6 ± 2.8	11.0 ± 2.8	1.43 <sup>NS</sup>	12.0 ± 3.5	11.0 ± 3.5	1.76 <sup>NS</sup>
Folic acid (µg)	140.4 ± 32.3	139.1 ± 32.3	0.46 <sup>NS</sup>	204.0 ± 63.7	203.0 ± 63.7	0.51 <sup>NS</sup>	235.2 ± 66.2	233.4 ± 66.2	0.64 <sup>NS</sup>
Vitamin C (mg)	36.4 ± 9.7	34.4 ± 9.7	2.10**	98.0 ± 38.0	96.0 ± 38.0	2.44**	119.6 ± 90.3	117.6 ± 90.3	2.28**

LIG-Low income group, MIG-Middle income group, HIG- High income group

<sup>NS</sup>-Non significant \* Significant at 1% \*\* Significant at 5%

A significant ( $p \leq 0.01$ ) difference was found between the pre and post test scores of the subjects in all the income groups. Hence it can be concluded that there was gain in knowledge of the subjects after the nutrition counseling.

The data revealed that there was a significant ( $p \leq 0.05$ ) decrease in the intake of cereals, milk and milk products, fats and oils and meat and poultry in the subjects after nutrition counseling whereas the intake of pulses and legumes, green leafy vegetables, other vegetables, roots and tubers, fruits and sugar showed a non-significant change (Table 6).

The data revealed that after counseling there was a significant ( $p \leq 0.05$ ) decrease in the intake of various nutrients viz. energy, fat, carbohydrate, calcium and vitamin C in the obese subjects whereas a non significant change was found in the intake of protein, total dietary fibre, iron, vitamin A, thiamine, riboflavin, niacin and folic acid by the subjects. Significant decrease was observed in carbohydrate intake after nutrition counseling in all the income groups and it was mainly due to avoidance of refined foods, sugar and jaggery and fast foods and also inclusion of high fiber fruits and vegetables, whole wheat flour, dalia etc. in their daily diets as explained during nutrition counseling. Similarly, a decrease in the percent adequacy of the nutrients was observed after nutrition counseling (Table 7). A significant decrease was found in the intake of milk and milk products and fats and oils which might be another contributory factor in lowering the energy, calcium and fat intake after nutrition counseling but total fat and milk intake was found still higher than the suggested intake in MIG and LIG groups. Others workers also reported significant reduction

in fat intakes after nutrition counseling [7].

## Conclusion

It may be concluded that the overall prevalence of overweight and obesity in the women of Hoshiarpur was found to be 56.6, 67.3 and 70.6 percent in low, middle and high income group respectively. Percent adequacy of food groups including roots and tubers, fruits, milk & milk products and fats & oils was higher in obese subjects as compared to their normal counterparts in all the three income groups. Nutrition counseling of the subjects resulted in a gain in their knowledge and quantum of improvement, which further led to a decrease in the intake of cereals, milk & milk products, fats & oils and roots and tubers which resulted in significant decrease in energy, fat, carbohydrates, calcium, and vitamin C values by the obese subjects as compared to normal subjects in all the three income groups. Among various nutrients fat and energy intake were found to be associated with the risk factors of obesity so there is a need to create awareness regarding the intake of low fat, low carbohydrate and high fiber diet among the masses. Imparting knowledge about the benefits of physical activity will further help in reducing the risk factors of obesity. Therefore, it is recommended that physical activity is a prerequisite for controlling weight; hence activities like jogging, walking, exercise etc should become a part of daily routine.

## References

1. James WPT, Jackson-Leach R, Mhurchu CN,

- Kalamara E, Shayeghi M, Rigby NJ, Nishida C and Rodgers. A Overweight and obesity (high Body Mass Index). In Comparative quantification of health risks: global and regional burden of disease attributable to selected major risk factors: Ezzati M, Lopez AD, Rodgers A, Murray C J L (ed) Geneva, (World Health Organization). 2004. pp.497-596.
2. Popkin BM, Adair SL and Ng WS. Global nutrition transition and the pandemic of obesity in developing countries. *Nutr Rev.* 2012;70:3-21.
  3. Garg C, Khan SA, Ansari SH and Garg M. Prevalence of obesity in Indian women. *Obes Rev.* 2010;11:105-08.
  4. WHO expert consultant. Appropriate Body Mass Index for Asian populations and its implications for policy and intervention strategies. *Lancet.* 2004;363:157-63.
  5. Antonio G and Chiara A. A natural diet versus modern western diets? A new approach to prevent Well-Being Syndromes. *Digestive Dis Sci.* 2005; 50:1-6.
  6. Nam JH. Effect of weight control program on obesity degree and blood lipid levels among middle-aged obese women. *Korean J Food Nutr.* 2006;19:70-78. [Google Scholar].
  7. Borbora, M, Bhattacharyya, R, Borthakur, S. Impact of nutrition counseling on subjects suffering from cardiovascular disease. *Asian J. Home Sci.* 2009;3:164-66.
  8. Wadden TA, Webb VL, Moran CH, Bailer BA. Lifestyle modification for obesity: new developments in diet, physical activity, and behavior therapy. *Circulation.* 2012;125:1157-70. [PMC free article][PubMed] [Google Scholar].
  9. Kaur G. DietCal-A tool for dietary assessment and planning. 2014. Department of Dietetics, AIIMS, New Delhi.
  10. ICMR. Nutrient requirement and safe dietary intake for Indians. 2010. National Institute of Nutrition, Indian Council of Medical Research, Hyderabad, India.
  11. Kalsi G. Nutritional status, energy expenditure pattern and risk factors in obese Punjabi women. 1989. Ph.D. Dissertation, Punjab Agricultural University, Ludhiana.
  12. Kaur A. Consumption of fat and essential fatty acids by adult population of various socio-economic groups. 1995. M.Sc. thesis, Punjab Agricultural University, Ludhiana.
  13. Gupta N, Shah P, Goel K, Misra A, Rastogi K, Vikram NK, Kumari V, Pandey RM, Kondal D, Wasir JS, Bhardwaj S and Gulati S. Imbalanced dietary profile, anthropometry and lipids in urban Asian Indian adolescents and young adults. *J Am Coll Nutr.* 2010;2:81-91.
  14. Vijayalaxmi PM, Kasturiba B, Naik KR and Malagi U. Influence of fats and oils intake on the lipid profile of adults belonging to different income groups. *JAMA.* 2007;274:131-36.
  15. Geraz T. Study on nature and etiology of obesity and related degenerative disease among urban adults. 1996. M.Sc. Thesis, CCS Haryana Agricultural University, Hisar.
  16. Sangha JK, Pandher AK and Kochhar A. Anthropometric profile and adiposity in the obese Punjabi children and their parents. *J Human Ecol.* 2006;19:159-62.
  17. Kaur G, Singh SP and Singh A. Prevalence of overweight and obesity in urban and rural women of Punjab. *Human Biol Rev.* 2013. pp.306-13.
  18. Dhir R. Effect of dietary protein and exercise on body composition during weight loss regimen in adult women. 2012. M.Sc. thesis, Punjab Agricultural University, Ludhiana, India.
  19. Bennett C, Reed GW, Peters JC, Abumrad N, Sun M and Hill JO. Short term effects of dietary fat ingestion on energy expenditure and nutrient balance. *Am J Clin Nutr.* 1992;55:1071-77.
  20. Bala P. Anthropometry and nutrient adequacy of well-to-do school girls of 13-15 years from Punjab. 2000 M.Sc. Thesis, Punjab Agricultural University, Ludhiana.
  21. Gargiula J. Adolescents and obesity. *Family Practice.* 2002;19:304-09.
-

## Vegetable Pigments: Effect of Ph and Heat- A Review

E. Lakshmi<sup>1</sup>, E. Saraswathy<sup>2</sup>

### How to cite this article:

E. Lakshmi, E. Saraswathy. Vegetable Pigments: Effect of Ph and Heat- A Review. Int J Food Nutr Diet. 2019;7(1):41-44.

### Abstract

Commonly, vegetables are prepared at home on the basis of convenience and taste rather than nutrient losses. many studies have been conducted in the recent past to investigate the impact of cooking methods and Ph on the retention of pigments and stability of nutrients in coloured vegetables. The results of these studies vary widely leading the consumer to question the best method of preparing and cooking coloured vegetables, A literature search was conducted to identify studies that evaluated the effects of cooking techniques on the levels of Chlorophyll and carotenoids in vegetables The objective of this review is to evaluate the most recent studies and draw conclusions that will enable: (a) the consumer to identify the impact of cooking methods and Ph on the retention of pigments of coloured vegetables and (b) identify the critical phases during cooking, when colour and nutrients might be lost. The present study concentrated on chlorophyll and carotenoid pigments.

**Keywords:** Chlorophyll; Carotenoids; Cooking methods; Ph; Vegetables.

### Introduction

The bright colors of many vegetables contribute a lot to their attractiveness. The colours result from the pigments contained in their tissues. The chief pigments of vegetables and fruits may be classified as water insoluble and water soluble [1].

Chlorophyll pigments are largely insoluble in water and dominant in unripe fruits. they're held close to the cytomembrane in tiny bodies referred to as chloroplasts. Chlorophyll-a is intense blue green

in colour and chlorophyll-b is dull yellow green in colour. This pigment is present in green leafy vegetables, capsicum, beans, peas and chillies [2].

Carotenoids are widely distributed natural pigments accountable for the yellow, orange, and red colours of fruits and vegetables. Carotenoids are often divided into carotenes containing solely carbon and hydrogen, and xanthophylls made up of carbon, hydrogen, and oxygen [3]. Carotenoids owe their name to carrots (*Daucus carota*), and xanthophyll is derived from the Greek words for yellow and leaf. This pigment remains unaffected by heat or acidic medium however turns slightly blue in alkaline medium. Thus, the colour of the vegetables and fruits containing carotene remains unchanged on cooking [4].

Report suggests that 5 to 78% of the  $\beta$ -carotene is degraded when vegetables are cooked by different domestic methods. Considerable quantities of carotenoids may also be lost during household cooking of vegetables [5]. Thus, information on the possible losses of carotenoids from vegetables, during traditional cooking methods is of major importance. There are documented evidence on

**Author Affiliation:** <sup>1</sup>Assistant Professor, Department of Nutrition, SRM College of Nursing, SRM Institute of Science and Technology, SRM University, Kattankulathur, Chennai, Tamil Nadu 603203, India. <sup>2</sup>Assistant Professor, Department of Botany, Stella Maris College, Chennai, Tamil Nadu 600086, India.

**Corresponding Author:** E. Lakshmi, Assistant Professor, Department of Nutrition, SRM College of Nursing, SRM Institute of Science and Technology, SRM University, Kattankulathur, Chennai, Tamil Nadu 603203, India.

**E-mail:** [elakshmi20@gmail.com](mailto:elakshmi20@gmail.com)

loss of  $\beta$ -carotene after boiling, stewing, frying, blanching, and pressure cooking cooking.

### Methods of cooking [6]

**Boiling in water:** This is one of the common methods used in homes. The vegetables are washed, cut, cooked in excess of water for 20-30 minutes and the excess of water is rejected. This leads to considerable loss of water soluble vitamins and minerals.

**Steaming:** The vegetables are washed and cut and placed in vessels containing small amounts of water and steamed in a cooker under ordinary pressure for 20-30 minutes. The losses of nutrients are minimal in this method of cooking.

**Pressure cooking:** The vegetables are washed and cut and placed in vessels containing small amounts of water and cooked in steam under pressure for 10-15 minutes. The losses of nutrients are similar to those occurring in steaming under atmospheric pressure.

During boiling and steaming, the cells expand suddenly, and gases escape from the spaces in between the cells. The heat damages chloroplasts hence releases natural cell acids to turn green pigments into olive-green. This change occurs in the first few seconds of cooking, and the air pockets change the color of the chloroplasts. when these molecules are collapsed by the sudden rush of heat on to the vegetable they become brighter.

### B. Texture

Cooking in general affects the texture of all foodstuffs. The cellulose present in them becomes softer and this makes the foodstuffs also softer for example, cooked vegetables are much softer than raw. Cooking in an alkaline medium softens the cellulose much faster and gives the foodstuffs a mushy appearance whereas cooking in an acidic medium makes the texture hard and such foodstuffs have to be cooked for a longer period. The texture of the food, after cooking is also dependent on the method which is used for cooking [7]. Moist heat methods i.e. boiling, steaming, pressure cooking and stewing makes the foodstuffs soft and tender.

The objective of the present study was to determine the effect of different domestic cooking methods on chlorophyll and carotenoids retention in string beans (*Phaseolus vulgaris*, L.) and carrots (*Daucus carota* L.) the commonly consumed vegetables. chlorophyll and  $\beta$ -carotene were chosen for the present studies since they form the

main pigments contributing to the health benefits of Vitamin A in human. The study was aimed to find the effect of culinary treatments like steaming boiling and prolonged cooking at different cooking durations (4 and 8 min) on the chlorophyll and carotenoid pigments of string beans (*Phaseolus vulgaris*, L.) and carrots (*Daucus carota* L.) in the first phase followed by change in ph of the cooking water in the second phase.

### Sample preparation

Upon arrival at the department of Nutrition and Dietetics laboratory, the fresh vegetables were washed under tap water and excessive water dripped off. Edible portions (600 g) of the vegetables were separated and cut to small pieces, in edible parts were removed. The vegetables were divided into five portions with 100g for each application. One portion was retained raw and others were subjected to different cooking treatments by altering the Ph. The colour change, texture and doneness were observed. The variations were compared with control (uncooked).

### Cooking Treatments

- a. Boiling in open pan for 10 minutes followed by closing
- b. Prolonged cooking
- c. Steaming
- d. Addition of vinegar (1 tsp)
- e. Addition of alkali- Baking soda (1/2 tsp)
- f. Control (uncooked).

### Boiling

Vegetable (100g) was added to 250 ml of water that has just reached the boiling temperature in a stainless steel pot, cooked for 4 or 8 min, Samples were drained off and cooled rapidly.

### Prolonged cooking

Vegetable (100 g) was added to 500 ml of water that has just reached the boiling temperature in a stainless steel pot, cooked for more than 10 minutes. Samples were drained off and cooled rapidly.

### Steaming

Vegetable (100g) was placed on tray in a steam cooker covered with a lid and steamed over boiling water for 7-10 min under atmospheric pressure. The samples were rapidly cooled.

### a. Green vegetable- Chlorophyll

All the variations were compared to the raw, cut vegetables.

Boiling of string beans in open pan for 10 minutes followed by closing resulted in bright green colour. The heat breaks down the cell structure in the pigment. This results in change of color. Cooking vegetables in an uncovered pan for few minutes helps to eliminate volatile acids and prevents formation of Pheophytin that results in formation of olive-green colour [8]. Boiling is more similar to the steaming but still had a brighter appearance compared to control (Fig. 1). Chlorophyll pigments are sensitive to heat and cooking. The magnesium atom present in the centre is easily replaced by hydrogen ions or other metal ions. As the cooking process continues (Prolonged cooking) the colour of the string beans changed from the bright green colour to the olive green. Excessive heating exposes the plants own acids to the chlorophyll which in turn makes the cooked product dull green [9]. Steaming turned string beans to a dark green colour. After few minutes the beans was a little bit darker and a more distinct green in colour compared to control.

In phase II when the PH was altered by addition of vinegar the colour of string beans turned almost immediately to a much lighter shade of green. After 5 minutes the color was considerably lighter. Magnesium present in the structure of chlorophyll is rather easily displaced by hydrogen molecule when heated in the presence of organic acids. A pale greenish grey compound known as Pheophytin-a or an olive-green Pheophytin-b results. Chlorophyll-a is more readily converted to pheophytin than chlorophyll-b [10]. when the PH was reduced to alkaline phase by adding baking soda there was

not much immediate reactions After 5 minutes the colour was slightly darker than the control. The phetyl and methyl groups of chlorophyll are displaced and a bright green water-soluble sodium salt of chlorophyllin compound is formed.

### b. Yellow-orange vegetable-carotenoids

The pigments that color these vegetables are deep orange colour called carotenoids. They are very stable pigments. All the variations were compared to the raw, peeled, carrots. Steaming resulted in darker and brighter orange colour of carrots. They appeared to have similar color to that of raw carrots. Boiling resulted in more yellow pigments than the raw carrots. Not only were they lighter, but they seemed more yellowish in color than orange (Fig. 2). The loss in intensity of colour is not only due to oxidation of the unsaturation of carotene but also shift from trans to cis form [11]. Keeping a lid on the cooking vessel had some advantage as there was about more retention of carotenes in these vegetable. seem more yellow than the raw carrots. Not only are they lighter, but they seem more yellowish in color than orange.

In phase II when the Ph was altered by addition of vinegar there was immediate change in the colour. The carrots reduced to a lighter shade compared to control. when the PH was reduced to alkaline phase by adding baking soda the colour turned darker. The colour is little affected by acid, alkali and the volume of the water. According to Nunn M et al. 2006, more investigations are needed to provide a better understanding of the oxidative phenomena of carotenoids during cooking since different vegetables have different chemical and physical characteristics.



Fig. 1: Chlorophyll pigments - Beans



Fig. 2: Carotenoid pigments - carrots

### Conclusion

The boiling and steaming methods on chlorophyll had the least color change, although they seemed little brighter. A second color change occurred in response to acidic water: The magnesium ion in the center of the chlorophyll molecule is replaced with a hydrogen atom, causing the green to dull. Chlorophyll-a becomes grey-green pheophytin-a, and chlorophyll-b turns into yellowish pheophytin-b. When the boiling water was slightly alkaline, then chlorophyll retained green. Accordingly boiling in open pan and latter closing can be followed for glv to maintain the natural colour of chlorophyll. Colors of carotenoids did not fade much in response to heat. Some change occurred, in carrots. The taproots change from red-orange to more yellow when cooked. Boiling resulted in intense yellow than the raw carrots. Steaming resulted in brighter orange than the rest of the carrots. There was an immediate change in the carrot when acid was added It turned lighter, while addition of baking soda led to little bit darker. cooking methods and Ph had a strong impact on the pigment retention of pigments in green and yellow vegetables. However, upon considering all the evidence steaming can be suggested as the best cooking method to preserve the most of the phenolic compounds, particularly flavonoids and glucosinolates.

### Key Messages

Cooking in general affects the texture of all foodstuffs Considerable quantities of vegetable pigments may also be lost during household cooking methods.

### References

1. Turkmen N, Poyrazoglu ES, Sari F, Velioglu YS. Effects of cooking methods on chlorophylls, pheophytins and colour of selected green vegetables Int. J. Food Sci. Technol. 2006;41(3):281-88.
2. Miglio C, Chiavaro E, Visconti A, Fogliano V., Pellegrini N. Effects of Different Cooking Methods on Nutritional and Physicochemical Characteristics of Selected Vegetables J. Agric. Food Chem. 2008;56(1):139-47. DOI: 10.1021/jf072304b.
3. DC Murador, DT da Cunha, VV de Rosso. Effects of cooking techniques on vegetable pigments: A meta-analytic approach to carotenoid and anthocyanin levels. Food Research International. 2014;65:177-83.
4. Turkmen N, Sari F, Velioglu YS. The effect of cooking methods on total phenolics and antioxidant activity of selected green vegetables. Food Chemistry, 2005; 93(4):713-18.
5. Archana GN, Prakash J, Asha MR, Chand N. Effects of processing on pigments of certain selected vegetables. J. Food Quality. 1995;18:91-101.
6. Miglio C, Chiavaro E, Visconti A, et al. Effects of different cooking methods on nutritional and physicochemical characteristics of selected vegetables, J. Agric. Food Chem. 2008;56:139-47. DOI: 10.1021/jf072304b.
7. Bao B. Chang KC. Carrot juice color, carotenoids, and nonstarchy polysaccharides as affected by processing conditions J. Food Sci. 1994;59:1155- 58.
8. Hiemori ME. Koh, Mitchell, AE. Influence of cooking on anthocyanins in black rice (*Oryza sativa* L.japonicavar. SBR). J. Agric. Food Chem. 2009; 57(5):1908-14.
9. Palermo M, Pellegrini N, Fogliano V . The effect of cooking on the phytochemical content of vegetables. J Sci Food Agric. 2014;94:1057-1070, DOI 10.1002/jsfa.6478.

10. Chen BH, Peng HY, Chen HE. Changes of carotenoids, color and vitamin A contents during processing of carrot juice J. Agric. Food Chem. 1995; 43:1912- 18.
  11. Nunn M, Giraud D, Parkhurst A, Hamouz F, Driskell J. Effects of cooking methods on sensory qualities and carotenoid retention in selected vegetables. J Food Qual. 2006;29:445-57 doi: 10.1111/j.1745-4557.2006.00071.x.
- 

## REDKART.NET

(A product of RF Library Services (P) Limited)  
(Publications available for purchase: Journals, Books, Articles and Single issues)  
(Date range: 1967 to till date)

The Red Kart is an e-commerce and is a product of RF Library Services (P) Ltd. It covers a broad range of journals, Books, Articles, Single issues (print & Online-PDF) in English and Hindi languages. All these publications are in stock for immediate shipping and online access in case of online.

**Benefits of shopping online are better than conventional way of buying.**

1. Convenience.
2. Better prices.
3. More variety.
4. Fewer expenses.
5. No crowds.
6. Less compulsive shopping.
7. Buying old or unused items at lower prices.
8. Discreet purchases are easier.

URL: [www.redkart.net](http://www.redkart.net)

Manuscripts must be prepared in accordance with “Uniform requirements for Manuscripts submitted to Biomedical Journal” developed by international committee of medical Journal Editors

### Types of Manuscripts and Limits

Original articles: Up to 3000 words excluding references and abstract and up to 10 references.

Review articles: Up to 2500 words excluding references and abstract and up to 10 references.

Case reports: Up to 1000 words excluding references and abstract and up to 10 references.

### Online Submission of the Manuscripts

Articles can also be submitted online from [http://rfppl.co.in/customer\\_index.php](http://rfppl.co.in/customer_index.php).

1) First Page File: Prepare the title page, covering letter, acknowledgement, etc. using a word processor program. All information which can reveal your identity should be here. use text/rtf/doc/PDF files. Do not zip the files.

2) Article file: The main text of the article, beginning from Abstract till References (including tables) should be in this file. Do not include any information (such as acknowledgement, your name in page headers, etc.) in this file. Use text/rtf/doc/PDF files. Do not zip the files. Limit the file size to 400 Kb. Do not incorporate images in the file. If file size is large, graphs can be submitted as images separately without incorporating them in the article file to reduce the size of the file.

3) Images: Submit good quality color images. Each image should be less than 100 Kb in size. Size of the image can be reduced by decreasing the actual height and width of the images (keep up to 400 pixels or 3 inches). All image formats (jpeg, tiff, gif, bmp, png, eps etc.) are acceptable; jpeg is most suitable.

Legends: Legends for the figures/images should be included at the end of the article file.

If the manuscript is submitted online, the contributors' form and copyright transfer form has to be submitted in original with the signatures of all the contributors within two weeks from submission. Hard copies of the images (3 sets), for articles submitted online, should be sent to the journal office at the time of submission of a revised manuscript. Editorial office: Red Flower Publication Pvt. Ltd., 48/41-42, DSIDC, Pocket-II, Mayur Vihar Phase-I, Delhi - 110 091, India, Phone: 91-11-22754205, 45796900, 22756995. E-mail: [author@rfppl.co.in](mailto:author@rfppl.co.in). Submission page: [http://rfppl.co.in/article\\_submission\\_system.php?mid=5](http://rfppl.co.in/article_submission_system.php?mid=5).

### Preparation of the Manuscript

The text of observational and experimental articles should be divided into sections with the headings: Introduction, Methods, Results, Discussion, References, Tables, Figures, Figure legends, and Acknowledgment. Do not make subheadings in these sections.

### Title Page

The title page should carry

- 1) Type of manuscript (e.g. Original article, Review article, Case Report)
- 2) The title of the article, should be concise and informative;
- 3) Running title or short title not more than 50 characters;
- 4) The name by which each contributor is known (Last name, First name and initials of middle name), with his or her highest academic degree(s) and institutional affiliation;
- 5) The name of the department(s) and institution(s) to which the work should be attributed;
- 6) The name, address, phone numbers, facsimile numbers and e-mail address of the contributor responsible for correspondence about the manuscript; should be mentioned.
- 7) The total number of pages, total number of photographs and word counts separately for abstract and for the text (excluding the references and abstract);
- 8) Source(s) of support in the form of grants, equipment, drugs, or all of these;
- 9) Acknowledgement, if any; and
- 10) If the manuscript was presented as part at a meeting, the organization, place, and exact date on which it was read.

### Abstract Page

The second page should carry the full title of the manuscript and an abstract (of no more than 150 words for case reports, brief reports and 250 words for original articles). The abstract should be structured and state the Context (Background), Aims, Settings and Design, Methods and Materials, Statistical analysis used, Results and Conclusions. Below the abstract should provide 3 to 10 keywords.

## Introduction

State the background of the study and purpose of the study and summarize the rationale for the study or observation.

## Methods

The methods section should include only information that was available at the time the plan or protocol for the study was written such as study approach, design, type of sample, sample size, sampling technique, setting of the study, description of data collection tools and methods; all information obtained during the conduct of the study belongs in the Results section.

Reports of randomized clinical trials should be based on the CONSORT Statement (<http://www.consort-statement.org>). When reporting experiments on human subjects, indicate whether the procedures followed were in accordance with the ethical standards of the responsible committee on human experimentation (institutional or regional) and with the Helsinki Declaration of 1975, as revised in 2000 (available at [http://www.wma.net/e/policy/17-c\\_e.html](http://www.wma.net/e/policy/17-c_e.html)).

## Results

Present your results in logical sequence in the text, tables, and illustrations, giving the main or most important findings first. Do not repeat in the text all the data in the tables or illustrations; emphasize or summarize only important observations. Extra or supplementary materials and technical details can be placed in an appendix where it will be accessible but will not interrupt the flow of the text; alternatively, it can be published only in the electronic version of the journal.

## Discussion

Include summary of key findings (primary outcome measures, secondary outcome measures, results as they relate to a prior hypothesis); Strengths and limitations of the study (study question, study design, data collection, analysis and interpretation); Interpretation and implications in the context of the totality of evidence (is there a systematic review to refer to, if not, could one be reasonably done here and now?, What this study adds to the available evidence, effects on patient care and health policy, possible mechanisms)? Controversies raised by this study; and Future research directions (for this particular research collaboration, underlying mechanisms, clinical research). Do not repeat in detail data or other

material given in the Introduction or the Results section.

## References

List references in alphabetical order. Each listed reference should be cited in text (not in alphabetic order), and each text citation should be listed in the References section. Identify references in text, tables, and legends by Arabic numerals in square bracket (e.g. [10]). Please refer to ICMJE Guidelines ([http://www.nlm.nih.gov/bsd/uniform\\_requirements.html](http://www.nlm.nih.gov/bsd/uniform_requirements.html)) for more examples.

### Standard journal article

[1] Flink H, Tegelberg Å, Thörn M, Lagerlöf F. Effect of oral iron supplementation on unstimulated salivary flow rate: A randomized, double-blind, placebo-controlled trial. *J Oral Pathol Med* 2006; 35: 540-7.

[2] Twetman S, Axelsson S, Dahlgren H, Holm AK, Källestål C, Lagerlöf F, et al. Caries-preventive effect of fluoride toothpaste: A systematic review. *Acta Odontol Scand* 2003; 61: 347-55.

### Article in supplement or special issue

[3] Fleischer W, Reimer K. Povidone iodine antiseptics. State of the art. *Dermatology* 1997; 195 Suppl 2: 3-9.

### Corporate (collective) author

[4] American Academy of Periodontology. Sonic and ultrasonic scalers in periodontics. *J Periodontol* 2000; 71: 1792-801.

### Unpublished article

[5] Garoushi S, Lassila LV, Tezvergil A, Vallittu PK. Static and fatigue compression test for particulate filler composite resin with fiber-reinforced composite substructure. *Dent Mater* 2006.

### Personal author(s)

[6] Hosmer D, Lemeshow S. Applied logistic regression, 2nd edn. New York: Wiley-Interscience; 2000.

### Chapter in book

[7] Nauntofte B, Tenovou J, Lagerlöf F. Secretion and composition of saliva. In: Fejerskov O,

Kidd EAM, editors. *Dental caries: The disease and its clinical management*. Oxford: Blackwell Munksgaard; 2003. p. 7-27.

### No author given

[8] World Health Organization. *Oral health surveys - basic methods*, 4th edn. Geneva: World Health Organization; 1997.

### Reference from electronic media

[9] National Statistics Online – Trends in suicide by method in England and Wales, 1979-2001. [www.statistics.gov.uk/downloads/theme\\_health/HSQ20.pdf](http://www.statistics.gov.uk/downloads/theme_health/HSQ20.pdf) (accessed Jan 24, 2005): 7-18. Only verified references against the original documents should be cited. Authors are responsible for the accuracy and completeness of their references and for correct text citation. The number of reference should be kept limited to 20 in case of major communications and 10 for short communications.

More information about other reference types is available at [www.nlm.nih.gov/bsd/uniform\\_requirements.html](http://www.nlm.nih.gov/bsd/uniform_requirements.html), but observes some minor deviations (no full stop after journal title, no issue or date after volume, etc).

### Tables

Tables should be self-explanatory and should not duplicate textual material.

Tables with more than 10 columns and 25 rows are not acceptable.

Table numbers should be in Arabic numerals, consecutively in the order of their first citation in the text and supply a brief title for each.

Explain in footnotes all non-standard abbreviations that are used in each table.

For footnotes use the following symbols, in this sequence: \*, †, ‡, §.

### Illustrations (Figures)

Graphics files are welcome if supplied as Tiff, EPS, or PowerPoint files of minimum 1200x1600 pixel size. The minimum line weight for line art is 0.5 point for optimal printing.

When possible, please place symbol legends below the figure instead of to the side.

Original color figures can be printed in color at the editor's and publisher's discretion provided the author agrees to pay.

Type or print out legends (maximum 40 words, excluding the credit line) for illustrations using double spacing, with Arabic numerals corresponding to the illustrations.

### Sending a revised manuscript

While submitting a revised manuscript, contributors are requested to include, along with single copy of the final revised manuscript, a photocopy of the revised manuscript with the changes underlined in red and copy of the comments with the point to point clarification to each comment. The manuscript number should be written on each of these documents. If the manuscript is submitted online, the contributors' form and copyright transfer form has to be submitted in original with the signatures of all the contributors within two weeks of submission. Hard copies of images should be sent to the office of the journal. There is no need to send printed manuscript for articles submitted online.

### Reprints

Journal provides no free printed reprints, however a author copy is sent to the main author and additional copies are available on payment (ask to the journal office).

### Copyrights

The whole of the literary matter in the journal is copyright and cannot be reproduced without the written permission.

### Declaration

A declaration should be submitted stating that the manuscript represents valid work and that neither this manuscript nor one with substantially similar content under the present authorship has been published or is being considered for publication elsewhere and the authorship of this article will not be contested by any one whose name (s) is/are not listed here, and that the order of authorship as placed in the manuscript is final and accepted by the co-authors. Declarations should be signed by all the authors in the order in which they are mentioned in the original manuscript. Matters appearing in the Journal are covered by copyright but no objection will be made to their reproduction provided permission is obtained from the Editor prior to publication and due acknowledgment of the source is made.

### Approval of Ethics Committee

We need the Ethics committee approval letter from an Institutional ethical committee (IEC) or an institutional review board (IRB) to publish your Research article or author should submit a statement that the study does not require ethics approval along with evidence. The evidence could either be consent from patients is available and there are no ethics issues in the paper or a letter from an IRB stating that the study in question does not require ethics approval.

### Abbreviations

Standard abbreviations should be used and be spelt out when first used in the text. Abbreviations should not be used in the title or abstract.

### Checklist

- Manuscript Title
- Covering letter: Signed by all contributors
- Previous publication/ presentations mentioned, Source of funding mentioned
- Conflicts of interest disclosed

### Authors

- Middle name initials provided.
- Author for correspondence, with e-mail address provided.
- Number of contributors restricted as per the instructions.
- Identity not revealed in paper except title page (e.g.name of the institute in Methods, citing previous study as 'our study')

### Presentation and Format

- Double spacing
- Margins 2.5 cm from all four sides
- Title page contains all the desired information. Running title provided (not more than 50 characters)
- Abstract page contains the full title of the manuscript
- Abstract provided: Structured abstract provided for an original article.
- Key words provided (three or more)
- Introduction of 75-100 words
- Headings in title case (not ALL CAPITALS).

References cited in square brackets

- References according to the journal's instructions

### Language and grammar

- Uniformly American English
- Abbreviations spelt out in full for the first time. Numerals from 1 to 10 spelt out
- Numerals at the beginning of the sentence spelt out

### Tables and figures

- No repetition of data in tables and graphs and in text.
- Actual numbers from which graphs drawn, provided.
- Figures necessary and of good quality (color)
- Table and figure numbers in Arabic letters (not Roman).
- Labels pasted on back of the photographs (no names written)
- Figure legends provided (not more than 40 words)
- Patients' privacy maintained, (if not permission taken)
- Credit note for borrowed figures/tables provided
- Manuscript provided on a CDROM (with double spacing)

### Submitting the Manuscript

- Is the journal editor's contact information current?
- Is the cover letter included with the manuscript? Does the letter:
  1. Include the author's postal address, e-mail address, telephone number, and fax number for future correspondence?
  2. State that the manuscript is original, not previously published, and not under concurrent consideration elsewhere?
  3. Inform the journal editor of the existence of any similar published manuscripts written by the author?
  4. Mention any supplemental material you are submitting for the online version of your article. Contributors' Form (to be modified as applicable and one signed copy attached with the manuscript)

## International Journal of Food, Nutrition and Dietetics

### Library Recommendation Form

If you would like to recommend this journal to your library, simply complete the form below and return it to us. Please type or print the information clearly. We will forward a sample copy to your library, along with this recommendation card.

#### Please send a sample copy to:

Name of Librarian

Name of Library

Address of Library

#### Recommended by:

Your Name/ Title

Department

Address

#### Dear Librarian,

I would like to recommend that your library subscribe to the International Journal of Food, Nutrition and Dietetics. I believe the major future uses of the journal for your library would provide:

1. useful information for members of my specialty.
2. an excellent research aid.
3. an invaluable student resource.

**I have a personal subscription and understand and appreciate the value an institutional subscription would mean to our staff.**

Should the journal you're reading right now be a part of your University or institution's library? To have a free sample sent to your librarian, simply fill out and mail this today!

Stock Manager

Red Flower Publication Pvt. Ltd.

48/41-42, DSIDC, Pocket-II

Mayur Vihar Phase-I

Delhi - 110 091(India)

Phone: Phone: 91-11-45796900, 22754205, 22756995, Cell: +91-9821671871

E-mail: sales@rfppl.co.in

## Red Flower Publication (P) Ltd.

*Presents its Book Publications for sale*

- |  |               |
|--|---------------|
| <b>1. Biostatistics Methods for Medical Research (2019)</b><br><i>by Sanjeev Sarmukaddam</i>   | INR549/USD44  |
| <b>2. MCQs in Medical Physiology (2019)</b> <i>by Bharati Mehta &amp; Bharti Bhandari Rathore</i>                                    | INR300/USD29  |
| <b>3. Synopsis of Anesthesia (2019)</b> <i>by Lalit Gupta MBBS &amp; Bhavna Gupta MBBS</i>   | INR1195/USD95 |
| <b>4. Shipping Economics (2018)</b> <i>by D. Amutha, Ph.D.</i>   | INR345/USD27  |
| <b>5. Breast Cancer: Biology, Prevention and Treatment (2015)</b><br><i>by Rana P. Singh, Ph.D. &amp; A. Ramesh Rao, Ph.D. (JNU)</i> | INR395/USD100 |
| <b>6. Child Intelligence (2005)</b> <i>by Rajesh Shukla, MD.</i>   | INR150/USD50  |
| <b>7. Pediatric Companion (2001)</b> <i>by Rajesh Shukla, MD.</i>  | INR250/USD50  |

### Order from

**Red Flower Publication Pvt. Ltd.**

48/41-42, DSIDC, Pocket-II

Mayur Vihar Phase-I

Delhi - 110 091(India)

Mobile: 8130750089, Phone: 91-11-45796900, 22754205, 22756995

E-mail: sales@rfppl.co.in

## Special Note!

Please note that our all Customers, Advertisers, Authors, Editorial Board Members and Editor-in-chief are advised to pay any type of charges against Article Processing, Editorial Board Membership Fees, Postage & Handling Charges of author copy, Purchase of Subscription, Single issue Purchase and Advertisement in any Journal directly to Red Flower Publication Pvt. Ltd.

Nobody is authorized to collect the payment on behalf of Red Flower Publication Pvt. Ltd. and company is not responsible of respective services ordered for.

Red Flower Publication Pvt. Ltd.

## CAPTURE YOUR MARKET

*For advertising in this journal*

Please contact:

**International print and online display advertising sales**

*Advertisement Manager*

Phone: 91-11-22756995, 22754205, 45796900, Cell: +91-9821671871

E-mail: sales@rfppl.co.in

**Recruitment and Classified Advertising**

*Advertisement Manager*

Phone: 91-11-22756995, 22754205, 45796900, Cell: +91-9821671871

E-mail: sales@rfppl.co.in