

A Study of Levels of Plasma Vitamin B12, Folate, and Homocysteine In Cataract Patients

Anand Reddi*, Meena Lakshman**

*Assistant Professor, Department of Ophthalmology, Kerala Medical College and Hospital, Mangode, Palakkad Dt, Kerala -679503. **Assistant Professor, Department of Ophthalmology, KMCT Medical College and Hospital, Kozhikode, Kerala -673602.

Abstract

Background: Elevated plasma Homocysteine level is an independent risk factor for age related (senile) cataract. Certain nutritional deficiencies, in particular Folate, Vitamin B12, Vitamin B6 relate inversely with Homocysteine level. *Aim:* This study was undertaken to evaluate the plasma level of Vitamin B12, Folate, and Homocysteine of cataract patients and to study the interplay between them. *Methods:* This study included 50 cataract patients and 30 age matched control subjects. Samples were collected from cataract patients admitted in Ophthalmology ward prior to cataract surgery. Assays were carried out by using well-established methods for Homocysteine, Folate, Vitamin B12 by Chemiluminescence. *Results:* Serum Homocysteine level is significantly increased in cataract patients when compared with control ($p < 0.001$). There was a significant decrease in the level of Folate as compared with control ($p < 0.001$). There was a negative correlation between Homocysteine vs. Vitamin B12 ($p < 0.01$) and Folate ($p < 0.01$) in the Cataract patients. *Conclusion:* An increased plasma Homocysteine level is associated with decreased plasma levels of Folate and Vitamin B12 in Cataract patients, which might have a possible role in the root cause of cataract pathogenesis.

Keywords: Homocysteine; Senile Cataract; Vitamin B12; Folate.

Introduction

Cataract may be defined as any light scatter opacity in the lens, not necessarily with any demonstrable effect on vision. Cataract that is significant enough to impair vision is the leading cause of blindness worldwide [1]. Senile cataract, which is most common in south India, describes any cataract that occurs after the age of 45 and has no evident cause. Age related cataract is a multifactorial disease in which genetic, environmental, socio-economic and biochemical factors may act synergistically. A number of indicators of poor nutrition have been found to be associated with increased risk of cataract in India [2,3]. Elevated Homocysteine levels are seen in various eye diseases such as exfoliation syndrome, glaucoma, and cataract [4,5]. It has been proposed that Homocysteine being a putative oxidant is involved in the pathogenesis of endothelial cell

injury and atherosclerotic vascular disease [6,7,8]. Vitamin B12 (Vit B12) and Folate are involved in the metabolism of Homocysteine in methylating it to Methionine. Vit B12 and Folate has strong protective effect against formation of Cataractogenesis [9, 10]. So, present study was under taken to assess the Homocysteine levels in age related cataract in relation to the levels of Vit B12 and Folate.

Methods

This study included 50 cataract patients and 30 age matched control subjects. Samples were collected

Reprint Request: Anand Reddi, Assistant Professor, Department of Ophthalmology, Kerala Medical College and Hospital, Mangode, Palakkad Dt, Kerala -679503.

E-mail: thesisusm@gmail.com

Recived on 12.12.2016, **Accepted on** 13.12.2016

from cataract patients admitted in Ophthalmology ward at , Kerala Medical College and Hospital, prior to cataract surgery. Cataract patients were selected based on the vision less than 6/18 and visible opacity in the lens. All these patients were above 50 years of age. They were normotensive and non-diabetic as their blood pressure was normal and random blood glucose was below 140mg%. Similarly controls were non-diabetic and normotensive . Both patients and control subjects were not on vitamin, mineral, any drug or such supplementation. Random blood samples were collected and non-hemolysed plasma was used for all biochemical parameters. . Assays were carried out by using well-established and sensitive methods for Homocysteine, Folate, Vitamin B12 by Chemiluminescence (Immolute analyzer) using reagent kits obtained from DPC [11]. Protein estimation was done by Biuret method [12], Albumin by BCG method [13] and Random blood glucose by GOD/POD method [14] in the Dadebehring Autoanalyser. The reagent kits for total protein and albumin were obtained from Biosystems S.A.(Costa Brava 30, Barcelona, Spain) and that for blood glucose estimation was obtained from Reckon Diagnostics P.Ltd (Gorwa,Baroda-390 016, India).

Statistical analysis was done by using student's 't' test for comparison between the groups and correlation was done by Pearson correlation in Microsoft excel and P values < 0.05 were considered as statistically significant.

Results

Our present study aimed at finding the levels of Homocysteine, Folate, and Vitamin B12 in senile cataract patients. We selected 40 cataract patients and 20 controls. Table 1 shows comparisons of Homocysteine, Vit B12 and Folate levels in cataract patients with control subjects. Statistically significant increase in the level of Homocysteine ($p < 0.001$) was found in cataract patients as compared to control subjects. Also there was a significantly decrease in the level of Folate ($p < 0.001$) and random blood glucose ($p < 0.05$) in cataract subjects as compared with controls. Table 2 showed the pair wise correlation analysis in cataract patients showed following results. Hcy was negatively correlated with Vit B12 ($r = -0.68$) and Folate ($r = -0.47$).

Table 1: Plasma levels of Homocysteine, Vitamin B12, Folate, Total Protein, Albumin and Random Blood Glucose in control subjects and Cataract patients

Parameters	Control (n = 30)	Case (n = 50)
Age	57.9 ± 6.82	59.85 ± 10.52
Homocysteine (µmol/L)	5.44 ± 3.20	28.06 ± 8.92**
Vitamin B12 (pg/ml)	279.85 ± 91.00	201.8 ± 178.60
Folate (ng/ml)	7.06 ± 2.33	2.42 ± 1.73**
Total Protein (gm/dl)	7.65 ± 0.47	7.69 ± 0.58
Albumin (gm/dl)	4.21 ± 0.39	4.24 ± 0.34
Random blood glucose (mg/dl)	97.3 ± 5.77	93.00 ± 15.8*

Values are expressed as mean ± SD; **P value < 0.001; *P value < 0.05

Table 2: Pairwise correlation analysis in cataract patients

Pairwise correlation of parameters	R value
Homocysteine and Vit B12	-0.66**
Homocysteine and Folate	-0.45*

**P value < 0.001; *P value < 0.001

Discussion

Homocysteine is derived from the dietary methionine. In normal person Homocysteine is metabolized to Methionine with the help of Folate and Vit B12. It may also be converted to cystathionine by cystathionine -b-synthase. Normal level of Homocysteine is 5-12 µmol/L. Under certain circumstances Homocysteine level is increased in blood. Those are 1) Vit B12, Folate deficiency and

heterozygous or homozygous trait for Methyl-tetrahydrofolate reductase 2) Heterozygous or homozygous trait for cystathionine -β-synthase activity 3) Renovascular stenosis and volume retention. In our study we found an increased level of Homocysteine, which correlate with age [15], Folate and Vit B12 deficiency [16]. It is well known that hyper homocysteinemia produces oxidative stress by generating reactive oxygen species spontaneously [17]. Sulfhydryl group of Homocysteine is believed to act catalytically with ferric and cupric ions in a

mixed function oxidation system to generate hydrogen peroxide, oxygen radicals [18,19]. Superoxide anion, hydrogen peroxide, and hydroxyl radical are produced during auto-oxidation of Homocysteine [20,21]. These could promote lipid peroxidation and damage vascular endothelium. Hyperhomocysteinemia is an independent graded risk factor for atherosclerotic disease in coronary, cerebral and peripheral blood vessels [8].

Contemporary hypothesis considers the oxidative stress as a crucial event in age related processes as well as in the age related cataract. Sulfhydryl group of lens protein (Crystallins) is the target of enhanced oxidative stress seen in senile Cataract [22]. Recently, it has been shown that post translational modification of lens proteins play crucial role in the formation of age related cataract and also found that oxidized amino acids accumulate in cataractous lenses which are probably due to altered redox balance, thus contribute to lens opaciation [23].

In our study, we found there was a significant decrease in Folate level in cataract patients as compared with controls. Vit B12 was also decreased but not statistically significant. This is in accordance with previous study, which reveals low Folate as the strong determinant of Hyperhomocysteinemia in older age [16]. In the present study, Homocysteine level correlates negatively with Vit B12 and Folate levels which is in agreement with previous studies proving low Folate, Vit B12, Vit B6 and older age were all independently associated with elevated Homocysteine [16]. Vit B12, Folate supplements has strong protective influence on reducing the prevalence of Cataractogenesis [9].

Accumulating evidences suggest that, prolonged hyperglycemia seen in type 2 diabetes mellitus enhances the glycation of proteins and may be the root cause of cataract formation [24,25]. But in our study, random blood glucose levels of patients were within the reference range and even it showed statistically significant decrease than control group. This suggests that elevated Homocysteine level even in the absence of hyper glycemia or any other virtual metabolic and nutritional disorder as evident by total protein and albumin levels, may play a possible role in the cataract formation. Hyperhomocysteinemia with low Vit B12 and Folate levels in older age group can be independent risk factor for Senile Cataract, which might be mediated through oxidative stress. Further studies are needed to unravel the role of oxidative stress arising from elevated Homocysteine level in older age groups in senile cataract formation.

References

1. Thylfors B, Negrel AD, Pararajasegaram R, Dadzie KY. Global data on blindness. *Bull World Health Organ* 1995; 73:115-21.
2. Livingston PM, Carson CA, Taylor HR. The epidemiology of cataract: a review of the literature. *Ophthalmic epidemiol* 1995; 2:151-64.
3. Sarma U, Brunner E, Evans J, Wormald R. Nutrition and the epidemiology of cataract and age related maculopathy. *Eur J Clin Nutr* 1994; 48:1-8.
4. Puustjarvi T, Blomster H, Kontkanen M, Punnonen K, Terasvirte M. Plasma and aqueous humour levels of homocysteine in exfoliation syndrome. *Graefe's Arch Clin Experimental ophthalmol* 2004; 242(9):749-54.
5. Cumurcu T, Salvin S, Ardin. Serum homocysteine, vitamin B12, Folic acid in different types of glaucoma. *BMC Ophthalmol* 2006; 6:1-6.
6. McCully KS. Homocysteine theory of Arteriosclerosis: development and current status. *Atheroscler Rev* 1983; 11:157-246.
7. Welch GN, Loscalzo J. Homocysteine, atherothrombosis. *N Engl J Med* 1998; 338: 1042-50.
8. Boushey CJ, Bertesford SA, Omenn GS, Motulsky AG. A quantitative assessment of plasma Homocysteine as a risk factor for vascular disease. Probable benefits of increasing folic acid intakes. *JAMA* 1995; 274:1049-57
9. Kuzniarz M, Mitchell P, Cumming RG, Flood VM. Use of vitamin supplements and cataract: the Blue Mountains Eye Study. *Am J Ophthal* 2001; 132(1):19-26.
10. Bergman B, Nilsson-Ehle H, Sjostrand J. Ocular changes, risk markers for eye disorders and effects of cataract surgery in elderly people: a study of an urban Swedish population followed from 70 to 97 years of age. *Acta Ophthalmologica Scandinavia* 2004; 82(2):166-74.
11. Estimation of Vit.B12, Folic acid, Homocysteine on Immulite as supplied by company kits(Diagnostic Products Corporation, Corporate Offices,5210 Pacific Concourse Drive, Losangeles, CA 90045-690 VCFL Sciences Journal ISSN 2331-9522
12. Gornall AG, Bardawill CS, David MM. Determination of serum proteins by means of the Biuret reaction. *J Biol Chem* 1949; 177:751-66.
13. Dumas BT, Watson WA, Biggs HG. Albumin standards and the measurement of serum albumin with bromocresol green. *Clin Chim Acta* 1971; 31:87-96.
14. Trinder P. Determination of blood glucose using an oxidase-peroxidase system with a non-carcinogenic chromogen. *J Clin Pathol* 1969 Mar; 22(2):158-61.
15. Selhub J, Jacques PF, Wilson PW, Rush D, Rosenberg IH. Vitamin status and intake as primary

- determinants of homocysteinemia in an elderly population. *JAMA* 1993; 270:2693-8.
16. Hao L, Ma J, Zhu J, Stampfer MJ, Tian Y, Willett WC et al. High prevalence of Hyperhomocysteinemia in Chinese Adults is associated with low Folate, Vitamin B12 and Vitamin B6 Status. *J Nutr* 2007; 137:407-13.
 17. Henecke JW, Rosen H, Suzuki LA, Chait A. The role of sulfur containing amino acids in superoxide production and modification of low density lipoprotein by arterial smooth muscle cells. *J Biol Chem* 1987; 262:10098-103.
 18. Olszewski AJ, McCully KS. Homocysteine metabolism and oxidative modification of proteins and lipids. *Free Radic Biol Med* 1993; 14:683-93.
 19. Stamler JS, Osborne JA, Jaraki O, Rabbani LE, Mullins M, Singel D, Loscalzo J. Adverse effect of homocysteine are modulated by endothelium-derived relaxing factor and related oxides of nitrogen. *J Clin Invest* 1993; 91:308-18.
 20. Misra HP. Generation of superoxide free radical during auto-oxidation of thiols. *J Biol Chem* 1974; 249:2151-5.
 21. Rowley DA, Halliwell B. Superoxide-dependent formation of hydroxyl radicals in the presence of thiol compounds. *FEBS Lett* 1982; 138:33-6.
 22. Zoric L. Parameters of oxidative stress in the lens aqueous humor and blood in patients with diabetes and senile cataracts. *Srpski arhiv za celokupno lekarstvo* 2003; 131(3-4):137-42.
 23. Malnar GA, Nemes V, Biro Z, Ludany A, Wagner Z, Wittmann I. Accumulation of hydroxyl free radical markers meta-ortho-tyrosine and DOPA in cataractous lenses is accompanied by a lower protein and phenylalanine content of the water-soluble phase. *Free Radical Research* 2005; 39:1359-66.
 24. Altomare E, Grattagliano I, Vendemaile G, Micelli-Ferrari T, Signorile A, Cardia L. Oxidative protein damage in human diabetic eye; evidnal participation. *Eur J Clin Invest (ENGLAND)* 1997; 27(2):141-7.
 25. Jain AK, Lim G, Langford M, Jain SK. Effects of high glucose levels on protein oxidation in cultured lens cells, and in crystalline and albumin solution and its inhibition by vitamin B6 and N-acetylcysteine: its possible relevance to cataract formation in diabetes. *Free Radic Biol Med* 2002; 33:1651-21.
-